

Clinical Practice Guideline: **Exercise Therapy for Treatment of Non-Specific Low Back Pain**

Date of Implementation: **September 18, 2008**

Product: **Specialty**

Table of Contents

GUIDELINES	1
DESCRIPTION/BACKGROUND	1
EVIDENCE REVIEW	2
PRACTITIONER SCOPE AND TRAINING	36
<i>References</i>	36

GUIDELINES

American Specialty Health – Specialty (ASH) considers exercise therapy medically necessary for treatment of patients with acute, sub-acute and chronic non-specific low back pain.

DESCRIPTION/BACKGROUND

The World Health Organization (WHO) defines low back pain (LBP) as discomfort experienced in the area between the bottom of the ribs and the buttocks. This pain can be classified as acute (lasting under 6 weeks), sub-acute (6-12 weeks), or chronic (over 12 weeks). Also, according to the WHO, LBP can be specific or non-specific. Specific LBP can be explained by an underlying disease (e.g., cancer), tissue damage (e.g., fracture), or may be referred from other organs (e.g., from kidney or aortic aneurysm). Non-specific means that the experience of pain cannot be confidently accounted for by another diagnosis such as an underlying disease, pathology or tissue damage. LBP is non-specific in about 90% of cases.

Chronic low back pain is a significant problem because of high health care utilization, rising health care costs, and perceived limitations of treatment effectiveness. Most patients with chronic low back pain have non-specific low back pain. Exercise therapy is one effective treatment option for chronic non-specific low back pain.

Exercise therapy represents a very diverse group of treatment approaches, which makes the discussion of “exercise therapy” as a whole difficult (Hayden et al., 2005).

Hayden et al. (2005) proposed the following specific characteristics of exercise: type, design, delivery, dose, and additional interventions.

Types of exercise therapy include muscle strengthening/stabilization/motor control exercises, stretching/flexibility, coordination/balance/proprioceptive exercises, and general fitness. Muscle strengthening typically involves repetitions of muscle contraction of specific muscle groups aimed to increase muscle strength and/or endurance (Abenhaim et al., 2000). Stretching/flexibility entail movements of one or more joints, intended to lengthen shortened muscles that can be static or dynamic in nature. Coordination and balance exercises involve training in specific movements aimed at improving proprioception and coordination of appropriate muscle groups (Johannsen et al., 1995; Kuukkanen & Malkia, 2000). Finally, general physical fitness routines typically include approaches involving the whole body (e.g., aerobic exercises) (Hayden et al., 2005).

Exercise therapy can also be categorized in terms of program design. Individualized programs are those tailored to the individual based on the history and physical examination. Partially individualized programs involve standard types of exercises, but at varied intensity and/or duration. Finally, standard exercise programs are ones in which all participants receive the same exercise program (Hayden et al., 2005).

Exercise programs can also be delivered in several ways: home, supervised home with follow up, group supervision, and individual supervision. Home exercise entails participants meeting initially with a therapist who provides them an exercise program to do at home, with no supervision or follow up. Home exercise with follow up involves the participants meeting initially with a therapist, doing the exercise program at home, and then having a follow up visit with the therapist at least every 6 weeks. In group supervised exercise, participants attend exercise sessions with 2 or more other individuals, under the guidance of a therapist. Finally, individually supervised exercise sessions entail individuals receiving one-on-one supervision while performing the prescribed exercise program (Hayden et al., 2005).

Dose or intensity (measured by the duration and number of treatment sessions) is also an important characteristic of exercise therapy (Hayden et al., 2005). Programs involving 20 or more hours of exercise are defined as high dose, and less than 20 hours of intervention time as low dose. Factors such as load, resistance, and frequency of repetitions (which can create a further categorization of strengthening exercise into strengthening vs. endurance) may also be important issues when addressing exercise dose (Manniche & Jordan, 1995; Jordan et al., 1998).

EVIDENCE REVIEW

Exercise is one of the few treatments for chronic low back pain with good literature support; however, the effect sizes reported have been small and the exact type of exercise that is most effective cannot be determined. In 2000, van Tulder et al. published a Cochrane review assessing exercise therapy for low back pain relative to pain relief, functional status, overall improvement and return to work. Thirty-nine randomized controlled trials (RCTs)

1 were included, and authors concluded that exercise therapy was not effective for acute low
 2 back pain but may be helpful for chronic low back pain. Since 2000, many new trials have
 3 been published, which precipitated the need for an updated review (Hayden et al., 2005).
 4 In this 2005 review, 61 RCTs were included in the analysis. These studies involved adult
 5 participants that could be categorized into acute, subacute, and chronic non-specific low
 6 back pain groups. Studies involving low back pain caused by a specific pathology or
 7 condition were excluded. Exercise therapy was defined as "a series of specific movements
 8 with the aim of training or developing the body by a routine practice or as physical training
 9 to promote good physical health." Studies included compared exercise therapy to a) no
 10 treatment or placebo treatment, b) other conservative treatment, or c) other exercise group.
 11 Outcomes of interest included self-reported pain intensity, condition-specific physical
 12 functioning, global improvement, and return to work/absenteeism. Both qualitative and
 13 quantitative rating systems were used to allow the most complete use of the available data.
 14 Of the total 61 RCTs, 43 trials (3,907 individuals) assessed chronic low back pain. Thirty-
 15 three exercise groups had non-exercise comparisons, and these trials provided strong
 16 evidence that exercise therapy is at least as effective as other conservative interventions.
 17 The evidence was conflicting as to whether exercise therapy was more effective than other
 18 treatments for chronic low back pain. It also appeared that exercise therapy is most
 19 effective when administered in a health care setting rather than as independent home
 20 exercises. In many of these trials, other conservative care was used in addition to exercise
 21 therapy; including behavioral and manual therapy, advice to stay active and education. As
 22 an aside, there is moderate effectiveness of graded-activity exercise programs for the
 23 subacute population. Only a small number of these studies were rated at high quality, which
 24 may have led to an overestimation of effect. Also, many of the studies lacked information
 25 to assess quality and clinical relevance. The most consistent outcome measure was for pain
 26 intensity, which limits the ability to discuss other outcome measures. Lastly, authors found
 27 potential publication bias, which also may have resulted in an overestimation of the
 28 effectiveness of exercise therapy in the chronic low back pain population. Authors also
 29 recommend that no further trials on the effectiveness of general exercise therapy for
 30 chronic low back pain should be initiated, but rather trials should focus on specific exercise
 31 intervention strategies in well-defined low back pain patient populations.

32
 33 Another review by Liddle et al. (2004) based on 16 RCTs of high to medium quality
 34 concluded that exercise as a primary intervention is an effective treatment for chronic low
 35 back pain, despite the wide variety of exercise programs offered. Positive results were
 36 maintained in 12 of the 16 trials, with supervision as a common feature. Again, authors felt
 37 studies did not explain exercise programs adequately and thus, no conclusions could be
 38 made regarding what type of exercise is most effective. The inclusion of exercise co-
 39 interventions introduced a confounding influence as well.

40
 41 To this end, a systematic review published in the Journal of Manipulative Physiological
 42 Therapeutics in 2007 attempted to determine the effect of unloaded movement facilitation

exercises on outcomes for people with non-specific chronic low back pain (NSCLBP) (Slade & Keating, 2007). In the previous systematic review reported by these authors, trunk strengthening was effective for improving function and reducing pain, compared to no exercise for patients with NSCLBP. Treatment effects increased with increasing exercise intensity and adding motivational strategies. Trunk strengthening exercises compared to aerobic training or the McKenzie approach showed no clear benefit (Slade & Keating, 2007). In their next review, 6 high quality RCTs were included. Participants were over 16 years of age with a current episode of low back pain lasting longer than 8 weeks (vs. the typical >12 weeks) with or without a history of low back surgery. Given this duration change, subjects could fall into the subacute category of low back pain rather than the chronic group. Authors stated that these parameters were used to capture the largest number of studies on exercise trials for chronic LBP that included the least number of participants likely to demonstrate a natural recovery process during the intervention time. They also defined low back pain as pain from below the scapulae to the buttock fold, with or without lower extremity radiation. Again, this varied from the previously described reviews. Interventions had to involve unloaded exercises that were likely to facilitate movement of the lumbar spine. If other interventions were involved, the unloaded exercise portion needed to be able to be partitioned out. Unloaded exercises basically referred to McKenzie exercises or yoga. Studies were excluded if they combined unloaded exercises with resistance exercises used to increase strengthening, spinal stabilization exercises or behavioral approaches and could not separate each component.

For NSCLBP without surgery, use of a McKenzie approach produced small effects for short and medium-term pain and short-term function compared to intensive trunk strengthening. There were no observable differences in outcomes when comparing the McKenzie approach to spinal stabilization exercises. When comparing yoga to trunk strengthening and aerobic training in subjects with NSCLBP without surgery, comparable effects were observed for short and medium-term outcomes. Compared to no exercise, yoga displayed a significantly large effect for medium term pain and function. Performing McKenzie exercises and yoga together compared with no exercise, significant, moderate effects on medium-term pain and function were noted in favor of the unloaded exercise. More specifically, within this review one of the RCTs published in the *Annals of Internal Medicine* (Sherman et al., 2005) attempted to determine whether yoga was more effective than conventional exercise or a self-care book for patients with chronic low back pain. One hundred one adults participated in a 12-week yoga program or conventional therapy program or just received a self-care book. They determined that yoga was more effective than a self-care book. The yoga group consistently reported superior outcomes compared with the exercise group, but these differences were not significant. Limitations included a relatively short follow up period (14 weeks), modest sample sizes, reliance on class instructors for intervention development and the inclusion of relatively highly educated and functional participants (Sherman et al., 2005). Authors stated that it would be virtually impossible to recreate these exercise programs, as minimal descriptions were reported.

1 Authors concluded that there is strong evidence that unloaded movement facilitation
 2 exercise compared to no exercise is effective for improving pain and function. However, it
 3 appears that when comparing unloaded exercise to other types of exercise, effects are
 4 comparable. It may be that for patients with NSCLBP, unloaded exercise is as effective as
 5 more vigorous forms of exercise that require more resources for relieving pain and
 6 increasing function.

7
 8 In another attempt to tease out what type of exercise is most beneficial, Kofotolis and Kellis
 9 (2006) studied the effects of two 4-week Proprioceptive Neuromuscular Facilitation (PNF)
 10 programs on muscle endurance, flexibility, and functional performance in women with
 11 chronic low back pain. Unfortunately, these programs were only compared to one another
 12 and not with another type of exercise program. Results demonstrated that both static and
 13 dynamic PNF programs were effective in improving short-term trunk muscle endurance
 14 and trunk mobility in people with chronic low back pain. Another RCT by Koumantakis et
 15 al. (2005) compared a general trunk muscle endurance exercise program enhanced with
 16 specific muscle stabilization exercises with a general exercise approach only. Fifty-five
 17 patients with recurrent LBP were randomized to the two groups. Both groups received an
 18 8-week intervention and written instructions. Results indicated that both the general
 19 exercise program and the enhanced exercise program provided benefits for patients with
 20 recurrent LBP. It appears to be the presence of physical exercise alone, rather than the
 21 specific exercise type that is the factor in patient improvement in those with chronic LBP.

22
 23 Another RCT compared general exercise, motor control exercise, and spinal manipulation
 24 therapy for chronic low back pain (Ferreira et al., 2007). Each group received 8 weeks of
 25 treatment. General exercise included strengthening, stretching and aerobic exercise, motor
 26 control exercise included retraining of specific trunk musculature using ultrasound and
 27 feedback, and spinal manipulation therapy involved both mobilization and manipulation.
 28 At 8 weeks the motor control group and manipulation group had slightly better outcomes
 29 than the general exercise group. At 6 and 12 months, these differences diminished, and
 30 similar outcomes were reported. It appears that motor control exercise has better short-term
 31 outcomes, while all three are equivalent over the medium and long-term with regards to
 32 perceived effectiveness and function (Ferreira et al., 2007). Costa et al. (2009) completed
 33 a randomized, placebo-controlled trial with subjects complaining of non-specific low back
 34 with or without leg pain for at least 3 months. Subjects were instructed in specific deep
 35 trunk muscle isolation exercise training which consisted of 12 individually supervised half-
 36 hour sessions over an 8-week period. The placebo group received 20 minutes of detuned
 37 short-wave diathermy and 5 minutes of detuned ultrasound for 12 sessions over an 8-week
 38 period. Outcomes were measured at 2, 6, and 12 months. This study found that motor
 39 control exercise produced short-term improvements in global impression of recovery and
 40 activity, but not pain, for people with chronic low back pain. Most of the effects observed
 41 in the short term were maintained at the six 6- and 12-month follow-ups (Costa, 2009).

In another review on use of the McKenzie method for chronic LBP by May and Donelson (2008), they suggest that the McKenzie method plays an important role in the classification of subgroups with different needs treatment-wise. It appears that as an intervention, this method produces more positive short-term outcomes than non-specific guideline-based care and equal or slightly better outcomes than stabilization or strengthening routines for patients with chronic LBP (May & Donelson, 2008). Another review on lumbar extension strengthening exercises for chronic LBP by Mayer et al. (2008) suggests that it is an effective intervention over no treatment or most passive modalities, whether used in isolation or as a co-intervention. These subjects report improved pain, disability, and other reported outcomes in the short term. Over the long term, this review suggests that some of the disability and pain benefits are lost. There also appears to be no clear benefit to lumbar extensor strengthening exercises over other exercise programs regarding improvements in pain, disability, strength, and endurance. Standaert et al. (2008) reported that lumbar stabilization exercises for chronic low back pain are effective at improving pain and function in a variety of patients with chronic LBP based on moderate evidence. Moderate evidence also suggests that lumbar stabilization exercises are no more effective than manual therapy. Strong evidence does exist that lumbar stabilization exercises are no more effective than a less specific, general exercise program (Standaert et al., 2008).

There are a few well-designed studies that demonstrate the effectiveness of activity or therapeutic exercise when used in conjunction with other manual interventions in the management of spinal pain. Research has demonstrated the benefit of matching sub-categories of patients to specific interventions. One of the interventions that has shown marked success in the treatment of LBP is manipulation combined with strengthening exercise. Flynn et al. (2002) reported 5 clinical predictors for success with spinal manipulation (Symptom duration <16 days, No symptoms distal to the knee, Fear Avoidance Belief Questionnaire Work Subscale <19, Hip IR >35 degrees, Positive lumbar spring test on at least one lumbar segment). Flynn found a Positive Likelihood Ratio (+LR) of 24 which provides a 95% chance of decreasing disability by >50% within the first two (2) treatments using manipulation. Childs et al. (2004) validated this rule in a multi-center trial and also determined the number needed to treat with thrust manipulation combined with exercise to prevent one patient from experiencing a worsening of disability was only ten. Childs et al. (2006) later reported that patients that met the clinical prediction rules above were 8 times more likely to experience an increase in disability within one week if they were not treated with a combined thrust manipulation/exercise intervention. This Clinical Prediction Rule has also been validated in the Primary Care setting by Fritz et al. (2005). The authors determined a +LR for success with thrust manipulation of 7.2 with the following two factors present: symptoms less than 16 days duration and no symptoms distal to the knee.

The literature demonstrates that an Extension Oriented Treatment Approach (EOTA) is beneficial in patients who demonstrate a directional preference (DP) of symptom

centralization with extension postures/exercises (Browder et al., 2007). The average duration of the patients' symptoms was 3 months. The authors compared an EOTA with strengthening exercises and reported the EOTA group demonstrated greater improvements in disability and pain at 1 week follow-up and greater improvement in disability at 4 weeks and 6-month follow-ups as well. The EOTA was provided over the course of 8 sessions (twice a week for 4 weeks) and included the following interventions:

1. Extension-oriented exercises (sustained and repeated) in prone and standing;
2. Posterior to Anterior (PA) lumbar mobilizations, grade I to IV, 10 to 20 oscillations;
3. Home exercise prescription (prone press-up) x10 repetitions every 2 to 3 waking hours (may substitute standing extension exercises).

The Orthopaedic Section of the American Physical Therapy Association (APTA) has an ongoing effort to create evidence-based practice guidelines for orthopaedic physical therapy management of patients with musculoskeletal impairments described in the World Health Organization's International Classification of Functioning, Disability, and Health (ICF). In 2012, Delitto et al. authored guidelines for low back pain. The purpose of these low back pain clinical practice guidelines was to describe the peer-reviewed literature and make recommendations related to (1) treatment matched to low back pain subgroup responder categories, (2) treatments that have evidence to prevent recurrence of low back pain, and (3) treatments that have evidence to influence the progression from acute to chronic low back pain and disability. Authors presented "A" level recommendations for treatment of low back pain which included manual therapy, trunk coordination, strengthening and endurance exercises, centralization and directional preference exercises and progressive endurance exercises and fitness activities. Research has determined thrust manipulation is effective in a subgroup of patients as part of a multi-component program including exercise. Lumbar coordination, strengthening and endurance exercises are a common treatment intervention for back pain. They are also referred to in the literature as motor control exercises, transversus abdominis training, lumbar multifidus training and dynamic lumbar stabilization exercises. Delitto et al. (2012) summarized the available literature indicating that clinicians should consider these exercises to reduce low back pain and disability in patients with subacute and chronic low back pain with movement dysfunction and in patients post microdiscectomy. Much of the research demonstrates that these exercises are effective but may be no more effective than a general exercise program. Centralization exercises appear to be beneficial for patients with acute low back pain with referred lower extremity pain. Clinicians should consider using repeated movements and exercises to promote centralization through reduction of lower extremity pain. Also, repeated movements in a specific direction, as noted by treatment response, should be utilized to reduce symptoms and improve mobility in all phases of low back pain. Lastly, progressive endurance exercises and fitness activities are endorsed by most current low back pain guidelines with moderate to high levels of evidence. Aerobic conditioning has

1 been hypothesized to reduce pain perception and improving function in patients with
2 chronic low back pain and other generalized pain.

3
4 A meta-analysis by Wang et al. (2012) concluded that core stability exercises are more
5 effective in decreasing pain and may improve physical function in patients with chronic
6 low back in the short-term relative to general exercise. However, over the long term, no
7 significant differences were noted. In 2013, Brumitt et al. (2013a) provided clinical
8 recommendations using the SORT (Strength of Recommendation Taxonomy) method.
9 They concluded that a therapeutic intervention program consisting of motor control
10 exercises OR general back strengthening exercises may be beneficial for patients with low
11 back pain lasting longer than 6 weeks. However, given the SORT evidence rating of 'B'
12 indicates that the evidence is inconsistent or of limited quality. Brumitt et al. (2013b)
13 published another paper analyzing randomized controlled trials that assessed the effects of
14 a motor control exercise approach, a general exercise approach, or both for patients with
15 low back pain that were published in scientific peer-reviewed journals. Fifteen studies were
16 identified (8, motor control exercise approach without general exercise comparison; 7,
17 general exercise approach with or without motor control exercise approach comparison).
18 Authors stated that current evidence suggests that exercise interventions may be effective
19 at reducing pain or disability in patients with low back pain, but it may not be necessary to
20 prescribe exercises purported to restore motor control of specific muscles. A systematic
21 review by Stuber et al. (2014) reviewed the effectiveness of core stability exercises for low
22 back pain in athletes. They concluded that given the low quantity and quality of available
23 literature, no strong conclusions could be formulated.

24
25 Lehtola et al. (2016) conducted a randomized controlled trial (RCT) to compare the effects
26 of general exercise versus specific movement control exercise (SMCE) on disability and
27 function in patients with MCI within the recurrent sub-acute LBP group. Subjects attended
28 5 sessions of either specific or general exercises. Both groups also received a short
29 application of manual therapy. The primary outcome was disability, assessed by the
30 Roland-Morris Disability Questionnaire (RMDQ). The measurements were taken at
31 baseline, immediately after the three months intervention and at twelve-month follow-up.
32 Measurements of 61 patients (SMCE $n = 30$ and general exercise $n = 31$) were completed
33 at 12 months. Patients in both groups reported significantly less disability at 12 months
34 follow up, with the SMCE group showing statistically significantly superior improvement.
35 However, the result did not reach the clinically significant three-point difference. There
36 was no statistical difference between the groups measured with Oswestry Disability Index
37 (ODI). Authors concluded for subjects with non-specific recurrent sub-acute LBP and MCI
38 an intervention consisting of SMCE and manual therapy combined may be superior to
39 general exercise combined with manual therapy. Saragiotto et al. (2016) authored a
40 Cochrane Review on motor control exercise for chronic non-specific low back pain.
41 (CNSLBP). As noted in the previous literature, exercise is a modestly effective treatment
42 for chronic LBP and current evidence suggests that no single form of exercise is superior

to another. Authors report that among the most commonly used exercise interventions are motor control exercise (MCE). To clarify, MCE intervention focuses on the activation of the deep trunk muscles and targets the restoration of control and co-ordination of these muscles, progressing to more complex and functional tasks integrating the activation of deep and global trunk muscles. Authors included trials comparing MCE with no treatment, another treatment or that added MCE as a supplement to other interventions. Primary outcomes were pain intensity and disability. They also considered function, quality of life, return to work or recurrence as secondary outcomes. They considered the following time points: short-term (less than 3 months after randomization); intermediate (at least three months but less than 12 months after randomization); and long-term (12 months or more after randomization) follow-up. 29 trials ($n = 2,431$) were included in this review. The study sample sizes ranged from 20 to 323 participants. Results demonstrate that there is low to high quality evidence that MCE is not clinically more effective than other exercises for all follow-up periods and outcomes tested. When compared to minimal intervention, there is low to moderate quality evidence that MCE is effective for improving pain at short, intermediate and long-term follow-up with medium effect sizes. There was also a clinically important difference for the outcomes function and global impression of recovery compared with minimal intervention. There was moderate to high quality evidence that there is no clinically important difference between MCE and manual therapy for all follow-up periods and outcomes tested. Finally, there was very low to low quality evidence that MCE is clinically more effective than exercise and electrophysical agents (EPA) for pain, disability, global impression of recovery and quality of life with medium to large effect sizes. Minor or no adverse events were reported in the included trials. Authors conclude that given the evidence that MCE is not superior to other forms of exercise, the choice of exercise for chronic LBP should probably depend on patient or therapist preferences, therapist training, costs, and safety.

Macedo et al. (2016) completed a Cochrane Review on the effectiveness of motor control exercise for acute non-specific low back pain. They only included RCTs examining the effectiveness of MCE for patients with acute non-specific LBP. Authors considered trials comparing MCE versus no treatment, versus another type of treatment or added as a supplement to other interventions. Primary outcomes were pain intensity and disability. Secondary outcomes were function, quality of life and recurrence. Authors considered the following follow-up intervals: short term (less than three months after randomization); intermediate term (at least three months but within 12 months after randomization); and long term (12 months or longer after randomization). Only 3 trials were included with study samples ranging from 33 to 123 participants. Evidence of very low to moderate quality indicates that MCE showed no benefit over spinal manipulative therapy, other forms of exercise or medical treatment in decreasing pain and disability among patients with acute and subacute low back pain. Whether MCE can prevent recurrences of LBP remains uncertain and no firm conclusions can be drawn regarding the effectiveness of MCE for acute LBP.

Pilates was also examined in a Cochrane Review as a treatment for non-specific LBP (Yamato et al., 2015). They included RCTs that examined the effectiveness of Pilates intervention in adults with acute, subacute, or chronic non-specific low back pain. The primary outcomes considered were pain, disability, global impression of recovery and quality of life. A total of 6 trials compared Pilates to minimal intervention. They did not find any high-quality evidence for any of the treatment comparisons, outcomes or follow-up periods investigated. However, there is low to moderate quality evidence that Pilates is more effective than minimal intervention for pain and disability. When Pilates was compared with other exercises, the authors found a small effect for function at intermediate-term follow-up. Thus, while there is some evidence for the effectiveness of Pilates for low back pain, there is no conclusive evidence that it is superior to other forms of exercises. The decision to use Pilates for low back pain may be based on the patient's or care provider's preferences, and costs.

A systematic review and meta-analysis by Carey and Freburger (2016) assessed research into the value of exercise as a way to treat and prevent LBP. The study found that exercise alone was linked to a 35% reduction in risk, while the combination of exercise and education was associated with a 45% risk reduction for up to one year. The use of exercise was also found to result in a 78% reduction in sick leave for LBP. Authors found that while education helped to further reduce that risk when combined with exercise, education alone doesn't seem to have much effect, according to authors. They also suggest that for exercise to remain protective against future LBP, it needs to be ongoing.

The Agency for Healthcare Research and Quality (AHRQ) published a Comparative Effectiveness Review in 2016 on noninvasive treatments for LBP. They summarized the research on exercise and LBP with the following key points:

1. For acute LBP, a systematic review found no differences between exercise therapy versus no exercise in pain or function; for subacute LBP, there were no differences in pain or function. Three other trials for acute to subacute LBP found inconsistent results of exercise vs. usual care to improve pain and function.
2. For chronic LBP, a systematic review found exercise was associated with greater pain relief versus no exercise and a more recent review using more restrictive criteria and additional trials were consistent with these earlier findings.
3. More specifically, for chronic LBP, a review found motor control exercise was associated with lower pain scores and better function in the short, intermediate and long term vs. minimal intervention. Another systematic review found MCE associated with lower pain intensity at the short term and intermediate term versus general exercise. No significant findings were noted in the long term. Better function was noted with MCE in the short and long term.
4. For radicular LBP, three trials not included in any systematic reviews found effects that favored exercise versus usual care or no exercise in pain and function, though effect sizes were small.

5. For comparisons of different exercise types, there were no clear differences for patients with acute or chronic LBP.
6. Adverse events were not often reported and if they were, typically muscle soreness and increased pain were reported. No serious harms were reported.

According to Qaseem et al. (2017), moderate-quality evidence showed that exercise therapy resulted in small improvements in pain and function. Specific components associated with greater effects on pain included individually designed programs, supervised home exercise, and group exercise; regimens that included stretching and strength training were most effective. In a systematic review, Vanti et al. (2019) found that pain, disability, quality of life and fear-avoidance similarly improve by walking or exercise in chronic low back pain. Walking may be considered as an alternative to other physical activity. Adding walking to exercise does not induce greater improvement in the short-term. Walking may be a less-expensive alternative to physical exercise in chronic low back pain. Wewege et al. (2018) compared progressive aerobic training (PAT) to progressive resistance training (PRT) for pain, disability, and quality of life (QoL) in people with chronic non-specific low back pain (CNSLBP). Six studies were included, comprising 333 participants. Exercise significantly reduced pain intensity although neither mode proved superior. PRT significantly improved the Short Form Health Survey-Mental Component Score. Authors concluded that PAT and PRT decreased pain intensity in individuals with CNSLBP although neither mode was superior. Resistance exercise improved psychological wellbeing. High-quality RCTs comparing PAT, PRT, and PAT + PRT, are required. Shi et al. (2018) analyzed all evidence available in the literature about effectiveness of the aquatic exercise. Eight trials involving 331 patients were included in the meta-analysis, and the results showed a relief of and physical function after aquatic exercise. However, there was no significant effectiveness with regard to general mental health in aquatic group. Authors concluded that aquatic exercise can statistically significantly reduce pain and increase physical function in patients with low back pain. Shiri et al. (2018) assessed the effect of exercise in population-based interventions to prevent low back pain (LBP) and associated disability. Thirteen randomized controlled trials (RCTs) and 3 nonrandomized controlled trials (NRCTs) qualified for the meta-analysis. Exercise alone reduced the risk of LBP by 33% and exercise combined with education reduced it by 27%. The severity of LBP and disability from LBP were also lower in exercise groups than in control groups. Authors concluded that exercise reduces the risk of LBP and associated disability, and a combination of strengthening with either stretching or aerobic exercises performed 2-3 times per week can reasonably be recommended for prevention of LBP in the general population. Suh et al. (2019) compared the efficiency between 2 exercises: the individualized graded lumbar stabilization exercise (IGLSE) and walking exercise (WE). A randomized controlled trial was conducted in 48 participants with chronic LBP. After screening, participants were randomized to 1 of 4 groups: flexibility exercise (FE), WE, stabilization exercise (SE), and stabilization with WE (SWE) groups. Participants underwent each exercise for 6 weeks. The primary outcome was visual analog scale (VAS)

of LBP during rest and physical activity. Secondary outcomes were as follows: VAS of radiating pain measured during rest and physical activity; frequency of medication use (number of times/day); Oswestry disability index; Beck Depression Inventory; endurance of specific posture; and strength of lumbar extensor muscles. The present study showed that lumbar SE and WE significantly improved chronic LBP. Both WE and stabilization with WE significantly improved muscular endurance of back muscles. Moreover, walking and SEs also improved the core stability. It is also worth noting that patients in the WE and SE groups were much more compliant than those in the other exercise groups. This study suggests that lumbar SE and WE should be recommended to patients with chronic LBP because they help not only to relieve back pain but also to prevent chronic back pain through the improvement of muscle endurance.

Many clinical practice guidelines recommend similar approaches for the assessment and management of low back pain. Recommendations include use of a biopsychosocial framework to guide management with initial non-pharmacological treatment, including education that supports self-management and resumption of normal activities and exercise, and psychological programs for those with persistent symptoms (Foster et al., 2018). Jones et al. (2020) discusses the use of pain education with therapeutic exercise to address the psychosocial aspects that are associated with chronic low back pain. Pain education is the umbrella term utilized to encompass any type of education to the patient about their chronic pain. Therapeutic exercise in combination with pain education may allow for more well-rounded and effective treatment for patients with chronic nonspecific low back pain (NS-LBP). They summarized key findings: A thorough literature review yielded 8 studies potentially relevant to the clinical question, and 3 studies that met the inclusion criteria were included. The 3 studies included reports that exercise therapy reduced symptoms. Two of the 3 included studies support the claim that exercise therapy reduces the symptoms of chronic NS-LBP when combined with pain education, whereas one study found no difference between pain education with therapeutic exercise. Authors concluded that there is moderate evidence to support the use of pain education along with therapeutic exercise when attempting to reduce symptoms of pain and disability in patients with chronic NS-LBP. Educational interventions should be created to educate patients about the foundation of pain, and pain education should be implemented as a part of the clinician's strategy for the rehabilitation of patients with chronic NS-LBP.

Owen et al. (2020) examined the effectiveness of specific modes of exercise training in non-specific chronic low back pain (NSCLBP). They included exercise training randomized controlled/clinical trials in adults with NSCLBP. Among 9,543 records, 89 studies (patients=5,578) were eligible for qualitative synthesis and 70 (pain), 63 (physical function), 16 (mental health) and 4 (trunk muscle strength) for Network Meta-analysis (NMA). The NMA consistency model revealed that the following exercise training modalities had the highest probability of being best when compared with true control: Pilates for pain, resistance and stabilization/motor control for physical function, and

1 resistance and aerobic for mental health. Stretching and McKenzie exercise effect sizes did
 2 not differ to true control for pain or function. NMA was not possible for trunk muscle
 3 endurance or analgesic medication. Authors concluded there is low quality evidence that
 4 Pilates, stabilization/motor control, resistance training and aerobic exercise training are the
 5 most effective treatments, pending outcome of interest, for adults with NSCLBP. Exercise
 6 training may also be more effective than therapist hands-on treatment. Heterogeneity
 7 among studies and the fact that there are few studies with low risk of bias are both
 8 limitations. Hayden et al. (2020) sought to determine which individuals might benefit the
 9 most from exercise for their low back pain. For studies included in this analysis, compared
 10 with no treatment/usual care, exercise therapy on average reduced pain, a result compatible
 11 with a clinically important 20% smallest worthwhile effect. Exercise therapy reduced
 12 functional limitations with a clinically important 23% improvement at short-term follow-
 13 up. Not having heavy physical demands at work and medication use for low back pain were
 14 potential treatment effect modifiers that were associated with superior exercise outcomes
 15 relative to non-exercise comparisons. Lower body mass index was also associated with
 16 better outcomes in exercise compared with no treatment/usual care. This study was limited
 17 by inconsistent availability and measurement of participant characteristics.

18
 19 Zhu et al. (2020) compared the effects of yoga for patients with chronic low back pain on
 20 pain, disability, quality of life with non-exercise (e.g., usual care, education), physical
 21 therapy exercise. A total of 18 randomized controlled trials were included in this meta-
 22 analysis. Yoga could significantly reduce pain at 4 to 8 weeks, 3 months, 6 to 7 months,
 23 and was not significant in 12 months compared with non-exercise. Yoga was better than
 24 non-exercise on disability at 4 to 8 weeks, 3 months, 6 months, 12 months. There was no
 25 significant difference on pain, disability compared with physical therapy exercise group.
 26 Furthermore, it suggested that there was a non-significant difference on physical and
 27 mental quality of life between yoga and any other interventions. Authors concluded that
 28 yoga might decrease pain from short term to intermediate term and improve functional
 29 disability status from short term to long term compared with non-exercise (e.g., usual care,
 30 education). Yoga had the same effect on pain and disability as any other exercise or
 31 physical therapy. Yoga might not improve the physical and mental quality of life based on
 32 the result of merging the 36 item short form health survey (SF-36) and the 12 item short
 33 form health survey (SF-12) data.

34
 35 Karlsson et al. (2020) assessed the overall certainty of evidence for the effects of exercise
 36 therapy, compared with other interventions, on pain, disability, recurrence, and adverse
 37 effects in adult patients with acute low back pain within a systematic review. Twenty-four
 38 reviews were included, in which 21 randomized controlled trials ($n = 2,685$) presented data
 39 for an acute population, related to 69 comparisons. Overlap was high, 76%, with a
 40 corrected covered area of 0.14. Methodological quality varied from low to high. Exercise
 41 therapy was categorized into general exercise therapy, stabilization exercise, and
 42 McKenzie therapy. No important difference in pain or disability was evident when exercise

therapy was compared with sham ultrasound, nor for the comparators of usual care, spinal manipulative therapy, advice to stay active, and educational booklet. Neither McKenzie therapy nor stabilization exercise yielded any important difference in effects compared with other types of exercise therapy. Certainty of evidence varied from very low to moderate. Authors concluded that these findings suggest very low to moderate certainty of evidence that exercise therapy may result in little or no important difference in pain or disability, compared with other interventions, in adult patients with acute low back pain.

Skelly et al. (2020) updated the evidence from their 2018 report assessing persistent improvement in outcomes following completion of therapy for noninvasive nonpharmacological treatment for selected chronic pain conditions. They included 233 RCTs (31 new to this update). Many were small ($N < 70$), and evidence beyond 12 months after treatment completion was sparse. The most common comparison was with usual care. Evidence on harms was limited, with no evidence suggesting increased risk for serious treatment-related harms for any intervention. Effect sizes were generally small for function and pain. For chronic low back pain, function improved over short and/or intermediate term for exercise (SOE moderate at short term for exercise). Improvements in pain at short term were seen for exercise (SOE: low). At intermediate term, exercise (SOE: low) were associated with improved pain. Compared with exercise, multidisciplinary rehabilitation improved both function and pain at short and intermediate terms (small effects, SOE: moderate.)

Hayden et al. (2021a) assessed the impact of exercise treatment on pain and functional limitations in adults with chronic non-specific low back pain compared to no treatment, usual care, placebo and other conservative treatments in a Cochrane review. The review includes data for trials identified in searches up to 27 April 2018. Authors included randomized controlled trials that assessed exercise treatment compared to no treatment, usual care, placebo or other conservative treatment on the outcomes of pain or functional limitations for a population of adult participants with chronic non-specific low back pain of more than 12 weeks' duration. They included 249 trials of exercise treatment, including studies conducted in Europe (122 studies), Asia (38 studies), North America (33 studies), and the Middle East (24 studies). Sixty-one per cent of studies (151 trials) examined the effectiveness of two or more different types of exercise treatment, and 57% (142 trials) compared exercise treatment to a non-exercise comparison treatment. Study participants had a mean age of 43.7 years and, on average, 59% of study populations were female. Most of the trials were judged to be at risk of bias, including 79% at risk of performance bias due to difficulty blinding exercise treatments. Authors found moderate-certainty evidence that exercise treatment is more effective for treatment of chronic low back pain compared to no treatment, usual care, or placebo comparisons for pain outcomes at earliest follow-up, a clinically important difference. Certainty of evidence was downgraded mainly due to heterogeneity. For the same comparison, there was moderate-certainty evidence for functional limitations outcomes; this finding did not meet the prespecified threshold for

minimal clinically important difference. Certainty of evidence was downgraded mainly due to some evidence of publication bias. Compared to all other investigated conservative treatments, exercise treatment was found to have improved pain and functional limitations outcomes. These effects did not meet the prespecified threshold for clinically important difference. Subgroup analysis of pain outcomes suggested that exercise treatment is probably more effective than education alone or non-exercise physical therapy, but with no differences observed for manual therapy. In studies that reported adverse effects (86 studies), one or more adverse effects were reported in 37 of 112 exercise groups (33%) and 12 of 42 comparison groups (29%). Twelve included studies reported measuring adverse effects in a systematic way, with a median of 0.14 per participant in the exercise groups (mostly minor harms, e.g., muscle soreness), and 0.12 in comparison groups. Authors concluded that moderate-certainty evidence exists that exercise is probably effective for treatment of chronic low back pain compared to no treatment, usual care or placebo for pain. The observed treatment effect for the exercise compared to no treatment, usual care or placebo comparisons is small for functional limitations, not meeting the threshold for minimal clinically important difference. They also found exercise to have improved pain (low-certainty evidence) and functional limitations outcomes (moderate-certainty evidence) compared to other conservative treatments; however, these effects were small and not clinically important when considering all comparisons together. Subgroup analysis suggested that exercise treatment is probably more effective than advice or education alone, or electrotherapy, but with no differences observed for manual therapy treatments. Hayden et al. (2021b) wanted to investigate what the effects of specific types of exercise treatments on pain intensity and functional limitation outcomes for adults with chronic low back pain are in a systematic review with network meta-analysis of randomized controlled trials. Exercise treatments prescribed or planned by a health professional that involved conducting specific activities, postures and/or movements with a goal to improve low back pain outcomes were included in the review. Outcome measures included pain intensity (e.g., visual analogue scale or numerical rating scale) and back-related functional limitations (e.g., Roland Morris Disability Questionnaire or Oswestry Disability Index), each standardized to range from 0 to 100. This review included 217 randomized controlled trials with 20,969 participants and 507 treatment groups. Most exercise types were more effective than minimal treatment for pain and functional limitation outcomes. Network meta-analysis results were compatible with moderate to clinically important treatment effects for Pilates, McKenzie therapy, and functional restoration (pain only) and flexibility exercises (function only) compared with minimal treatment, other effective treatments, and other exercise types. This review found evidence that Pilates, McKenzie therapy and functional restoration were more effective than other types of exercise treatment for reducing pain intensity and functional limitations. Nevertheless, people with chronic low back pain should be encouraged to perform the exercise that they enjoy to promote adherence.

Thorton et al. (2021) summarized the evidence for non-pharmacological management of low back pain (LBP) in athletes, a common problem in sport that can negatively impact performance and contribute to early retirement. Among 1,629 references, 14 randomized controlled trials (RCTs) involving 541 athletes were included. The trials had biases across multiple domains including performance, attrition, and reporting. Treatments included exercise, biomechanical modifications, and manual therapy. There were no trials evaluating the efficacy of surgery or injections. Exercise was the most frequently investigated treatment; no RTS data were reported for any exercise intervention. There was a reduction in pain and disability reported after all treatments. Authors concluded that while several treatments for LBP in athletes improved pain and function, it was unclear what the most effective treatments were, and for whom. Exercise approaches generally reduced pain and improved function in athletes with LBP, but the effect on RTS is unknown. No conclusions regarding the value of manual therapy (massage, spinal manipulation) or biomechanical modifications alone could be drawn because of insufficient evidence. High-quality RCTs are urgently needed to determine the effect of commonly used interventions in treating LBP in athletes. Quentin et al. (2021) conducted a systemic review and meta-analysis on the effects of home-based exercise on pain and functional limitation in LBP. They included 33 studies and 9,588 patients. They found that pain intensity decreased in the exclusive home exercise group in the group which conducted exercise both at-home and at another setting. Similarly, functional limitation also decreased in both groups. Relaxation and postural exercise seemed to be ineffective in decreasing pain intensity, whereas trunk, pelvic or leg stretching decreased pain intensity. Yoga improved functional limitation.

Supervised training was the most effective method to improve pain intensity. Insufficient data precluded robust conclusions around the duration and frequency of the sessions and program. Authors concluded that home-based exercise training improved pain intensity and functional limitation parameters in LBP. Van Dillen et al. (2021) sought to determine whether an exercise-based treatment of person-specific motor skill training (MST) in performance of functional activities is more effective in improving function than strength and flexibility exercise (SFE) immediately, 6 months, and 12 months following treatment. The effect of booster treatments 6 months following treatment also was examined. A total of 154 people with at least 12 months of chronic, nonspecific LBP, aged 18 to 60 years, with modified Oswestry Disability Questionnaire (MODQ) score of at least 20% were randomized to either MST or SFE. Data were analyzed between December 1, 2017, and October 6, 2020. Participants received 6 weekly 1-hour sessions of MST in functional activity performance or SFE of the trunk and lower limbs. Half of the participants in each group received up to 3 booster treatments 6 months following treatment. A total of 149 participants (91 women; mean [SD] age, 42.5 [11.7] years) received some treatment and were included in the intention-to-treat analysis. Following treatment, MODQ scores were lower for MST than SFE by 7.9 (95% CI, 4.7 to 11.0; $P < .001$). During the follow-up phase, the MST group maintained lower MODQ scores than the SFE group, 5.6 lower at 6

months (95% CI, 2.1 to 9.1) and 5.7 lower at 12 months (95% CI, 2.2 to 9.1). Booster sessions did not change MODQ scores in either treatment. Authors concluded that people with chronic LBP who received MST had greater short-term and long-term improvements in function than those who received SFE. Person-specific MST in functional activities limited owing to LBP should be considered in the treatment of people with chronic LBP.

According to Chou (2021), low back pain is a common problem that is the leading cause of disability and is associated with high costs. Evaluation focuses on identification of risk factors indicating a serious underlying condition and increased risk for persistent disabling symptoms in order to guide selective use of diagnostic testing (including imaging) and treatments. Nonpharmacologic therapies, including exercise and psychosocial management, are preferred for most patients with low back pain and may be supplemented with adjunctive drug therapies. Surgery and interventional procedures are options in a minority of patients who do not respond to standard treatments. Hlaing et al. (2021) compared the effects of two different exercise regimes, Core stabilization exercises (CSE) and strengthening exercise (STE), on proprioception, balance, muscle thickness and pain-related outcomes in patients with subacute non-specific low back pain (NSLBP). Thirty-six subacute NSLBP patients, [mean age, 34.78 ± 9.07 years; BMI, 24.03 ± 3.20 Kg/m²; and duration of current pain, 8.22 ± 1.61 weeks], were included in this study. They were randomly allocated into either CSE ($n = 18$) or STE groups ($n = 18$). Exercise training was given for 30 min, three times per week, for up to 4 weeks. Proprioception, standing balance, muscle thickness of transversus abdominis (TrA) and lumbar multifidus (LM), and pain-related outcomes, comprising pain, functional disability and fear of movement, were assessed at baseline and after 4 weeks of intervention. The CSE group demonstrated significantly more improvement than the STE group after 4 weeks of intervention. Improvements were in: proprioception, balance: single leg standing with eyes open and eyes closed on both stable and unstable surfaces, and percentage change of muscle thickness of TrA and LM. Although both exercise groups gained relief from pain, the CSE group demonstrated greater reduction of functional disability and fear of movement. There were no significant adverse effects in either type of exercise program. Authors concluded that despite both core stabilization and strengthening exercises reducing pain, core stabilization exercise is superior to strengthening exercise. It is effective in improving proprioception, balance, and percentage change of muscle thickness of TrA and LM, and reducing functional disability and fear of movement in patients with subacute NSLBP.

Rathnayake et al. (2021) systematically reviewed the evidence for the effect of self-management interventions (SMIs) with an exercise component added, on pain and disability in people with CLBP. Authors concluded that there is low-quality evidence that SMIs with exercises added have moderately positive effects on pain and disability in patients with CLBP compared to control interventions involving usual care, typically consisting of access to medication, exercise, advice, education, and manual therapy.

Drummond et al. (2021) assessed the effectiveness of sling exercise therapy (SET) in individuals with chronic low back pain (LBP). The search identified 1,204 studies, with 12 studies meeting the inclusion criteria. Meta-analysis comparing SET with general exercise revealed a nonsignificant effect for pain. Meta-analysis comparing SET with motor control training/lumbar stabilization revealed a significant effect favoring SET for pain and disability. Meta-analysis comparing SET with no treatment revealed a significant effect favoring SET for pain. Meta-analysis comparing SET plus modalities with modalities revealed a significant effect favoring the SET plus modalities group for pain and a nonsignificant effect for disability. Sling exercise therapy was more effective than all comparisons for various muscle attributes. The overall level of evidence ranged from very low to moderate. Sling exercise therapy is effective in reducing pain, disability, and improving core muscle activation, strength, thickness, and onset in patients with chronic LBP. Because SET demonstrated comparable outcomes with common active interventions, it provides an opportunity to implement pain-free exercises based on the patient's initial functional level early in the plan of care.

Ferreira et al. (2021) assessed whether an exercise and education program was more effective than an education booklet for preventing recurrence of low back pain (LBP). Participants aged 18 years or older who had recovered from an episode of LBP within the previous week were recruited from primary care practices and the community. Participants were randomized to receive either 12 weeks of exercise and education (8 supervised exercise sessions and 3 one-on-one sessions) or a control (education booklet). The primary outcome was time to recurrence of LBP during the 1-year follow-up. Times to recurrence of LBP leading to activity limitation, care seeking, and work absence were secondary outcomes. Data were analyzed with Cox regression using intention-to-treat principles. The same size was 111 (exercise and education, $n = 57$; educational booklet, $n = 54$). At the end of the study period, data completeness was 84.2%. Thirty-six (63%) participants in the exercise and education group and 31 (57%) participants in the control group had a recurrence of LBP. There was no statistically significant difference in time to recurrence of pain between groups (hazard ratio = 1.09; 95% confidence interval: 0.7, 1.8). There was no statistically significant effect for any of the secondary outcomes. Authors concluded that among people recently recovered from LBP, exercise and education may not meaningfully reduce risk of recurrence compared to providing an educational booklet.

Burns et al. (2021) determined whether adding hip treatment to usual care for low back pain (LBP) improved disability and pain in individuals with LBP and a concurrent hip impairment. Seventy-six participants (age, 18 years or older; Oswestry Disability Index, 20% or greater; numeric pain-rating scale, 2 or more points) with LBP and a concurrent hip impairment were randomly assigned to a group that received treatment to the lumbar spine only (LBO group) ($n = 39$) or to one that received both lumbar spine and hip treatments (LBH group) ($n = 37$). The individual treating clinicians decided which specific low back treatments to administer to the LBO group. Treatments aimed at the hip (LBH

group) included manual therapy, exercise, and education, selected by the therapist from a predetermined set of treatments. Primary outcomes were disability and pain, measured by the Oswestry Disability Index and the numeric pain-rating scale, respectively, at baseline, 2 weeks, discharge, 6 months, and 12 months. The secondary outcomes were fear-avoidance beliefs (work and physical activity subscales of the Fear-Avoidance Beliefs Questionnaire), global rating of change, the Patient Acceptable Symptom State, and physical activity level. Investigators used mixed-model 2-by-3 analyses of variance to examine group-by-time interaction effects (intention-to-treat analysis). Data were available for 68 patients at discharge (LBH group, $n = 33$; LBO group, $n = 35$) and 48 at 12 months ($n = 24$ for both groups). There were no between-group differences in disability at discharge, 12 months, and all other time points. There were no between-group differences in pain at discharge, 12 months, and all other time points. There were no between-group differences in secondary outcomes, except for higher Fear-Avoidance Beliefs Questionnaire (work subscale) scores in the LBH group at 2 weeks and discharge. Authors concluded that adding treatments aimed at the hip to usual low back physical therapy did not provide additional short- or long-term benefits in reducing disability and pain in individuals with LBP and a concurrent hip impairment. Clinicians may not need to include hip treatments to achieve reductions in low back disability and pain in individuals with LBP and a concurrent hip impairment.

Nava-Bringas et al. (2021) compared the effectiveness of lumbar stabilization exercises and flexion exercises for pain control and improvements of disability in individuals with chronic low back pain (CLBP) and degenerative spondylolisthesis (DS). A randomized controlled trial was conducted in a tertiary public hospital and included 92 individuals over the age of 50 years who were randomly allocated to lumbar stabilization exercises or flexion exercises. Participants received 6 sessions of physical therapy (monthly appointments) and were instructed to execute exercises daily at home during the 6 months of the study. The primary outcome (measured at baseline, 1 month, 3 months, and 6 months) was pain intensity (visual analog scale, 0-100 mm) and disability (Oswestry Disability Index, from 0% to 100%). Secondary outcomes were disability (Roland-Morris Disability Questionnaire, from 0 to 24 points), changes in body mass index, and flexibility (fingertip to floor, in centimeters) at baseline and 6 months, and also the total of days of analgesic use at 6-month follow-up. Mean differences between groups were not significant for lumbar pain, radicular pain, for Oswestry scores, and for Roland Morris scores. Authors state that the findings from the present study reveal that flexion exercises are not inferior to and offer a similar response to stabilization exercises for the control of pain and improvements of disability in individuals with CLBP and DS.

De Campos et al. (2021) evaluate the evidence from randomized controlled trials (RCTs) on the effectiveness of prevention strategies to reduce future impact of low back pain (LBP), where impact is measured by LBP intensity and associated disability. 27 published reports of 25 different trials including a total of 8341 participants fulfilled the inclusion

criteria. The pooled results, from three RCTs (612 participants), found moderate-quality evidence that an exercise program can prevent future LBP intensity, and from 4 RCTs (471 participants) that an exercise and education program can prevent future disability due to LBP. It is uncertain whether prevention programs improve future quality of life (QoL) and workability due to the overall low-quality and very low-quality available evidence. Authors concluded that this review provides moderate-quality evidence that an exercise program, and a program combining exercise and education, are effective to reduce future LBP intensity and associated disability. It is uncertain whether prevention programs can improve future QoL and workability. Further high-quality RCTs evaluating prevention programs aiming to reduce future impact of LBP are needed.

George et al. (2021) updated a clinical practice guideline for treatment of low back pain. Findings relative to exercise included the following:

- Exercise For Acute Low Back Pain
 - Physical therapists can use exercise training interventions, including specific trunk muscle activation, for patients with acute low back pain (LBP) (grade C).
- Exercise For Acute Low Back Pain With Leg Pain
 - Physical therapists may use exercise training interventions, including trunk muscle strengthening and endurance and specific trunk muscle activation, to reduce pain and disability for patients with acute LBP with leg pain (grade B).
- Exercise For Chronic Low Back Pain
 - Physical therapists should use exercise training interventions, including trunk muscle strengthening and endurance, multimodal exercise interventions, specific trunk muscle activation exercise, aerobic exercise, aquatic exercise, and general exercise, for patients with chronic LBP (grade A).
 - Physical therapists may provide movement control exercise or trunk mobility exercise for patients with chronic LBP (grade B).
- Exercise For Chronic Low Back Pain With Leg Pain
 - Physical therapists may use exercise training interventions, including specific trunk muscle activation and movement control, for patients with chronic LBP with leg pain (grade B).
- Exercise For Chronic Low Back Pain With Movement Control Impairment
 - Physical therapists should use specific trunk muscle activation and movement control exercise for patients with chronic LBP and movement control impairment (grade A).
- Exercise For Chronic Low Back Pain In Older Adults
 - Physical therapists should use general exercise training to reduce pain and disability in older adults with chronic LBP (grade A).

• Exercise For Postoperative Low Back Pain

- Physical therapists can use general exercise training for patients with LBP following lumbar spine surgery (grade C).

Gianola et al. (2022) assessed the effectiveness of interventions for acute and subacute non-specific low back pain (NS-LBP) based on pain and disability outcomes in a systematic review with network meta-analysis. Forty-six RCTs ($n=8,765$) were included. At immediate-term follow-up, for pain decrease, exercise was considered one of the most efficacious treatments against an inert therapy. Similar findings were confirmed for disability. Fernández-Rodríguez et al. (2022) sought to determine which type of exercise is best for reducing pain and disability in adults with chronic low back pain (LBP) in a systematic review with a network meta-analysis (NMA) of randomized controlled trials (RCTs). Authors included 118 trials (9,710 participants). There were 28 head-to-head comparisons, 7 indirect comparisons for pain, and 8 indirect comparisons for disability. Compared with control, all types of physical exercises were effective for improving pain and disability, except for stretching exercises (for reducing pain) and the McKenzie method (for reducing disability). The most effective interventions for reducing pain were Pilates, mind-body, and core-based exercises. The most effective interventions for reducing disability were Pilates, strength, and core-based exercises. On SUCRA analysis, Pilates had the highest likelihood for reducing pain (93%) and disability (98%). Authors concluded that although most exercise interventions had benefits for managing pain and disability in chronic LBP, the most beneficial programs were those that included (1) at least 1 to 2 sessions per week of Pilates or strength exercises; (2) sessions of less than 60 minutes of core-based, strength, or mind-body exercises; and (3) training programs from 3 to 9 weeks of Pilates and core-based exercises.

Grooten et al. (2022) aimed to identify systematic reviews of common exercise types used in CLBP, to appraise their quality, and to summarize and compare their effect on pain and disability. The included reviews were grouped into nine exercise types: aerobic training, aquatic exercises, motor control exercises (MCE), resistance training, Pilates, sling exercises, traditional Chinese exercises (TCE), walking, and yoga. Out of the 253 full texts that were screened, we included 45 systematic reviews and meta-analyses. The quality of the included reviews ranged from high to critically low. Due to large heterogeneity, no meta-analyses were performed. Authors found low-to-moderate evidence of mainly short-term and small beneficial effects on pain and disability for MCE, Pilates, resistance training, TCE, and yoga compared to no or minimal intervention. Authors conclude that findings show that the effect of various exercise types used in CLBP on pain and disability varies with no major difference between exercise types. Essman and Lin (2022) highlighted the role of exercise in preventing and managing acute and chronic axial low back pain (LBP). They note that no single exercise method has been shown to be more effective than another. Overall, their review summarizes the beneficial role of a personalized exercise program and related counseling strategies in the prevention and management of LBP.

1 Bagg et al. (2022) estimated the effect of a graded sensorimotor retraining intervention
 2 (RESOLVE) on pain intensity in people with chronic low back pain. This parallel, 2-group,
 3 randomized clinical trial recruited participants with chronic (>3 months) nonspecific low
 4 back pain from primary care and community settings. A total of 276 adults were
 5 randomized (in a 1:1 ratio) to the intervention or sham procedure and attention control
 6 groups delivered by clinicians at a medical research institute in Sydney, Australia.
 7 Participants randomized to the intervention group ($n = 138$) were asked to participate in 12
 8 weekly clinical sessions and home training designed to educate them about and assist them
 9 with movement and physical activity while experiencing lower back pain. Participants
 10 randomized to the control group ($n = 138$) were asked to participate in 12 weekly clinical
 11 sessions and home training that required similar time as the intervention but did not focus
 12 on education, movement, and physical activity. The control group included sham laser and
 13 shortwave diathermy applied to the back and sham noninvasive brain stimulation. Among
 14 276 randomized patients completed follow-up at 18 weeks. The mean pain intensity was
 15 5.6 at baseline and 3.1 at 18 weeks in the intervention group and 5.8 at baseline and 4.0 at
 16 18 weeks in the control group, with an estimated between-group mean difference at 18
 17 weeks of -1.0 point, favoring the intervention group. In this randomized clinical trial
 18 conducted at a single center among patients with chronic low back pain, graded
 19 sensorimotor retraining, compared with a sham procedure and attention control,
 20 significantly improved pain intensity at 18 weeks. The improvements in pain intensity were
 21 small, and further research is needed to understand the generalizability of the findings.

22
 23 Fleckenstein et al. (2022) investigated the effects of individualized interventions, based on
 24 exercise alone or combined with psychological treatment, on pain intensity and disability
 25 in patients with chronic non-specific low-back-pain. Fifty-eight studies ($n = 10,084$) were
 26 included. At short-term follow-up (12 weeks), low-certainty evidence for pain intensity
 27 and very low-certainty evidence for disability indicates effects of individualized versus
 28 active exercises, and very low-certainty evidence for pain intensity, but not (low-certainty
 29 evidence) for disability compared to passive controls. At long-term follow-up (1 year),
 30 moderate-certainty evidence for pain intensity and disability indicates effects versus
 31 passive controls. Sensitivity analyses indicates that the effects on pain, but not on disability
 32 (always short-term and versus active treatments) were robust. Pain reduction caused by
 33 individualized exercise treatments in combination with psychological interventions (in
 34 particular behavioral-cognitive therapies) is of clinical importance. Certainty of evidence
 35 was downgraded mainly due to evidence of risk of bias, publication bias and inconsistency
 36 that could not be explained. Individualized exercise can treat pain and disability in chronic
 37 non-specific low-back-pain. The effects at short term are of clinical importance (relative
 38 differences versus active 38% and versus passive interventions 77%), especially in regard
 39 to the little extra effort to individualize exercise. Sub-group analysis suggests a
 40 combination of individualized exercise (especially motor-control based treatments) with
 41 behavioral therapy interventions to booster effects.

Niederer et al. (2022) investigated how risk of bias and intervention type modify effect sizes of exercise interventions that are intended to reduce chronic low back pain intensity. Potential effect modifiers were risk of bias, exercise modes, study, and meta-analyses characteristics. Data from 26 systematic reviews ($k = 349$ effect sizes, $n = 18,879$ participants) were analyzed. There was a clinically relevant effect overestimation in studies with a high risk of bias due to missing outcomes and low sample size. There was a clinically relevant underestimation of the effect when studies were at high risk of bias and outcome measurement. Motor control and stabilization training had the largest effects; stretching had the smallest effect. Authors concluded that the effects of exercise trials at high risk of bias may be overestimated or underestimated. After accounting for risk of bias, motor control and stabilization exercises may represent the most effective exercise therapies for chronic low back pain. Cashin et al. (2022) aimed to synthesize and appraise the current research to provide practical, evidence-based guidance concerning exercise prescription for non-specific CLBP. Systematic reviews show exercise is effective for small, short-term reductions in pain and disability, when compared with placebo, usual care, or waiting list control, and serious adverse events are rare. A range of individualized or group-based exercise modalities have been demonstrated as effective in reducing pain and disability. Authors conclude that to promote recovery, sustainable outcomes and self-management, exercise can be coupled with education strategies, as well as interventions that enhance adherence, motivation, and patient self-efficacy.

García-Moreno et al. (2022) upgraded the evidence of the most effective preventive physiotherapy interventions to improve back care in children and adolescents. Twenty studies were finally included. The most common physiotherapy interventions were exercise, postural hygiene, and physical activity. The mean age of the total sample was 11.79 years. Authors concluded that recent studies provide strong support for the use of physiotherapy in the improvement of back care and prevention of non-specific low back pain in children and adolescents. Based on GRADE methodology, they found that the evidence was from very low to moderate quality and interventions involving physical exercise, postural hygiene and physical activity should be preferred. Lindberg and Leggit (2022) summarized that there is low- to moderate-quality evidence that exercise reduces pain and improves function in patients with chronic low back pain compared with no treatment, usual care, and other conservative interventions such as education, manual therapy, and electrotherapy. This effect is clinically significant in the short term (six to 12 weeks) but less pronounced six months after treatment completion. The review does not recommend a specific exercise regimen to treat chronic low back pain.

Prat-Luri et al. (2023) analyzed the effect of trunk-focused exercise programs (TEPs) and moderator factors on chronic nonspecific low back pain (LBP). Forty randomized controlled trials ($n = 2,391$) were included. TEPs showed positive effects for all outcomes versus control. There were small effects in favor of TEPs versus general exercises for pain and disability. Trunk and/or hip range-of-motion improvements were associated with

greater reductions in pain and disability. Low body mass was associated with higher pain reduction. Authors concluded that trunk-focused exercise programs had positive effects on pain, disability, quality of life, and trunk performance compared to control groups, and on pain and disability compared to general exercises. Increasing trunk and/or hip range of motion was associated with greater pain and disability reduction, and lower body mass with higher pain reduction.

Ijzelenberg et al. (2023) evaluated the benefits and harms of exercise therapy for acute non-specific low back pain in adults compared to sham/placebo treatment or no treatment at short-term, intermediate-term, and long-term follow-up. This is an update of a Cochrane Review first published in 2005. Authors included RCTs that examined the effects of exercise therapy on non-specific LBP lasting six weeks or less in adults. Major outcomes for this review were pain, functional status, and perceived recovery. Minor outcomes were return to work, health-related quality of life, and adverse events. Main comparisons were exercise therapy versus sham/placebo treatment and exercise therapy versus no treatment. Outcomes were evaluated at short-term follow-up (time point within three months and closest to six weeks after randomization; main follow-up), intermediate-term follow-up (between nine months and closest to six months), and long-term follow-up (after nine months and closest to 12 months). Authors included 23 studies (13 from the previous review, 10 new studies) that involved 2,674 participants and provided data for 2,637 participants. Included studies were conducted in Europe ($N = 9$), the Asia-Pacific region ($N = 9$), and North America ($N = 5$); and most took place in a primary care setting ($N = 12$), secondary care setting ($N = 6$), or both ($N = 1$). In most studies, the population was middle-aged and included men and women. They judged 10 studies (43%) at low risk of bias with regard to sequence generation and allocation concealment. There is very low-certainty evidence that exercise therapy compared with sham/placebo treatment has no clinically relevant effect on pain scores in the short term. There is very low-certainty evidence that exercise therapy compared with sham/placebo treatment has no clinically relevant effect on functional status scores in the short term. There is very low-certainty evidence that exercise therapy compared with no treatment has no clinically relevant effect on pain or functional status in the short term. Owing to insufficient reporting of adverse events, authors were unable to reach any conclusions on the safety or harms related to exercise therapy. Authors concluded that exercise therapy compared to sham/placebo treatment may have no clinically relevant effect on pain or functional status in the short term in people with acute non-specific LBP, but the evidence is very uncertain. Exercise therapy compared to no treatment may have no clinically relevant effect on pain or functional status in the short term in people with acute non-specific LBP, but the evidence is very uncertain.

Li et al. (2023) evaluated the effects of different exercise therapies on chronic low back pain and provided a reference for exercise regimens in CLBP patients. This study included 75 randomized controlled trials (RCTs) with 5,254 participants. Network meta-analysis

results showed that tai chi, yoga (SMD, -1.76; 95% CI -2.72 to -0.81), Pilates exercise, and sling exercise showed a better pain improvement than conventional rehabilitation. Tai chi and yoga showed a better pain improvement than no intervention provided. Yoga and core or stabilization exercises showed a better physical function improvement than conventional rehabilitation. Yoga and core or stabilization exercises showed a better physical function improvement than no intervention provided. Authors concluded that compared with conventional rehabilitation and no intervention provided, tai chi, yoga, Pilates exercise, sling exercise, motor control exercise, and core or stabilization exercises significantly improved CLBP in patients. Compared with conventional rehabilitation and no intervention provided, yoga and core or stabilization exercises were statistically significant in improving physical function in patients with CLBP. Due to the limitations of the quality and quantity of the included studies, it is difficult to make a definitive recommendation before more large-scale and high-quality RCTs are conducted.

Kazeminia et al. (2023) aimed to estimate the results of randomized clinical trials (RCT) about the effect of pelvic floor muscle-strengthening exercises on reducing low back pain. Nineteen RCTs with a sample size of 456 subjects in the intervention group and 470 in the control group were included in the meta-analysis. Authors concluded that based on the results of the present meta-analysis, pelvic floor muscle-strengthening exercises significantly reduce the low back pain intensity. Therefore, these exercises can be regarded as a part of a low back pain management plan.

Wong et al. (2023) compared the effects of Pilates exercise (PE) with other forms of exercise on pain and disability in individuals with chronic non-specific low back pain (CNSLBP) and to inform clinical practice and future research. Eleven RCTs were included. A low certainty of evidence supported PE was more effective than general exercise (GE) in pain reduction. Moreover, very low levels of certainty were revealed for effectiveness of PE compared with direction-specific exercise (DSE) for pain reduction and equivalence of PE and spinal stabilization exercise (SSE) for pain and disability. Authors concluded that their review found no strong evidence for using one type of exercise intervention over another when managing patients with CNSLBP. Existing evidence does not allow this review to draw definitive recommendations. In the absence of a superior exercise form clinicians should work collaboratively with the patient, using the individual's goals and preferences to guide exercise selection. Further appropriately designed research is warranted to explore this topic further.

Zaina et al. (2023) identified evidence-based rehabilitation interventions for persons with non-specific low back pain (LBP) with and without radiculopathy and developed recommendations from high-quality clinical practice guidelines (CPGs) to inform the World Health Organization's (WHO) Package of Interventions for Rehabilitation (PIR). Four high-quality CPGs were identified. Recommended interventions included (1) education about recovery expectations, self-management strategies, and maintenance of

usual activities; (2) multimodal approaches incorporating education, exercise, and spinal manipulation; (3) nonsteroidal anti-inflammatory drugs combined with education in the acute stage; and (4) intensive interdisciplinary rehabilitation that includes exercise and cognitive/behavioral interventions for persistent pain. No high-quality CPGs for people younger than 16 years of age were found. Authors concluded that these recommendations emphasize the potential benefits of education, exercise, manual therapy, and cognitive/behavioral interventions.

Gilliam et al. (2023) assessed the effectiveness of mind-body (MB) exercise interventions provided by physical therapists for reducing pain and disability in people with low back pain (LBP). Randomized controlled trials evaluating the effects of Pilates, yoga, and tai chi interventions performed by physical therapists on pain or disability outcomes in adults with musculoskeletal LBP were included. Eight trials, 7 reporting on Pilates and 1 reporting on yoga, were included. Short-term outcomes for pain and indicated MB exercise was more effective than control intervention. Tests for subgroup differences between studies with exercise vs non-exercise control groups revealed a moderating effect on short-term outcomes where larger effects were observed in studies with non-exercise comparators. Long-term outcomes for pain and disability suggested that MB exercise is not more effective than control interventions for pain or disability. Quality of the evidence ranged from very low to low. Authors concluded that physical therapist-delivered MB exercise interventions, which overwhelmingly consisted of Pilates, were more effective than control in the short and long-term for pain and in the short-term for disability, with differences in the short-term effects lessened when compared with an active intervention. Pilates interventions delivered by physical therapists represent a viable tool for the clinical management of chronic LBP.

Ram et al. (2023) determined the effect of higher versus lower intensity exercise intensity on pain, disability, quality of life and adverse events in people with CLBP. Four trials (n = 214 participants, 84% male) reported across five studies were included. Higher intensity exercise reduced disability more than lower intensity exercise at end-treatment but not at 6-month follow-up. Higher intensity exercise did not reliably improve pain and quality of life more than lower intensity exercise. Adverse events did not differ between exercise intensities. All studies were at high risk of bias. Based on very low certainty evidence from a limited number of studies, exercise intensity does not appear to meaningfully influence clinical outcomes in people with CLBP.

Almeida et al. (2023) evaluated the effectiveness of the McKenzie method in people with (sub)acute non-specific low back pain in a Cochrane review. This review included five RCTs with a total of 563 participants recruited from primary or tertiary care. Three trials were conducted in the USA, one in Australia, and one in Scotland. Three trials received financial support from non-commercial funders and two did not provide information on funding sources. All trials were at high risk of performance and detection bias. None of the

1 included trials measured adverse events. McKenzie method versus minimal intervention
 2 (educational booklet; McKenzie method as a supplement to other intervention - main
 3 comparison): There is low-certainty evidence that the McKenzie method may result in a
 4 slight reduction in pain in the short term but not in the intermediate term. There is low-
 5 certainty evidence that the McKenzie method may not reduce disability in the short term
 6 nor in the intermediate term. McKenzie method versus manual therapy: There is low-
 7 certainty evidence that the McKenzie method may not reduce pain in the short term and
 8 may result in a slight increase in pain in the intermediate term. There is low-certainty
 9 evidence that the McKenzie method may not reduce disability in the short term nor in the
 10 intermediate term. McKenzie method versus other interventions (massage and advice):
 11 There is very low-certainty evidence that the McKenzie method may not reduce disability
 12 in the short term nor in the intermediate term. Authors concluded that based on low- to
 13 very low-certainty evidence, the treatment effects for pain and disability found in our
 14 review were not clinically important. Thus, they can conclude that the McKenzie method
 15 is not an effective treatment for (sub)acute NSLBP.

16
 17 Gilanyi et al. (2023) determined the effect of exercise on pain self-efficacy in adults with
 18 nonspecific chronic low back pain (NSCLBP). Authors included randomized controlled
 19 trials that compared the effect of exercise on pain self-efficacy to control, in adults with
 20 NSCLBP. Seventeen trials were included, of which eight ($n = 1,121$ participants; 60.6%
 21 female; mean age: 49.6 years) were included in the meta-analysis. Exercise increased pain
 22 self-efficacy by 3.02 points on the 60-point Pain Self-Efficacy Questionnaire. The certainty
 23 of evidence was moderate; all trials were at high risk of bias. Authors concluded that there
 24 was moderate-certainty evidence that exercise increased pain self-efficacy in adults with
 25 NSCLBP.

26
 27 Santos et al. (2023) evaluated the efficacy of Pilates on pain, functional disorders, and
 28 quality of life in patients with chronic low back pain (CLBP). Nineteen randomized
 29 controlled trials with a total of 1108 patients were included. Compared with the controls,
 30 this meta-analysis revealed that Pilates may have positive efficacy for pain relief and the
 31 improvement of functional disorders in CLBP patients, but the improvement in quality of
 32 life seems to be less obvious.

33
 34 Verville et al. (2023) evaluated benefits and harms of structured exercise programs for
 35 chronic primary low back pain (CPLBP) in adults to inform a World Health Organization
 36 (WHO) standard clinical guideline. Thirteen RCTs rated with overall low or unclear risk
 37 of bias were synthesized. Assessing individual exercise types (predominantly very low
 38 certainty evidence), pain reduction was associated with aerobic exercise and Pilates vs. no
 39 intervention, and motor control exercise vs. sham. Improved function was associated with
 40 mixed exercise vs. usual care, and Pilates vs. no intervention. Temporary increased minor
 41 pain was associated with mixed exercise vs. no intervention, and yoga vs. usual care. Little
 42 to no difference was found for other comparisons and outcomes. When pooling exercise

types, exercise vs. no intervention probably reduces pain in adults and functional limitations in adults and older adults (moderate certainty evidence). Authors concluded with moderate certainty that structured exercise programs probably reduce pain and functional limitations in adults and older people with CLBP.

Zhang et al. (2023) compared the efficacy of different exercises therapy on CLBP, dysfunction, quality of life, and mobility in the elderly. Sixteen articles (18 RCTs) were included, comprising a total of 989 participants. The quality of included studies was relatively high. Meta-analysis results indicated that exercise therapy could improve visual analog scale (VAS), Oswestry disability index (ODI), short-form 36-item health survey physical composite summary (SF-36PCS), short-form 36-item health survey mental composite summary (SF-36MCS), and timed up and go test (TUG). Exercise therapy effectively improved VAS, ODI, and SF-36 indexes in the elderly. Based on the subgroup, when designing the exercise therapy regimen, aerobics, strength, and mind-body exercise (≥ 12 weeks, ≥ 3 times/week, ≥ 60 min) should be considered carefully, to ensure the safety and effectiveness for the rehabilitation of CLBP patients.

Roren et al. (2023) critically reviewed available evidence regarding the efficacy of physical activity for people with LBP. They reported that in acute and subacute LBP, exercise did not reduce pain compared to no exercise. In chronic low back pain (CLBP), exercise reduced pain at the earliest follow-up compared with no exercise. In a recent systematic review, exercise improved function both at the end of treatment and in the long-term compared with usual care. Exercise also reduced work disability in the long-term. Authors were unable to establish a clear hierarchy between different exercise modalities. Multidisciplinary functional programs consistently improved pain and function in the short- and long-term compared with usual care and physiotherapy and improved the long-term likelihood of returning to work compared to non-multidisciplinary programs.

Heidari et al. (2023) aimed to systematically analyze the efficacy of aquatic exercise on pain intensity, disability, and quality of life among adults with low back pain. Out of 856 articles, 14 RCTs ($n = 484$ participants; 257 in the experimental groups and 227 in the control groups) met inclusion criteria. Pooled results illustrated that aquatic exercises significantly reduced pain, improved disability, and improved quality of life in both the physical component score and the mental component score when compared with a control group. Authors concluded that the current review showed that aquatic exercise regimens were effective among adults with low back pain. High-quality clinical investigations are still needed to support the use of therapeutic aquatic exercise in a clinical setting.

Babiloni-Lopez et al. (2023) aimed to systematically review and synthesize evidence (i.e., active [land-based training] and nonactive controls [e.g., receiving usual care]) regarding the effects of water-based training on patients with nonspecific chronic low-back pain (NSCLBP). The included studies satisfied the following criteria: (a) NSCLBP (≥ 12 weeks)

patients, (b) water-based intervention, (c) control group (land-based trained; nonactive group), and (d) outcomes related to pain, disability, quality of life, or flexibility. The main outcome analyzed in the meta-analysis was pain intensity. Secondary outcomes included disability, body mass index, and flexibility. After intervention, pain intensity was reduced compared with nonactive controls and a similar reduction was noted when compared with land-based trained group. Greater decrease in disability and greater increase in sit-and-reach were noted after intervention compared with the nonactive group. In conclusion, water-based exercise therapy reduces pain intensity, disability, and increases flexibility in NSCLBP compared with nonactive subjects and was equally effective compared with land-based exercise to reduce pain. Favorable effects may be expected at ≤ 8 weeks. However, due to several methodological issues (e.g., high heterogeneity), for the improvement of most outcomes, authors were unable to provide other than a weak recommendation in favor of intervention compared with control treatment.

Ceballos-Laito et al. (2023) evaluated the effectiveness of hip interventions on pain and disability in patients with LBP in the short-, medium-, and long-term. A total of 2,581 studies were screened. Eight were included in the meta-analysis involving 508 patients with LBP. The results provided very low certainty that both hip strengthening and hip stretching improved pain and disability in the short-term, respectively. No benefits were found in the medium- or long-term. The risk of bias, heterogeneity, and imprecision of the results downgraded the level of evidence. Very low certainty evidence suggests a positive effect of hip strengthening in isolation or combined with specific low back exercise and hip stretching combined with specific low back exercise for decreasing pain intensity and disability in the short-term, in patients with LBP.

Tikhile and Patil (2024) evaluated the efficacy of various physiotherapy strategies in alleviating LBP, considering a range of interventions and their associated outcomes. Through a thorough examination of existing literature from January 2017 to October 2023, this review synthesizes evidence on the effectiveness of interventions such as manual therapy, exercise therapy, electrotherapy modalities, and education-based approaches. The review also scrutinizes the comparative effectiveness of different physiotherapy modalities and their suitability for specific patient populations, considering factors such as chronicity, severity, and underlying pathology. This review aims to provide insights into the most effective physiotherapy strategies for alleviating LBP, chronic low back pain (CLBP) and chronic nonspecific low back pain (CNLBP) and guiding clinical practice toward evidence-based interventions. Twenty-one studies that fulfilled the criteria for inclusion (aged 20 to 50 years and of both genders) were added to the review. Exercises for core stability, strengthening, orthosis, transcutaneous electrical nerve stimulation, heat massage therapy, interferential current, Mulligan's mobilization, low-level laser therapy, and McGill stabilization exercises (core exercises) were among the therapeutic strategies. The McKenzie method (back exercises), ultrasound, sensory-motor training, Swiss ball exercises, and other techniques reduced pain and enhanced strength, balance, and ease of

daily activities. Every therapeutic approach has an impact on recovery rates ranging from minimal to maximal. Conventional physical therapy is less effective than most recent advanced techniques like mobilization and exercises. In summary, the integration of manual techniques, orthoses and alternative intervention strategies with conservative therapeutic approaches can effectively alleviate pain, enhance function and yield better overall outcomes. To get more information about the optimal dosage, therapeutic modalities and long-term effects of these treatments, more research is required.

El Melhat et al. (2024) explored the effectiveness and patient-related outcomes of various conservative approaches, including physical therapy modalities and alternative therapies in the treatment of lumbar disc herniation associated with radiculopathy (LDHR). The objective of this article was to introduce advanced and new treatment techniques, supplementing existing knowledge on various conservative treatments. Authors identified the following interventions to yield moderate evidence (Level B) of effectiveness for the conservative treatment of LDHR: patient education and self-management, McKenzie method, mobilization and manipulation, exercise therapy, traction (short-term outcomes), neural mobilization, and epidural injections. Two interventions were identified to have weak evidence of effectiveness (Level C): traction for long-term outcomes and dry needling. Three interventions were identified to have conflicting or no evidence (Level D) of effectiveness: electro-diagnostic-based management, laser and ultrasound, and electrotherapy.

Maharty et al. (2024) summarized evaluation and management of chronic LBP in adults. Patients with chronic low back pain should have a history and physical examination to identify red flags that may indicate serious conditions that warrant immediate intervention or yellow flags (i.e., psychological, environmental, and social factors) that indicate risk of disability. The examination should include an evaluation for radicular symptoms. Routine imaging is not recommended but is indicated when red flags are present, there is a neuromuscular deficit, or if pain does not resolve with conservative therapy. Patients should avoid bed rest. Nonpharmacologic treatment is first-line management and may include therapies with varying evidence of support, such as counseling, exercise therapy, spinal manipulation, massage, heat, dry needling, acupuncture, transcutaneous electrical nerve stimulation, and physical therapy. Pharmacologic interventions are second-line treatment. Nonsteroidal anti-inflammatory drugs are the initial medication of choice; duloxetine may also be beneficial. Evidence is inconclusive to recommend the use of benzodiazepines, muscle relaxants, antidepressants, corticosteroids, insomnia agents, anticonvulsants, cannabis, acetaminophen, or long-term opioids. Epidural corticosteroid injections are not recommended except for short-term symptom relief in patients with radicular pain. Most patients with chronic low back pain will not require surgery; evaluation for surgery may be considered in those with persistent functional disabilities and pain from progressive spinal stenosis, worsening spondylolisthesis, or herniated disk.

Zhou et al. (2024) aimed to identify and compare the recommendations of recent clinical practice guidelines (CPGs) for the management of LBP across the world. Analysis identified a total of 22 CPGs that met the inclusion criteria and were of middle and high methodological quality. The guidelines exhibited heterogeneity in their recommendations, particularly in the approach to different stages of LBP. For acute LBP, the guidelines recommended the use of non-steroidal anti-inflammatory drugs (NSAIDs), therapeutic exercise, staying active, and spinal manipulation. For subacute LBP, the guidelines recommended the use of NSAIDs, therapeutic exercise, staying active, and spinal manipulation. For chronic LBP, the guidelines recommended therapeutic exercise, the use of NSAIDs, spinal manipulation, and acupuncture. Authors state that current CPGs provide recommendations for almost all major aspects of the management of LBP, but there is marked heterogeneity between them. Some recommendations lack clarity and overlap with other treatments within the guidelines.

Patti et al. (2024) investigated the effectiveness of Pilates exercise on pain intensity and functional disability caused by LBP. Randomized controlled trials (RCTs) evaluating LBP in which the primary treatment was based on Pilates exercise compared with no exercise, or non-specific exercise were included in the literature search. The search returned 1566 records of which 36 articles were included in the systematic review and 19 in the meta-analysis. Twenty-two studies compared the effects of Pilates exercise vs no exercise and 13 studies examined the effects of Pilates exercise vs non-specific exercise. Analysis showed that Pilates had a positive effect on the perception of LBP vs no exercise. A similar trend occurred with non-specific exercise. Authors concluded that Pilates exercise can decrease LBP compared to no exercise and non-specific exercise. General practitioners should consider Pilates exercise as an effective strategy to manage LBP and counteract the growing health.

Liang et al. (2024) quantified the dose-response relationship between overall and specific exercise modalities and pain, in patients with nonspecific chronic low back pain. Authors included randomized controlled trials of exercise interventions in adults with nonspecific chronic LBP and at least 1 pain outcome reported at the main trial end point. Eighty-two trials were included (n = 5033 participants). They found a nonlinear dose-response relationship between total exercise and pain in patients with nonspecific chronic LBP. The maximum significant response was observed at 920 MET minutes. The minimal clinically important difference for achieving meaningful pain improvement was 520 MET minutes per week. The dose to achieve minimal clinically important difference varied by type of exercise; Pilates was the most effective. The certainty of the evidence was very low to moderate for all outcomes. Authors concluded that the dose-response relationship of different exercise modalities to improve pain in patients with nonspecific chronic LBP had a U-shaped trajectory and low- to moderate-certainty evidence. The clinical effect was most pronounced with Pilates exercise.

IJzelenberg et al. (2024) assessed the effectiveness of exercise for acute non-specific low back pain (LBP) vs : (1) sham treatment and (2) no treatment at short term (main follow-up time). The primary outcomes were back pain, back-specific functional status, and recovery. Authors identified 23 randomized controlled trials (2674 participants). There is very low-certainty evidence that exercise therapy compared with sham/placebo treatment has no clinically relevant effect on pain and on functional status in the short term. There is very low-quality evidence which suggests no difference in effect on pain and functional status for exercise vs no treatment at short-term follow-up. Similar results were found for the other follow-up moments. The certainty of the evidence was downgraded because many randomized controlled trials had a high risk of bias, were small in size, and/or there was substantial heterogeneity. Authors concluded that exercise therapy compared with sham/placebo and no treatment may have no clinically relevant effect on pain or functional status in the short term in people with acute non-specific LBP, but the evidence is very uncertain. Owing to insufficient reporting of adverse events, authors were unable to reach any conclusions on the safety or harms related to exercise therapy.

de Roode et al. (2024) investigated the proportion of improvement in pain and disability that can be attributed to contextual effects in the outcome of exercise therapy for patients with low back pain in a systematic review. Low back pain is the leading cause of global disability for which exercise therapy is a widely recommended treatment. Research indicates that contextual factors may also influence treatment outcomes in low back pain. Examples include the patient-therapist relationship and other treatment-related circumstances that affect patient expectations. By focusing on the specific treatment effect, clinical trials often ignore the effect of contextual factors, thereby contributing to the so-called efficacy paradox. This means that treatment effects observed in clinical practice are often greater than those reported in clinical trials. Authors conducted a meta-analysis with eligible articles reporting randomized controlled trials that compared exercise therapy to placebo interventions. Outcomes of interest were pain and disability. Meta-analysis was carried out to calculate the proportion attributable to contextual effects for both pain and disability. Eight studies met the inclusion criteria and were included in the meta-analysis. Five studies were rated as having a moderate risk of bias and two studies had a low risk of bias. Proportion attributable to contextual effects was 0.60 for pain and 0.69 for disability. Authors concluded that a large extent of pain and disability improvement after exercise therapy in low back pain is attributable to contextual effects although this conclusion is based on low certainty evidence.

Diez-Buil et al. (2024) compared whether the combination of exercise with education is more effective for the treatment of low back and/or pelvic pain (PP) than each of these interventions separately in pregnant women. A total of 13 articles were selected. There is a significant decrease in pain in the combination of exercise and education compared with education alone. With respect to disability, there is a significant decrease in the exercise and education group compared with the group that only addressed education). One article

analyzed kinesophobia, reporting no significant changes. Authors concluded that the combination of exercise and education seems to be more effective in reducing pain and disability in pregnant women with low back and/or PP than the use of education alone. In kinesophobia, the results found are not significant.

Herrero et al. (2024) analyzed whether interventions aimed at modifying lifestyle can be effective in improving pain intensity and functional disability in CNSLBP. A total of 20 studies were included for qualitative analysis, of which 16 were randomized clinical trials with a moderate-high methodological quality and were part of the quantitative analysis. The interventions that had the greatest effect in reducing pain intensity were cognitive therapy combined with functional exercise programs, lumbar stabilization exercise and resistance exercise; meanwhile, for functional disability, they were functional exercise programs, aerobic exercise and standard care. In conclusion, a multimodal intervention aimed at changing one's lifestyle that encompasses cognitive, behavioral, and physical aspects seems to be highly effective in improving pain intensity and functional disability caused by CNSLBP; however, it is not yet known if these improvements are maintained in the long term.

García-Moreno et al. (2024) compared the different physiotherapy treatments and determine the most effective treatment to reduce the nonspecific low back pain (NSLBP) intensity in children and adolescents. A meta-analysis of 11 controlled trials with 827 participants found that physiotherapy treatments effectively reduced NSLBP intensity on posttest measurement and 6-month follow-up. Network meta-analysis showed both therapeutic exercise and a combination of therapeutic exercise and manual therapy were effective compared to no treatment. There were no significant differences between therapeutic exercise and the combination of therapeutic exercise and manual therapy. Authors concluded that physical exercise has proven to be the most effective treatment for addressing the intensity of NSLBP in children and adolescents. While combining it with manual therapy may yield even better results, it is crucial to emphasize that physical exercise should serve as the cornerstone in the physiotherapeutic approach to managing NSLBP intensity in this age group.

Salehi et al. (2024) assessed the efficacy of specific exercises in general population with non-specific low back pain (LBP). Fifty-four trials met the inclusion criteria for this study. Additionally, 46 of these trials were randomized controlled trials and were further evaluated for the meta-analysis. Authors included trials investigating the effectiveness of exercise therapy, including isometric activation of deep trunk muscles, strengthening exercises, stabilization exercises, stretching exercises, and proprioceptive neuromuscular facilitation exercises (PNF) in LBP patients. The primary outcome was pain intensity, measured using tools such as the visual analogue scale (VAS) and numeric pain rating scale (NPRS). The secondary outcome was disability, assessed through instruments such as the Roland Morris Disability Questionnaire (RMDQ) and Oswestry Disability Index (ODI).

The meta-analysis indicated a small efficacy in favor of isometric activation of deep trunk muscles, a moderate efficacy in favor of stabilization exercises, and a large efficacy in favor of PNF exercises for reducing pain intensity as assessed by VAS or NPRS tools. Moreover, the meta-analysis revealed a moderate efficacy for isometric activation of deep trunk muscles, and a large efficacy for PNF exercises in improving disability, assessed using RMDQ or ODI questionnaires. The level of certainty in the evidence was very low to low. Authors concluded that these findings emphasize the importance of incorporating localized therapeutic exercises as a fundamental aspect of managing non-specific LBP. Clinicians should consider utilizing localized therapeutic exercise tailored to individual patient needs. Furthermore, further research investigating optimal exercise therapy, optimal dose of the exercises, durations, and long-term adherence is warranted to enhance the precision and efficacy of exercise-based interventions for non-specific LBP.

Gou et al. (2024) synthesized the available data on the effectiveness of hamstring stretching exercises in relieving pain intensity and improving function for patients with low back pain. Searches retrieved 344 trials, of which 14 met the inclusion criteria for this review (n = 735 participants). The combined meta-analysis showed hamstring stretching resulted in lower pain scores in different categories of low back pain. Subgroup analysis showed that hamstring stretching led to a larger range of motion for cases of back pain with radiating pain. The combined meta-analysis revealed that hamstring stretching resulted in lower Oswestry Disability Index scores in comparison to regular treatment, particularly in individuals suffering from low back pain across all subtypes. Authors concluded that this meta-analysis demonstrated the effectiveness of hamstring stretching exercises in reducing pain intensity in various categories of low back pain and improving the straight leg raise in patients experiencing back pain with radiating pain. Additionally, it highlights the improvement in function for patients with back pain across all subtypes.

Hennemann et al. (2024) determined the effectiveness of the McKenzie Method compared to any conservative interventions on pain and disability in patients with chronic low back pain (LBP) with directional preference (DP). Five trials (n = 743) were included. There was low-certainty evidence that the McKenzie Method, compared to all other interventions combined, produced clinically important reductions in short-term pain and in intermediate-term disability. Low-to-moderate certainty evidence showed that the McKenzie Method also resulted in clinically important improvements in short-term pain and disability when compared specifically to other exercise approaches, and in intermediate-term pain and disability as well as long-term disability when compared to minimal intervention. Low-certainty evidence showed usually small, clinically unimportant effects in comparison to manual therapy. Authors found low-to-moderate certainty evidence that the McKenzie Method was superior to all other interventions combined for up to 6 months for pain and up to 12 months for disability, with clinically important differences versus exercise in the short term and versus minimal interventions in the intermediate term. The only clinically important long-term effect was on disability compared to minimal intervention.

1 Lim et al. (2024) assessed the efficacy of pelvic floor muscle training and physical therapy
 2 interventions in patients with low back pain. Studies providing pelvic floor muscle training
 3 in individuals with low back pain were included. Nineteen studies were included in this
 4 review. Pelvic floor muscle training showed low certainty evidence for improving pain and
 5 reflected a clinically meaningful reduction in pain. The evidence for disability
 6 improvement had a low certainty due to high heterogeneity. Substantial improvements in
 7 pain and disability were observed when pelvic floor muscle training was added to standard
 8 physical therapy, with low certainty of evidence supporting these findings. Whereas pelvic
 9 floor muscle training substantially improved pain compared to other interventions, there
 10 was no marked improvement in disability. Authors concluded that pelvic floor muscle
 11 training is potentially beneficial in addition to physical therapy for reducing low back pain,
 12 particularly in pregnancy-related cases. However, the evidence should be interpreted
 13 considering the quality and risk of bias.

14
 15 Alonso-Sal et al. (2024) evaluated the effectiveness of exercise interventions for managing
 16 nonspecific low back pain (NSLBP) and explores their impact on related biopsychosocial
 17 factors, physical health variables, and inflammatory biomarkers. Materials and Methods:
 18 Fifteen randomized controlled trials involving 1338 participants aged 18 to 65 years with
 19 NSLBP were included. Exercise significantly reduced pain intensity and improved
 20 biopsychosocial factors such as depression, disability, functionality, quality of life, and
 21 kinesiophobia. Additionally, it enhanced physical parameters like proprioception, muscle
 22 thickness, and physical performance. However, the review found insufficient evidence
 23 regarding the effects of exercise on inflammatory biomarkers in NSLBP patients. Authors
 24 concluded that findings suggest that physical exercise is an effective intervention for pain
 25 reduction and the improvement of overall health in NSLBP, though further research is
 26 needed to clarify its impact on inflammation.

27
 28 Lim et al. (2025) summarized the content and critically appraised the quality and
 29 applicability of recent clinical practice guidelines (CPGs) for nonpharmacological,
 30 nonsurgical management of spine pain. Authors included 30 CPGs, primarily (90%)
 31 developed in Western countries, which contained 404 recommendations. High-quality
 32 CPGs consistently recommended exercise therapy and multimodal care, encompassing a
 33 combination of exercises, mobilization/manipulation, education, alternative medicine, and
 34 cognitive-behavioral treatments. Generally, CPGs did not recommend assistive (eg, corsets
 35 and orthosis) devices or electro/thermotherapies (eg, therapeutic ultrasound and
 36 transcutaneous electrical nerve stimulation). Approximately half of the CPGs demonstrated
 37 good methodological quality according to AGREE II, whereas the rest were of poor quality.
 38 On the AGREE-REX assessment, one third of the recommendations were of excellent
 39 quality. Although recent guidelines frequently recommended exercise therapy and
 40 multimodal care for the management of spine pain, their recommendations often
 41 overlooked demographics and comorbidities. Despite methodological improvements, most
 42 CPGs lacked simple clinical applicability and considerations of knowledge users' values.

1 PRACTITIONER SCOPE AND TRAINING

2 Practitioners should practice only in the areas in which they are competent based on their
3 education training and experience. Levels of education, experience, and proficiency may
4 vary among individual practitioners. It is ethically and legally incumbent on a practitioner
5 to determine where they have the knowledge and skills necessary to perform such services.

6
7 It is best practice for the practitioner to appropriately render services to a patient only if
8 they are trained, equally skilled, and adequately competent to deliver a service compared
9 to others trained to perform the same procedure. If the service would be most competently
10 delivered by another health care practitioner who has more skill and expert training, it
11 would be best practice to refer the patient to the more expert practitioner.

12
13 Best practice can be defined as a clinical, scientific, or professional technique, method, or
14 process that is typically evidence-based and consensus driven and is recognized by a
15 majority of professionals in a particular field as more effective at delivering a particular
16 outcome than any other practice (Joint Commission International Accreditation Standards
17 for Hospitals, 2020).

18
19 Depending on the practitioner's scope of practice, training, and experience, a member's
20 condition and/or symptoms during examination or the course of treatment may indicate the
21 need for referral to another practitioner or even emergency care. In such cases it is prudent
22 for the practitioner to refer the member for appropriate co-management (e.g., to their
23 primary care physician) or if immediate emergency care is warranted, to contact 911 as
24 appropriate. See the *Managing Medical Emergencies (CPG 159 – S)* clinical practice
25 guideline for information.

26 References

27
28 Abenhaim, L., Rossignol, M., Valat, J. P., Nordin, M., Avouac, B., Blotman, F., Charlot,
29 J., Dreiser, R. L., Legrand, E., Rozenberg, S., & Vautravers, P. (2000). The role of
30 activity in the therapeutic management of back pain. Report of the International Paris
31 Task Force on Back Pain. *Spine*, 25(4 Suppl), 1S–33S

32
33 Almeida MO, Narciso Garcia A, Menezes Costa LC, van Tulder MW, Lin CC, Machado
34 LA. The McKenzie method for (sub)acute non-specific low back pain. *Cochrane*
35 *Database Syst Rev*. 2023 Apr 5;4(4):CD009711

36
37 Alonso-Sal A, Alonso-Perez JL, Sosa-Reina MD, et al. Effectiveness of Physical Activity
38 in the Management of Nonspecific Low Back Pain: A Systematic Review. *Medicina*
39 (Kaunas). 2024;60(12):2065. Published 2024 Dec 16

40
41 Andersson G. B. (1999). Epidemiological features of chronic low-back pain. *Lancet*
42 (*London, England*), 354(9178), 581–585

- 1 Babiloni-Lopez C, Fritz N, Ramirez-Campillo R, Colado JC. Water-Based Exercise in
2 Patients With Nonspecific Chronic Low-Back Pain: A Systematic Review With Meta-
3 Analysis. *J Strength Cond Res*. 2024 Jan 1;38(1):206-219
4
- 5 Bagg MK, Wand BM, Cashin AG, et al. Effect of Graded Sensorimotor Retraining on Pain
6 Intensity in Patients With Chronic Low Back Pain: A Randomized Clinical Trial.
7 *JAMA*. 2022;328(5):430-439. Doi:10.1001/jama.2022.9930
8
- 9 Bogduk, N. and Jull, G. (1985). The theoretical pathology of acute locked back. A basis
10 for manipulative therapy. *23 (4)* 77-81
11
- 12 Bronfort, G., Haas, M., Evans, R., Kawchuk, G., & Dagenais, S. (2008). Evidence-
13 informed management of chronic low back pain with spinal manipulation and
14 mobilization. *The spine journal: official journal of the North American Spine*
15 *Society*, 8(1), 213–225
16
- 17 Broszko C, Golden K, Holmes CR, Fulleborn S, Biglow C, Guthmann R. Is exercise
18 therapy effective treatment for low back pain? *J Fam Pract*. 2021;70(8):E2-E3. Doi
19
- 20 Browder, D. A., Childs, J. D., Cleland, J. A., & Fritz, J. M. (2007). Effectiveness of an
21 extension-oriented treatment approach in a subgroup of subjects with low back pain: a
22 randomized clinical trial. *Physical therapy*, 87(12), 1608–1579
23
- 24 Brox, J. I., Storheim, K., Grotle, M., Tveito, T. H., Indahl, A., & Eriksen, H. R. (2008).
25 Systematic review of back schools, brief education, and fear-avoidance training for
26 chronic low back pain. *The spine journal: official journal of the North American Spine*
27 *Society*, 8(6), 948–958
28
- 29 Brumitt, J., Matheson, J. W., & Meira, E. P. (2013a). Core stabilization exercise
30 prescription, part I: current concepts in assessment and intervention. *Sports*
31 *health*, 5(6), 504–509
32
- 33 Brumitt, J., Matheson, J. W., & Meira, E. P. (2013b). Core stabilization exercise
34 prescription, part 2: a systematic review of motor control and general (global) exercise
35 rehabilitation approaches for patients with low back pain. *Sports health*, 5(6), 510–513
36
- 37 Burns SA, Cleland JA, Rivett DA, et al. When Treating Coexisting Low Back Pain and
38 Hip Impairments, Focus on the Back: Adding Specific Hip Treatment Does Not Yield
39 Additional Benefits-A Randomized Controlled Trial. *J Orthop Sports Phys Ther*.
40 2021;51(12):581-601

- 1 Carey, T. S., & Freburger, J. K. (2016). Exercise and the Prevention of Low Back Pain:
2 Ready for Implementation. *JAMA internal medicine*, 176(2), 208–209
- 3
- 4 Cashin AG, Booth J, McAuley JH, et al. Making exercise count: Considerations for the
5 role of exercise in back pain treatment. *Musculoskeletal Care*. 2022;20(2):259-270.
6 Doi:10.1002/msc.1597
- 7
- 8 Ceballos-Laita L, Estébanez-de-Miguel E, Jiménez-Rejano JJ, Bueno-Gracia E, Jiménez-
9 Del-Barrio S. The effectiveness of hip interventions in patients with low-back pain: A
10 systematic review and meta-analysis. *Braz J Phys Ther*. 2023 Mar-Apr;27(2):100502
- 11
- 12 Childs, J. D., Flynn, T. W., & Fritz, J. M. (2006). A perspective for considering the risks
13 and benefits of spinal manipulation in patients with low back pain. *Manual*
14 *therapy*, 11(4), 316–320
- 15
- 16 Childs, J. D., Fritz, J. M., Flynn, T. W., Irrgang, J. J., Johnson, K. K., Majkowski, G. R.,
17 & Delitto, A. (2004). A clinical prediction rule to identify patients with low back pain
18 most likely to benefit from spinal manipulation: a validation study. *Annals of internal*
19 *medicine*, 141(12), 920–928
- 20
- 21 Chou, R., Deyo, R., Friedly, J., Skelly, A., Hashimoto, R., Weimer, M., Fu, R., Dana, T.,
22 Kraegel, P., Griffin, J., Grusing, S., & Brodt, E. (2016). *Noninvasive Treatments for*
23 *Low Back Pain*. Agency for Healthcare Research and Quality (US)
- 24
- 25 Chou R. Low Back Pain. *Ann Intern Med*. 2021;174(8):ITC113-ITC128
- 26
- 27 Costa, L. O., Maher, C. G., Latimer, J., Hodges, P. W., Herbert, R. D., Refshauge, K. M.,
28 McAuley, J. H., & Jennings, M. D. (2009). Motor control exercise for chronic low back
29 pain: a randomized placebo-controlled trial. *Physical therapy*, 89(12), 1275–1286
- 30
- 31 De Campos TF, Maher CG, Fuller JT, Steffens D, Attwell S, Hancock MJ. Prevention
32 strategies to reduce future impact of low back pain: a systematic review and meta-
33 analysis. *Br J Sports Med*. 2021;55(9):468-476. Doi:10.1136/bjsports-2019-101436
- 34
- 35 de Roode A, Heymans MW, van Lankveld W, Staal JB. The impact of contextual effects
36 in exercise therapy for low back pain: a systematic review and meta-analysis. *BMC*
37 *Med*. 2024;22(1):484. Published 2024 Oct 23
- 38
- 39 Delitto, A., George, S. Z., Van Dillen, L., Whitman, J. M., Sowa, G., Shekelle, P.,
40 Denninger, T. R., Godges, J. J., & Orthopaedic Section of the American Physical
41 Therapy Association (2012). Low back pain. *The Journal of orthopaedic and sports*
42 *physical therapy*, 42(4), A1–A57

- 1 Diez-Buil H, Hernandez-Lucas P, Leirós-Rodríguez R, Echeverría-García O. Effects of the
2 combination of exercise and education in the treatment of low back and/or pelvic pain
3 in pregnant women: Systematic review and meta-analysis. *Int J Gynaecol Obstet.*
4 2024;164(3):811-822
- 5
- 6 Drummond C, Lebedeva V, Kirker K, Masaracchio M. Sling Exercise in the Management
7 of Chronic Low Back Pain: A Systematic Review and Meta-Analysis [published online
8 ahead of print, 2021 Sep 23]. *J Strength Cond Res.*
9 2021;10.1519/JSC.0000000000004135
- 10
- 11 El Melhat AM, Youssef ASA, Zebdawi MR, Hafez MA, Khalil LH, Harrison DE. Non-
12 Surgical Approaches to the Management of Lumbar Disc Herniation Associated with
13 Radiculopathy: A Narrative Review. *J Clin Med.* 2024;13(4):974. Published 2024
14 Feb 8
- 15
- 16 Essman M, Lin CY. The Role of Exercise in Treating Low Back Pain. *Curr Sports Med*
17 *Rep.* 2022;21(8):267-271. Published 2022 Aug 1
- 18
- 19 Fernández-Rodríguez R, Álvarez-Bueno C, Cavero-Redondo I, et al. Best Exercise Options
20 for Reducing Pain and Disability in Adults With Chronic Low Back Pain: Pilates,
21 Strength, Core-Based, and Mind-Body. A Network Meta-analysis. *J Orthop Sports*
22 *Phys Ther.* 2022;52(8):505-521
- 23
- 24 Ferreira, M. L., Ferreira, P. H., Latimer, J., Herbert, R. D., Hodges, P. W., Jennings, M.
25 D., Maher, C. G., & Refshauge, K. M. (2007). Comparison of general exercise, motor
26 control exercise and spinal manipulative therapy for chronic low back pain: A
27 randomized trial. *Pain, 131*(1-2), 31–37
- 28
- 29 Ferreira GE, Lin CC, Stevens ML, et al. Exercise Is Medicine, But Perhaps Not for
30 Preventing Low Back Pain: A Randomized Trial of Exercise and Education to Prevent
31 Low Back Pain Recurrence. *J Orthop Sports Phys Ther.* 2021;51(4):188-195
- 32
- 33 Fleckenstein J, Floessel P, Engel T, et al. Individualized Exercise in Chronic Non-Specific
34 Low Back Pain: A Systematic Review with Meta-Analysis on the Effects of Exercise
35 Alone or in Combination with Psychological Interventions on Pain and Disability. *J*
36 *Pain.* 2022;23(11):1856-1873
- 37
- 38 Flynn, T., Fritz, J., Whitman, J., Wainner, R., Magel, J., Rendeiro, D., Butler, B., Garber,
39 M., & Allison, S. (2002). A clinical prediction rule for classifying patients with low
40 back pain who demonstrate short-term improvement with spinal
41 manipulation. *Spine, 27*(24), 2835–2843

- 1 Foster, N. E., Anema, J. R., Cherkin, D., Chou, R., Cohen, S. P., Gross, D. P., Ferreira, P.
2 H., Fritz, J. M., Koes, B. W., Peul, W., Turner, J. A., Maher, C. G., & Lancet Low Back
3 Pain Series Working Group (2018). Prevention and treatment of low back pain:
4 evidence, challenges, and promising directions. *Lancet (London, England)*, 391(10137), 2368–2383
- 6
7 Fritz, J. M., Childs, J. D., & Flynn, T. W. (2005). Pragmatic application of a clinical
8 prediction rule in primary care to identify patients with low back pain with a good
9 prognosis following a brief spinal manipulation intervention. *BMC family
10 practice*, 6(1), 29
- 11
12 García-Moreno JM, Calvo-Muñoz I, Gómez-Conesa A, López-López JA. Effectiveness of
13 physiotherapy interventions for back care and the prevention of non-specific low back
14 pain in children and adolescents: a systematic review and meta-analysis. *BMC
15 Musculoskelet Disord*. 2022;23(1):314. Published 2022 Apr 2
- 16
17 García-Moreno JM, Calvo-Muñoz I, Gómez-Conesa A, López-López JA. Therapeutic
18 Exercise is Effective in Reducing the Intensity of Nonspecific Low Back Pain in
19 Children and Adolescents: A Systematic Review and Network Meta-analysis. *Arch
20 Phys Med Rehabil*. Published online November 16, 2024
- 21
22 Gatchel, R. J., & Mayer, T. G. (2008). Evidence-informed management of chronic low
23 back pain with functional restoration. *The spine journal: official journal of the North
24 American Spine Society*, 8(1), 65–69
- 25
26 Gatchel, R. J., & Rollings, K. H. (2008). Evidence-informed management of chronic low
27 back pain with cognitive behavioral therapy. *The spine journal: official journal of the
28 North American Spine Society*, 8(1), 40–44
- 29
30 George SZ, Fritz JM, Silfies SP, et al. Interventions for the Management of Acute and
31 Chronic Low Back Pain: Revision 2021. *J Orthop Sports Phys Ther*.
32 2021;51(11):CPG1-CPG60
- 33
34 Gianola S, Barger S, Del Castillo G, et al. Effectiveness of treatments for acute and
35 subacute mechanical non-specific low back pain: a systematic review with network
36 meta-analysis. *Br J Sports Med*. 2022;56(1):41-50
- 37
38 Gilanyi YL, Wewege MA, Shah B, Cashin AG, Williams CM, Davidson SRE, McAuley
39 JH, Jones MD. Exercise Increases Pain Self-efficacy in Adults With Nonspecific
40 Chronic Low Back Pain: A Systematic Review and Meta-analysis. *J Orthop Sports
41 Phys Ther*. 2023 Jun;53(6):335–342

- 1 Gilliam JR, George SZ, Norman KS, Hendren S, Sahu PK, Silfies SP. Mind-Body Exercise
2 Performed by Physical Therapists for Reducing Pain and Disability in Low Back Pain:
3 A Systematic Review With Meta-analysis. *Arch Phys Med Rehabil*. 2023
4 May;104(5):776-789
- 5
- 6 Gou Y, Lei H, Chen X, Wang X. The effects of hamstring stretching exercises on pain
7 intensity and function in low back pain patients: A systematic review with meta-
8 analysis of randomized controlled trials. *SAGE Open Med*.
9 2024;12:20503121241252251. Published 2024 Jul 29
- 10
- 11 Grooten WJA, Boström C, Dederig Å, et al. Summarizing the effects of different exercise
12 types in chronic low back pain – a systematic review of systematic reviews. *BMC*
13 *Musculoskelet Disord*. 2022;23(1):801. Published 2022 Aug 22
- 14
- 15 Guzmán, J., Esmail, R., Karjalainen, K., Malmivaara, A., Irvin, E., & Bombardier, C.
16 (2001). Multidisciplinary rehabilitation for chronic low back pain: systematic
17 review. *BMJ (Clinical research ed.)*, 322(7301), 1511–1516
- 18
- 19 Haldeman, S., & Dagenais, S. (2008). What have we learned about the evidence-informed
20 management of chronic low back pain? *The spine journal: official journal of the North*
21 *American Spine Society*, 8(1), 266–277
- 22
- 23 Hayden JA, van Tulder MW, Malmivaara A, Koes BW. Exercise therapy for treatment of
24 non-specific low back pain. *Cochrane Database Syst Rev*. 2005 Jul 20;(3):CD000335
- 25
- 26 Hayden JA, Wilson MN, Stewart S, Cartwright JL, Smith AO, Riley RD, van Tulder M,
27 Bendix T, Cecchi F, Costa LOP, Dufour N, Ferreira ML, Foster NE, Gudavalli MR,
28 Hartvigsen J, Helmhout P, Kool J, Koumantakis GA, Kovacs FM, Kuukkanen T, Long
29 A, Macedo LG, Machado LAC, Maher CG, Mehling W, Morone G, Peterson T,
30 Rasmussen-Barr E, Ryan CG, Sjögren T, Smeets R, Staal JB, Unsgaard-Tøndel M,
31 Wajswelner H, Yeung EW; Chronic Low Back Pain IPD Meta-Analysis Group.
32 Exercise treatment effect modifiers in persistent low back pain: an individual
33 participant data meta-analysis of 3514 participants from 27 randomised controlled
34 trials. *Br J Sports Med*. 2020 Nov;54(21):1277-1278
- 35
- 36 Hayden JA, Ellis J, Ogilvie R, Malmivaara A, van Tulder MW. Exercise therapy for
37 chronic low back pain. *Cochrane Database Syst Rev*. 2021a;9(9):CD009790. Published
38 2021 Sep 28
- 39
- 40 Hayden JA, Ellis J, Ogilvie R, et al. Some types of exercise are more effective than others
41 in people with chronic low back pain: a network meta-analysis. *J Physiother*.
42 2021b;67(4):252-262

- 1 Heidari F, Mohammad Rahimi N, Aminzadeh R. Aquatic Exercise Impact on Pain
2 Intensity, Disability and Quality of Life in Adults with Low Back Pain: A Systematic
3 Review and Meta-analysis. *Biol Res Nurs*. 2023 Oct;25(4):527-541
4
- 5 Hennemann V, Ziegelmann PK, Marcolino MAZ, Duncan BB. The McKenzie Method
6 delivered by credentialed therapists for chronic low back pain with directional
7 preference: systematic review with meta-analysis. *J Man Manip Ther*. Published online
8 October 9, 2024
9
- 10 Herrero P, Val P, Lapuente-Hernández D, Cuenca-Zaldívar JN, Calvo S, Gómez-Trullén
11 EM. Effects of Lifestyle Interventions on the Improvement of Chronic Non-Specific
12 Low Back Pain: A Systematic Review and Network Meta-Analysis. *Healthcare*
13 (Basel). 2024;12(5):505. Published 2024 Feb 20
14
- 15 Hicks, G. E., Fritz, J. M., Delitto, A., & McGill, S. M. (2005). Preliminary development of
16 a clinical prediction rule for determining which patients with low back pain will
17 respond to a stabilization exercise program. *Archives of physical medicine and*
18 *rehabilitation*, 86(9), 1753–1762
19
- 20 Hides, J. A., Jull, G. A., & Richardson, C. A. (2001). Long-term effects of specific
21 stabilizing exercises for first-episode low back pain. *Spine*, 26(11), E243–E248
22
- 23 Hlaing SS, Puntumetakul R, Khine EE, Boucaut R. Effects of core stabilization exercise
24 and strengthening exercise on proprioception, balance, muscle thickness and pain
25 related outcomes in patients with subacute nonspecific low back pain: a randomized
26 controlled trial. *BMC Musculoskelet Disord*. 2021;22(1):998. Published 2021 Nov 30
27
- 28 Hoffman, B. M., Papas, R. K., Chatkoff, D. K., & Kerns, R. D. (2007). Meta-analysis of
29 psychological interventions for chronic low back pain. *Health psychology: official*
30 *journal of the Division of Health Psychology, American Psychological*
31 *Association*, 26(1), 1–9
32
- 33 IJzelenberg W, Oosterhuis T, Hayden JA, Koes BW, van Tulder MW, Rubinstein SM, de
34 Zoete A. Exercise therapy for treatment of acute non-specific low back pain. *Cochrane*
35 *Database Syst Rev*. 2023 Aug 30;8(8):CD009365
36
- 37 IJzelenberg W, Oosterhuis T, Hayden JA, et al. Exercise Therapy for Treatment of Acute
38 Non-specific Low Back Pain: A Cochrane Systematic Review and Meta-analysis of
39 Randomized Controlled Trials. *Arch Phys Med Rehabil*. 2024;105(8):1571-1584

- Johannsen, F., Remvig, L., Kryger, P., Beck, P., Warming, S., Lybeck, K., Dreyer, V., & Larsen, L. H. (1995). Exercises for chronic low back pain: a clinical trial. *The Journal of orthopaedic and sports physical therapy*, 22(2), 52–59
- Jones, K. C., Tocco, E. C., Marshall, A. N., Valovich McLeod, T. C., & Welch Bacon, C. E. (2020). Pain Education With Therapeutic Exercise in Chronic Nonspecific Low Back Pain Rehabilitation: A Critically Appraised Topic. *Journal of sport rehabilitation*, 29(8), 1204–1209
- Jordan, A., Bendix, T., Nielsen, H., Hansen, F. R., Høst, D., & Winkel, A. (1998). Intensive training, physiotherapy, or manipulation for patients with chronic neck pain. A prospective, single-blinded, randomized clinical trial. *Spine*, 23(3), 311–319
- Karlsson M, Bergenheim A, Larsson MEH, Nordeman L, van Tulder M, Bernhardtsson S. Effects of exercise therapy in patients with acute low back pain: a systematic review of systematic reviews. *Syst Rev*. 2020 Aug 14;9(1):182
- Kazeminia M, Rajati F, Rajati M. The effect of pelvic floor muscle-strengthening exercises on low back pain: a systematic review and meta-analysis on randomized clinical trials. *Neurol Sci*. 2023 Mar;44(3):859-872
- Kofotolis, N., & Kellis, E. (2006). Effects of two 4-week proprioceptive neuromuscular facilitation programs on muscle endurance, flexibility, and functional performance in women with chronic low back pain. *Physical therapy*, 86(7), 1001–1012
- Koumantakis, G. A., Watson, P. J., & Oldham, J. A. (2005). Supplementation of general endurance exercise with randomized training versus general exercise only. Physiological and functional outcomes of a randomized controlled trial of patients with recurrent low back pain. *Clinical biomechanics (Bristol, Avon)*, 20(5), 474–482
- Kuukkanen, T. M., & Mätkiä, E. A. (2000). An experimental controlled study on postural sway and therapeutic exercise in subjects with low back pain. *Clinical rehabilitation*, 14(2), 192–202
- Lehtola, V., Luomajoki, H., Leinonen, V., Gibbons, S., & Airaksinen, O. (2016). Sub-classification based specific movement control exercises are superior to general exercise in sub-acute low back pain when both are combined with manual therapy: A randomized controlled trial. *BMC musculoskeletal disorders*, 17, 135
- Li Y, Yan L, Hou L, Zhang X, Zhao H, Yan C, Li X, Li Y, Chen X, Ding X. Exercise intervention for patients with chronic low back pain: a systematic review and network meta-analysis. *Front Public Health*. 2023 Nov 17;11:1155225

- 1 Liang Z, Tian S, Wang C, et al. The Best Exercise Modality and Dose for Reducing Pain
2 in Adults With Low Back Pain: A Systematic Review With Model-Based Bayesian
3 Network Meta-analysis. *J Orthop Sports Phys Ther.* 2024;54(5):315-327
4
- 5 Liddle, S. D., Baxter, G. D., & Gracey, J. H. (2004). Exercise and chronic low back pain:
6 what works? *Pain*, 107(1-2), 176–190
7
- 8 Lim Y, Do Y, Lee SH, Lee H. Efficacy of pelvic floor muscle training with physical therapy
9 for low back pain: A systematic review and meta-analysis. *Clin Rehabil.*
10 2024;38(12):1590-1608
11
- 12 Lim TH, Mak HY, Man Ngai SM, et al. Nonpharmacological Spine Pain Management in
13 Clinical Practice Guidelines: A Systematic Review Using AGREE II and AGREE-
14 REX Tools. *J Orthop Sports Phys Ther.* 2025;55(1):12-25
15
- 16 Lindberg B, Leggit JC. Effectiveness of Exercise Therapy in Patients With Chronic Low
17 Back Pain. *Am Fam Physician.* 2022;106(4):380-381
18
- 19 Macedo, L. G., Saragiotto, B. T., Yamato, T. P., Costa, L. O., Menezes Costa, L. C., Ostelo,
20 R. W., & Maher, C. G. (2016). Motor control exercise for acute non-specific low back
21 pain. *The Cochrane database of systematic reviews*, 2, CD012085
22
- 23 Maharty DC, Hines SC, Brown RB. Chronic Low Back Pain in Adults: Evaluation and
24 Management. *Am Fam Physician.* 2024;109(3):233-244
25
- 26 Manniche, C., & Jordan, A. (1995). The value of exercise therapy. *Spine*, 20(11), 1221–
27 1222
28
- 29 May, S., & Donelson, R. (2008). Evidence-informed management of chronic low back pain
30 with the McKenzie method. *The spine journal: official journal of the North American*
31 *Spine Society*, 8(1), 134–141
32
- 33 Mayer, J., Mooney, V., & Dagenais, S. (2008). Evidence-informed management of chronic
34 low back pain with lumbar extensor strengthening exercises. *The spine journal : official*
35 *journal of the North American Spine Society*, 8(1), 96–113
36
- 37 Nava-Bringas TI, Romero-Fierro LO, Trani-Chagoya YP, et al. Stabilization Exercises
38 Versus Flexion Exercises in Degenerative Spondylolisthesis: A Randomized
39 Controlled Trial. *Phys Ther.* 2021;101(8):pzab108

- 1 Niederer D, Weippert M, Behrens M. What Modifies the Effect of an Exercise Treatment
2 for Chronic Low Back Pain? A Meta-epidemiologic Regression Analysis of Risk of
3 Bias and Comparative Effectiveness. *J Orthop Sports Phys Ther.* 2022;52(12):792-802
4
- 5 Ostelo, R. W., van Tulder, M. W., Vlaeyen, J. W., Linton, S. J., Morley, S. J., & Assendelft,
6 W. J. (2005). Behavioural treatment for chronic low-back pain. *The Cochrane database*
7 *of systematic reviews*, (1), CD002014
8
- 9 Owen PJ, Miller CT, Mundell NL, Verswijveren SJJM, Tagliaferri SD, Brisby H, Bowe
10 SJ, Belavy DL. Which specific modes of exercise training are most effective for
11 treating low back pain? Network meta-analysis. *Br J Sports Med.* 2020
12 Nov;54(21):1279-1287
13
- 14 Patti A, Thornton JS, Giustino V, et al. Effectiveness of Pilates exercise on low back pain:
15 a systematic review with meta-analysis. *Disabil Rehabil.* 2024;46(16):3535-3548
16
- 17 Philadelphia Panel (2001). Philadelphia Panel evidence-based clinical practice guidelines
18 on selected rehabilitation interventions for low back pain. *Physical therapy*, 81(10),
19 1641–1674
20
- 21 Prat-Luri A, de Los Rios-Calonge J, Moreno-Navarro P, Manresa-Rocamora A, Vera-
22 Garcia FJ, Barbado D. Effect of Trunk-Focused Exercises on Pain, Disability, Quality
23 of Life, and Trunk Physical Fitness in Low Back Pain and How Potential Effect
24 Modifiers Modulate Their Effects: A Systematic Review With Meta-analyses. *J Orthop*
25 *Sports Phys Ther.* 2023;53(2):64-93
26
- 27 Qaseem, A., Wilt, T. J., McLean, R. M., Forciea, M. A., & Clinical Guidelines Committee
28 of the American College of Physicians (2017). Noninvasive Treatments for Acute,
29 Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline From the
30 American College of Physicians. *Annals of internal medicine*, 166(7), 514–530
31
- 32 Quentin C, Bagheri R, Ugbolue UC, et al. Effect of Home Exercise Training in Patients
33 with Nonspecific Low-Back Pain: A Systematic Review and Meta-Analysis. *Int J*
34 *Environ Res Public Health.* 2021;18(16):8430. Published 2021 Aug 10
35
- 36 Ram AK, Summers SJ, Booth J, Gibbs MT, Jones MD. Higher intensity exercise reduces
37 disability more than lower intensity exercise in adults with chronic low back pain: A
38 systematic review and meta-analysis. *Musculoskeletal Care.* 2023 Sep;21(3):611-622
39
- 40 Rathnayake APS, Sparkes V, Sheeran L. What is the effect of low back pain self-
41 management interventions with exercise components added? A systematic review with
42 meta-analysis. *Musculoskelet Sci Pract.* 2021;56:102469

- 1 Roren A, Daste C, Coleman M, Rannou F, Freyssenet D, Moro C, Lefèvre-Colau MM,
2 Nguyen C. Physical activity and low back pain: A critical narrative review. *Ann Phys
3 Rehabil Med.* 2023 Mar;66(2):101650
4
- 5 Saragiotto, B. T., Maher, C. G., Yamato, T. P., Costa, L. O., Menezes Costa, L. C., Ostelo,
6 R. W., & Macedo, L. G. (2016). Motor control exercise for chronic non-specific low-
7 back pain. *The Cochrane database of systematic reviews*, (1), CD012004
8
- 9 Salehi S, Sobhani V, Mir SM, Keivanfar N, Shamsoddini A, Hashemi SE. Efficacy of
10 specific exercises in general population with non-specific low back pain: A systematic
11 review and meta-analysis of randomized controlled trials. *J Bodyw Mov Ther.*
12 2024;39:673-705
13
- 14 Sherman, K. J., Cherkin, D. C., Erro, J., Miglioretti, D. L., & Deyo, R. A. (2005).
15 Comparing yoga, exercise, and a self-care book for chronic low back pain: a
16 randomized, controlled trial. *Annals of internal medicine*, 143(12), 849–856
17
- 18 Shi, Z., Zhou, H., Lu, L., Pan, B., Wei, Z., Yao, X., Kang, Y., Liu, L., & Feng, S. (2018).
19 Aquatic Exercises in the Treatment of Low Back Pain: A Systematic Review of the
20 Literature and Meta-Analysis of Eight Studies. *American journal of physical medicine
21 & rehabilitation*, 97(2), 116–122
22
- 23 Shiri, R., Coggon, D., & Falah-Hassani, K. (2018). Exercise for the Prevention of Low
24 Back Pain: Systematic Review and Meta-Analysis of Controlled Trials. *American
25 journal of epidemiology*, 187(5), 1093–1101
26
- 27 Skelly AC, Chou R, Dettori JR, Turner JA, Friedly JL, Rundell SD, Fu R, Brodt ED,
28 Wasson N, Kantner S, Ferguson AJR. Noninvasive Nonpharmacological Treatment for
29 Chronic Pain: A Systematic Review Update [Internet]. Rockville (MD): Agency for
30 Healthcare Research and Quality (US); 2020 Apr. Report No.: 20-EHC009. PMID:
31 32338846
32
- 33 Slade, S. C., & Keating, J. L. (2007). Unloaded movement facilitation exercise compared
34 to no exercise or alternative therapy on outcomes for people with nonspecific chronic
35 low back pain: a systematic review. *Journal of manipulative and physiological
36 therapeutics*, 30(4), 301–311
37
- 38 Standaert, C. J., Weinstein, S. M., & Rumpeltes, J. (2008). Evidence-informed
39 management of chronic low back pain with lumbar stabilization exercises. *The spine
40 journal: official journal of the North American Spine Society*, 8(1), 114–120

- 1 Stuber, K. J., Bruno, P., Sajko, S., & Hayden, J. A. (2014). Core stability exercises for low
2 back pain in athletes: a systematic review of the literature. *Clinical journal of sport*
3 *medicine: official journal of the Canadian Academy of Sport Medicine*, 24(6), 448–456
4
- 5 Suh, J. H., Kim, H., Jung, G. P., Ko, J. Y., & Ryu, J. S. (2019). The effect of lumbar
6 stabilization and walking exercises on chronic low back pain: A randomized controlled
7 trial. *Medicine*, 98(26), e16173
8
- 9 Thornton JS, Caneiro JP, Hartvigsen J, et al. Treating low back pain in athletes: a
10 systematic review with meta-analysis. *Br J Sports Med*. 2021;55(12):656-662
11
- 12 Tikhile P, Patil DS. Unveiling the Efficacy of Physiotherapy Strategies in Alleviating Low
13 Back Pain: A Comprehensive Review of Interventions and Outcomes. *Cureus*.
14 2024;16(3):e56013. Published 2024 Mar 12
15
- 16 Traeger AC, Qaseem A, McAuley JH. Low Back Pain. *JAMA*. 2021;326(3):286
17
- 18 van Dillen LR, Lanier VM, Steger-May K, et al. Effect of Motor Skill Training in
19 Functional Activities vs Strength and Flexibility Exercise on Function in People With
20 Chronic Low Back Pain: A Randomized Clinical Trial [published correction appears
21 in *JAMA Neurol*. 2021 Jan 19;:]. *JAMA Neurol*. 2021;78(4):385-395
22
- 23 van Tulder, M. W., Ostelo, R., Vlaeyen, J. W., Linton, S. J., Morley, S. J., & Assendelft,
24 W. J. (2000). Behavioral treatment for chronic low back pain: a systematic review
25 within the framework of the Cochrane Back Review Group. *Spine*, 25(20), 2688–2699
26
- 27 Vanti, C., Andreatta, S., Borghi, S., Guccione, A. A., Pillastrini, P., & Bertozzi, L. (2019).
28 The effectiveness of walking versus exercise on pain and function in chronic low back
29 pain: a systematic review and meta-analysis of randomized trials. *Disability and*
30 *rehabilitation*, 41(6), 622–632
31
- 32 Verville L, Ogilvie R, Hincapié CA, Southerst D, Yu H, Bussi res A, Gross DP, Pereira P,
33 Mior S, Tricco AC, Cedraschi C, Brunton G, Nordin M, Connell G, Wong JJ, Shearer
34 HM, Lee JGB, Wang D, Hayden JA, Cancelliere C. Systematic Review to Inform a
35 World Health Organization (WHO) Clinical Practice Guideline: Benefits and Harms of
36 Structured Exercise Programs for Chronic Primary Low Back Pain in Adults. *J Occup*
37 *Rehabil*. 2023 Dec;33(4):636-650
38
- 39 Wai, E. K., Rodriguez, S., Dagenais, S., & Hall, H. (2008). Evidence-informed
40 management of chronic low back pain with physical activity, smoking cessation, and
41 weight loss. *The spine journal: official journal of the North American Spine*
42 *Society*, 8(1), 195–202

- 1 Wang, X. Q., Zheng, J. J., Yu, Z. W., Bi, X., Lou, S. J., Liu, J., Cai, B., Hua, Y. H., Wu,
2 M., Wei, M. L., Shen, H. M., Chen, Y., Pan, Y. J., Xu, G. H., & Chen, P. J. (2012). A
3 meta-analysis of core stability exercise versus general exercise for chronic low back
4 pain. *PloS one*, 7(12), e52082
- 5
- 6 Wewege, M. A., Booth, J., & Parmenter, B. J. (2018). Aerobic vs. resistance exercise for
7 chronic non-specific low back pain: A systematic review and meta-analysis. *Journal of*
8 *back and musculoskeletal rehabilitation*, 31(5), 889–899
- 9
- 10 Whitman, J. M., Flynn, T. W., Childs, J. D., Wainner, R. S., Gill, H. E., Ryder, M. G.,
11 Garber, M. B., Bennett, A. C., & Fritz, J. M. (2006). A comparison between two
12 physical therapy treatment programs for patients with lumbar spinal stenosis: a
13 randomized clinical trial. *Spine*, 31(22), 2541–2549
- 14
- 15 Wong CM, Rugg B, Geere JA. The effects of Pilates exercise in comparison to other forms
16 of exercise on pain and disability in individuals with chronic non-specific low back
17 pain: A systematic review with meta-analysis. *Musculoskeletal Care*. 2023
18 Mar;21(1):78-96
- 19
- 20 World Health Organization (WHO). (2023). Low back pain. Retrieved on February 17,
21 2025 from <https://www.who.int/news-room/fact-sheets/detail/low-back-pain>
- 22
- 23 Yamato, T. P., Maher, C. G., Saragiotto, B. T., Hancock, M. J., Ostelo, R. W., Cabral, C.
24 M., Menezes Costa, L. C., & Costa, L. O. (2015). Pilates for low back pain. *The*
25 *Cochrane database of systematic reviews*, (7), CD010265
- 26
- 27 Yu Z, Yin Y, Wang J, Zhang X, Cai H, Peng F. Efficacy of Pilates on Pain, Functional
28 Disorders and Quality of Life in Patients with Chronic Low Back Pain: A Systematic
29 Review and Meta-Analysis. *Int J Environ Res Public Health*. 2023 Feb 6;20(4):2850
- 30
- 31 Zaina F, Côté P, Cancelliere C, Di Felice F, Donzelli S, Rauch A, Verville L, Negrini S,
32 Nordin M. A Systematic Review of Clinical Practice Guidelines for Persons With Non-
33 specific Low Back Pain With and Without Radiculopathy: Identification of Best
34 Evidence for Rehabilitation to Develop the WHO's Package of Interventions for
35 Rehabilitation. *Arch Phys Med Rehabil*. 2023 Nov;104(11):1913-1927
- 36
- 37 Zhang SK, Gu ML, Zhang T, Xu H, Mao SJ, Zhou WS. Effects of exercise therapy on
38 disability, mobility, and quality of life in the elderly with chronic low back pain: a
39 systematic review and meta-analysis of randomized controlled trials. *J Orthop Surg*
40 *Res*. 2023 Jul 19;18(1):513

- 1 Zhou T, Salman D, McGregor AH. Recent clinical practice guidelines for the management
2 of low back pain: a global comparison. BMC Musculoskelet Disord. 2024;25(1):344.
3 Published 2024 May 1
4
- 5 Zhu F, Zhang M, Wang D, Hong Q, Zeng C, Chen W. Yoga compared to non-exercise or
6 physical therapy exercise on pain, disability, and quality of life for patients with chronic
7 low back pain: A systematic review and meta-analysis of randomized controlled trials.
8 PLoS One. 2020 Sep 1;15(9):e0238544