

1	Clinical Practice Guideline:	Exercise Therapy for Treatment of Non-Specific Low Back Pain
2		
3		
4	Date of Implementation:	September 18, 2008
5		
6	Effective Date:	March 19, 2026
7		
8	Product:	Specialty
9		

10 **Table of Contents**

11	GUIDELINES	1
12	DESCRIPTION/BACKGROUND	1
13	EVIDENCE REVIEW.....	2
14	PRACTITIONER SCOPE AND TRAINING	42
15	<i>References</i>	43

16
17 **GUIDELINES**

18 American Specialty Health – Specialty (ASH) considers exercise therapy medically
19 necessary for treatment of patients with acute, sub-acute and chronic non-specific low back
20 pain.

21
22 **DESCRIPTION/BACKGROUND**

23 The World Health Organization (WHO) defines low back pain (LBP) as discomfort
24 experienced in the area between the bottom of the ribs and the buttocks. This pain can be
25 classified as acute (lasting under 6 weeks), sub-acute (6-12 weeks), or chronic (over 12
26 weeks). Also, according to the WHO, LBP can be specific or non-specific. Specific LBP
27 can be explained by an underlying disease (e.g., cancer), tissue damage (e.g., fracture), or
28 may be referred from other organs (e.g., from kidney or aortic aneurysm). Non-specific
29 means that the experience of pain cannot be confidently accounted for by another diagnosis
30 such as an underlying disease, pathology or tissue damage. LBP is non-specific in about
31 90% of cases.

32
33 Chronic low back pain is a significant problem because of high health care utilization,
34 rising health care costs, and perceived limitations of treatment effectiveness. Most patients
35 with chronic low back pain have non-specific low back pain. Exercise therapy is one
36 effective treatment option for chronic non-specific low back pain.

37
38 Exercise therapy represents a very diverse group of treatment approaches, which makes the
39 discussion of “exercise therapy” as a whole difficult (Hayden et al., 2005).

1 Hayden et al. (2005) proposed the following specific characteristics of exercise: type,
2 design, delivery, dose, and additional interventions.

3
4 Types of exercise therapy include muscle strengthening/stabilization/motor control
5 exercises, stretching/flexibility, coordination/balance/proprioceptive exercises, and
6 general fitness. Muscle strengthening typically involves repetitions of muscle contraction
7 of specific muscle groups aimed to increase muscle strength and/or endurance (Abenheim
8 et al., 2000). Stretching/flexibility entail movements of one or more joints, intended to
9 lengthen shortened muscles that can be static or dynamic in nature. Coordination and
10 balance exercises involve training in specific movements aimed at improving
11 proprioception and coordination of appropriate muscle groups (Johannsen et al., 1995;
12 Kuukkanen & Malkia, 2000). Finally, general physical fitness routines typically include
13 approaches involving the whole body (e.g., aerobic exercises) (Hayden et al., 2005).

14
15 Exercise therapy can also be categorized in terms of program design. Individualized
16 programs are those tailored to the individual based on the history and physical examination.
17 Partially individualized programs involve standard types of exercises, but at varied
18 intensity and/or duration. Finally, standard exercise programs are ones in which all
19 participants receive the same exercise program (Hayden et al., 2005).

20
21 Exercise programs can also be delivered in several ways: home, supervised home with
22 follow up, group supervision, and individual supervision. Home exercise entails
23 participants meeting initially with a therapist who provides them an exercise program to
24 do at home, with no supervision or follow up. Home exercise with follow up involves the
25 participants meeting initially with a therapist, doing the exercise program at home, and then
26 having a follow up visit with the therapist at least every 6 weeks. In group supervised
27 exercise, participants attend exercise sessions with 2 or more other individuals, under the
28 guidance of a therapist. Finally, individually supervised exercise sessions entail individuals
29 receiving one-on-one supervision while performing the prescribed exercise program
30 (Hayden et al., 2005).

31
32 Dose or intensity (measured by the duration and number of treatment sessions) is also an
33 important characteristic of exercise therapy (Hayden et al., 2005). Programs involving 20
34 or more hours of exercise are defined as high dose, and less than 20 hours of intervention
35 time as low dose. Factors such as load, resistance, and frequency of repetitions (which can
36 create a further categorization of strengthening exercise into strengthening vs. endurance)
37 may also be important issues when addressing exercise dose (Manniche & Jordan, 1995;
38 Jordan et al., 1998).

39 40 **EVIDENCE REVIEW**

41 Exercise is one of the few treatments for chronic low back pain with good literature
42 support; however, the effect sizes reported have been small and the exact type of exercise

1 that is most effective cannot be determined. In 2000, van Tulder et al. published a Cochrane
2 review assessing exercise therapy for low back pain relative to pain relief, functional status,
3 overall improvement and return to work. Thirty-nine randomized controlled trials (RCTs)
4 were included, and authors concluded that exercise therapy was not effective for acute low
5 back pain but may be helpful for chronic low back pain. Since 2000, many new trials have
6 been published, which precipitated the need for an updated review (Hayden et al., 2005).
7 In this 2005 review, 61 RCTs were included in the analysis. These studies involved adult
8 participants that could be categorized into acute, subacute, and chronic non-specific low
9 back pain groups. Studies involving low back pain caused by a specific pathology or
10 condition were excluded. Exercise therapy was defined as "a series of specific movements
11 with the aim of training or developing the body by a routine practice or as physical training
12 to promote good physical health." Studies included compared exercise therapy to a) no
13 treatment or placebo treatment, b) other conservative treatment, or c) other exercise group.
14 Outcomes of interest included self-reported pain intensity, condition-specific physical
15 functioning, global improvement, and return to work/absenteeism. Both qualitative and
16 quantitative rating systems were used to allow the most complete use of the available data.
17 Of the total 61 RCTs, 43 trials (3,907 individuals) assessed chronic low back pain. Thirty-
18 three exercise groups had non-exercise comparisons, and these trials provided strong
19 evidence that exercise therapy is at least as effective as other conservative interventions.
20 The evidence was conflicting as to whether exercise therapy was more effective than other
21 treatments for chronic low back pain. It also appeared that exercise therapy is most
22 effective when administered in a health care setting rather than as independent home
23 exercises. In many of these trials, other conservative care was used in addition to exercise
24 therapy; including behavioral and manual therapy, advice to stay active and education. As
25 an aside, there is moderate effectiveness of graded-activity exercise programs for the
26 subacute population. Only a small number of these studies were rated at high quality, which
27 may have led to an overestimation of effect. Also, many of the studies lacked information
28 to assess quality and clinical relevance. The most consistent outcome measure was for pain
29 intensity, which limits the ability to discuss other outcome measures. Lastly, authors found
30 potential publication bias, which also may have resulted in an overestimation of the
31 effectiveness of exercise therapy in the chronic low back pain population. Authors also
32 recommend that no further trials on the effectiveness of general exercise therapy for
33 chronic low back pain should be initiated, but rather trials should focus on specific exercise
34 intervention strategies in well-defined low back pain patient populations.

35
36 Another review by Liddle et al. (2004) based on 16 RCTs of high to medium quality
37 concluded that exercise as a primary intervention is an effective treatment for chronic low
38 back pain, despite the wide variety of exercise programs offered. Positive results were
39 maintained in 12 of the 16 trials, with supervision as a common feature. Again, authors felt
40 studies did not explain exercise programs adequately and thus, no conclusions could be
41 made regarding what type of exercise is most effective. The inclusion of exercise co-
42 interventions introduced a confounding influence as well.

1 To this end, a systematic review published in the Journal of Manipulative Physiological
2 Therapeutics in 2007 attempted to determine the effect of unloaded movement facilitation
3 exercises on outcomes for people with non-specific chronic low back pain (NSCLBP)
4 (Slade & Keating, 2007). In the previous systematic review reported by these authors, trunk
5 strengthening was effective for improving function and reducing pain, compared to no
6 exercise for patients with NSCLBP. Treatment effects increased with increasing exercise
7 intensity and adding motivational strategies. Trunk strengthening exercises compared to
8 aerobic training or the McKenzie approach showed no clear benefit (Slade & Keating,
9 2007). In their next review, 6 high quality RCTs were included. Participants were over 16
10 years of age with a current episode of low back pain lasting longer than 8 weeks (vs. the
11 typical >12 weeks) with or without a history of low back surgery. Given this duration
12 change, subjects could fall into the subacute category of low back pain rather than the
13 chronic group. Authors stated that these parameters were used to capture the largest number
14 of studies on exercise trials for chronic LBP that included the least number of participants
15 likely to demonstrate a natural recovery process during the intervention time. They also
16 defined low back pain as pain from below the scapulae to the buttock fold, with or without
17 lower extremity radiation. Again, this varied from the previously described reviews.
18 Interventions had to involve unloaded exercises that were likely to facilitate movement of
19 the lumbar spine. If other interventions were involved, the unloaded exercise portion
20 needed to be able to be partitioned out. Unloaded exercises basically referred to McKenzie
21 exercises or yoga. Studies were excluded if they combined unloaded exercises with
22 resistance exercises used to increase strengthening, spinal stabilization exercises or
23 behavioral approaches and could not separate each component.

24
25 For NSCLBP without surgery, use of a McKenzie approach produced small effects for
26 short and medium-term pain and short-term function compared to intensive trunk
27 strengthening. There were no observable differences in outcomes when comparing the
28 McKenzie approach to spinal stabilization exercises. When comparing yoga to trunk
29 strengthening and aerobic training in subjects with NSCLBP without surgery, comparable
30 effects were observed for short and medium-term outcomes. Compared to no exercise,
31 yoga displayed a significantly large effect for medium term pain and function. Performing
32 McKenzie exercises and yoga together compared with no exercise, significant, moderate
33 effects on medium-term pain and function were noted in favor of the unloaded exercise.
34 More specifically, within this review one of the RCTs published in the Annals of Internal
35 Medicine (Sherman et al., 2005) attempted to determine whether yoga was more effective
36 than conventional exercise or a self-care book for patients with chronic low back pain. One
37 hundred one adults participated in a 12-week yoga program or conventional therapy
38 program or just received a self-care book. They determined that yoga was more effective
39 than a self-care book. The yoga group consistently reported superior outcomes compared
40 with the exercise group, but these differences were not significant. Limitations included a
41 relatively short follow up period (14 weeks), modest sample sizes, reliance on class
42 instructors for intervention development and the inclusion of relatively highly educated

1 and functional participants (Sherman et al., 2005). Authors stated that it would be virtually
2 impossible to recreate these exercise programs, as minimal descriptions were reported.
3 Authors concluded that there is strong evidence that unloaded movement facilitation
4 exercise compared to no exercise is effective for improving pain and function. However, it
5 appears that when comparing unloaded exercise to other types of exercise, effects are
6 comparable. It may be that for patients with NSCLBP, unloaded exercise is as effective as
7 more vigorous forms of exercise that require more resources for relieving pain and
8 increasing function.

9
10 In another attempt to tease out what type of exercise is most beneficial, Kofotolis and Kellis
11 (2006) studied the effects of two 4-week Proprioceptive Neuromuscular Facilitation (PNF)
12 programs on muscle endurance, flexibility, and functional performance in women with
13 chronic low back pain. Unfortunately, these programs were only compared to one another
14 and not with another type of exercise program. Results demonstrated that both static and
15 dynamic PNF programs were effective in improving short-term trunk muscle endurance
16 and trunk mobility in people with chronic low back pain. Another RCT by Koumantakis et
17 al. (2005) compared a general trunk muscle endurance exercise program enhanced with
18 specific muscle stabilization exercises with a general exercise approach only. Fifty-five
19 patients with recurrent LBP were randomized to the two groups. Both groups received an
20 8-week intervention and written instructions. Results indicated that both the general
21 exercise program and the enhanced exercise program provided benefits for patients with
22 recurrent LBP. It appears to be the presence of physical exercise alone, rather than the
23 specific exercise type that is the factor in patient improvement in those with chronic LBP.

24
25 Another RCT compared general exercise, motor control exercise, and spinal manipulation
26 therapy for chronic low back pain (Ferreira et al., 2007). Each group received 8 weeks of
27 treatment. General exercise included strengthening, stretching and aerobic exercise, motor
28 control exercise included retraining of specific trunk musculature using ultrasound and
29 feedback, and spinal manipulation therapy involved both mobilization and manipulation.
30 At 8 weeks the motor control group and manipulation group had slightly better outcomes
31 than the general exercise group. At 6 and 12 months, these differences diminished, and
32 similar outcomes were reported. It appears that motor control exercise has better short-term
33 outcomes, while all three are equivalent over the medium and long-term with regards to
34 perceived effectiveness and function (Ferreira et al., 2007). Costa et al. (2009) completed
35 a randomized, placebo-controlled trial with subjects complaining of non-specific low back
36 with or without leg pain for at least 3 months. Subjects were instructed in specific deep
37 trunk muscle isolation exercise training which consisted of 12 individually supervised half-
38 hour sessions over an 8-week period. The placebo group received 20 minutes of detuned
39 short-wave diathermy and 5 minutes of detuned ultrasound for 12 sessions over an 8-week
40 period. Outcomes were measured at 2, 6, and 12 months. This study found that motor
41 control exercise produced short-term improvements in global impression of recovery and

1 activity, but not pain, for people with chronic low back pain. Most of the effects observed
2 in the short term were maintained at the six 6- and 12-month follow-ups (Costa, 2009).

3
4 In another review on use of the McKenzie method for chronic LBP by May and Donelson
5 (2008), they suggest that the McKenzie method plays an important role in the classification
6 of subgroups with different needs treatment-wise. It appears that as an intervention, this
7 method produces more positive short-term outcomes than non-specific guideline-based
8 care and equal or slightly better outcomes than stabilization or strengthening routines for
9 patients with chronic LBP (May & Donelson, 2008). Another review on lumbar extension
10 strengthening exercises for chronic LBP by Mayer et al. (2008) suggests that it is an
11 effective intervention over no treatment or most passive modalities, whether used in
12 isolation or as a co-intervention. These subjects report improved pain, disability, and other
13 reported outcomes in the short term. Over the long term, this review suggests that some of
14 the disability and pain benefits are lost. There also appears to be no clear benefit to lumbar
15 extensor strengthening exercises over other exercise programs regarding improvements in
16 pain, disability, strength, and endurance. Standaert et al. (2008) reported that lumbar
17 stabilization exercises for chronic low back pain are effective at improving pain and
18 function in a variety of patients with chronic LBP based on moderate evidence. Moderate
19 evidence also suggests that lumbar stabilization exercises are no more effective than
20 manual therapy. Strong evidence does exist that lumbar stabilization exercises are no more
21 effective than a less specific, general exercise program (Standaert et al., 2008).

22
23 There are a few well-designed studies that demonstrate the effectiveness of activity or
24 therapeutic exercise when used in conjunction with other manual interventions in the
25 management of spinal pain. Research has demonstrated the benefit of matching sub-
26 categories of patients to specific interventions. One of the interventions that has shown
27 marked success in the treatment of LBP is manipulation combined with strengthening
28 exercise. Flynn et al. (2002) reported 5 clinical predictors for success with spinal
29 manipulation (Symptom duration <16 days, No symptoms distal to the knee, Fear
30 Avoidance Belief Questionnaire Work Subscale <19, Hip IR >35 degrees, Positive lumbar
31 spring test on at least one lumbar segment). Flynn found a Positive Likelihood Ratio (+LR)
32 of 24 which provides a 95% chance of decreasing disability by >50% within the first two
33 (2) treatments using manipulation. Childs et al. (2004) validated this rule in a multi-center
34 trial and also determined the number needed to treat with thrust manipulation combined
35 with exercise to prevent one patient from experiencing a worsening of disability was only
36 ten. Childs et al. (2006) later reported that patients that met the clinical prediction rules
37 above were 8 times more likely to experience an increase in disability within one week if
38 they were not treated with a combined thrust manipulation/exercise intervention. This
39 Clinical Prediction Rule has also been validated in the Primary Care setting by Fritz et al.
40 (2005). The authors determined a +LR for success with thrust manipulation of 7.2 with the
41 following two factors present: symptoms less than 16 days duration and no symptoms distal
42 to the knee.

1 The literature demonstrates that an Extension Oriented Treatment Approach (EOTA) is
 2 beneficial in patients who demonstrate a directional preference (DP) of symptom
 3 centralization with extension postures/exercises (Browder et al., 2007). The average
 4 duration of the patients’ symptoms was 3 months. The authors compared an EOTA with
 5 strengthening exercises and reported the EOTA group demonstrated greater improvements
 6 in disability and pain at 1 week follow-up and greater improvement in disability at 4 weeks
 7 and 6-month follow-ups as well. The EOTA was provided over the course of 8 sessions
 8 (twice a week for 4 weeks) and included the following interventions:

- 9 1. Extension-oriented exercises (sustained and repeated) in prone and standing;
- 10 2. Posterior to Anterior (PA) lumbar mobilizations, grade I to IV, 10 to 20
 11 oscillations;
- 12 3. Home exercise prescription (prone press-up) x10 repetitions every 2 to 3 waking
 13 hours (may substitute standing extension exercises).

14
 15 The Orthopaedic Section of the American Physical Therapy Association (APTA) has an
 16 ongoing effort to create evidence-based practice guidelines for orthopaedic physical
 17 therapy management of patients with musculoskeletal impairments described in the World
 18 Health Organization’s International Classification of Functioning, Disability, and Health
 19 (ICF). In 2012, Delitto et al. authored guidelines for low back pain. The purpose of these
 20 low back pain clinical practice guidelines was to describe the peer-reviewed literature and
 21 make recommendations related to (1) treatment matched to low back pain subgroup
 22 responder categories, (2) treatments that have evidence to prevent recurrence of low back
 23 pain, and (3) treatments that have evidence to influence the progression from acute to
 24 chronic low back pain and disability. Authors presented “A” level recommendations for
 25 treatment of low back pain which included manual therapy, trunk coordination,
 26 strengthening and endurance exercises, centralization and directional preference exercises
 27 and progressive endurance exercises and fitness activities. Research has determined thrust
 28 manipulation is effective in a subgroup of patients as part of a multi-component program
 29 including exercise. Lumbar coordination, strengthening and endurance exercises are a
 30 common treatment intervention for back pain. They are also referred to in the literature as
 31 motor control exercises, transversus abdominis training, lumbar multifidus training and
 32 dynamic lumbar stabilization exercises. Delitto et al. (2012) summarized the available
 33 literature indicating that clinicians should consider these exercises to reduce low back pain
 34 and disability in patients with subacute and chronic low back pain with movement
 35 dysfunction and in patients post microdiscectomy. Much of the research demonstrates that
 36 these exercises are effective but may be no more effective than a general exercise program.
 37 Centralization exercises appear to be beneficial for patients with acute low back pain with
 38 referred lower extremity pain. Clinicians should consider using repeated movements and
 39 exercises to promote centralization through reduction of lower extremity pain. Also,
 40 repeated movements in a specific direction, as noted by treatment response, should be
 41 utilized to reduce symptoms and improve mobility in all phases of low back pain. Lastly,
 42 progressive endurance exercises and fitness activities are endorsed by most current low

1 back pain guidelines with moderate to high levels of evidence. Aerobic conditioning has
2 been hypothesized to reduce pain perception and improving function in patients with
3 chronic low back pain and other generalized pain.

4
5 A meta-analysis by Wang et al. (2012) concluded that core stability exercises are more
6 effective in decreasing pain and may improve physical function in patients with chronic
7 low back in the short-term relative to general exercise. However, over the long term, no
8 significant differences were noted. In 2013, Brumitt et al. (2013a) provided clinical
9 recommendations using the SORT (Strength of Recommendation Taxonomy) method.
10 They concluded that a therapeutic intervention program consisting of motor control
11 exercises OR general back strengthening exercises may be beneficial for patients with low
12 back pain lasting longer than 6 weeks. However, given the SORT evidence rating of ‘B’
13 indicates that the evidence is inconsistent or of limited quality. Brumitt et al. (2013b)
14 published another paper analyzing randomized controlled trials that assessed the effects of
15 a motor control exercise approach, a general exercise approach, or both for patients with
16 low back pain that were published in scientific peer-reviewed journals. Fifteen studies were
17 identified (8, motor control exercise approach without general exercise comparison; 7,
18 general exercise approach with or without motor control exercise approach comparison).
19 Authors stated that current evidence suggests that exercise interventions may be effective
20 at reducing pain or disability in patients with low back pain, but it may not be necessary to
21 prescribe exercises purported to restore motor control of specific muscles. A systematic
22 review by Stuber et al. (2014) reviewed the effectiveness of core stability exercises for low
23 back pain in athletes. They concluded that given the low quantity and quality of available
24 literature, no strong conclusions could be formulated.

25
26 Lehtola et al. (2016) conducted a randomized controlled trial (RCT) to compare the effects
27 of general exercise versus specific movement control exercise (SMCE) on disability and
28 function in patients with MCI within the recurrent sub-acute LBP group. Subjects attended
29 5 sessions of either specific or general exercises. Both groups also received a short
30 application of manual therapy. The primary outcome was disability, assessed by the
31 Roland-Morris Disability Questionnaire (RMDQ). The measurements were taken at
32 baseline, immediately after the 3 months intervention and at twelve-month follow-up.
33 Measurements of 61 patients (SMCE $n = 30$ and general exercise $n = 31$) were completed
34 at 12 months. Patients in both groups reported significantly less disability at 12 months
35 follow up, with the SMCE group showing statistically significantly superior improvement.
36 However, the result did not reach the clinically significant three-point difference. There
37 was no statistical difference between the groups measured with Oswestry Disability Index
38 (ODI). Authors concluded for subjects with non-specific recurrent sub-acute LBP and MCI
39 an intervention consisting of SMCE and manual therapy combined may be superior to
40 general exercise combined with manual therapy. Saragiotto et al. (2016) authored a
41 Cochrane Review on motor control exercise for chronic non-specific low back pain.
42 (CNSLBP). As noted in the previous literature, exercise is a modestly effective treatment

1 for chronic LBP and current evidence suggests that no single form of exercise is superior
2 to another. Authors report that among the most commonly used exercise interventions are
3 motor control exercise (MCE). To clarify, MCE intervention focuses on the activation of
4 the deep trunk muscles and targets the restoration of control and co-ordination of these
5 muscles, progressing to more complex and functional tasks integrating the activation of
6 deep and global trunk muscles. Authors included trials comparing MCE with no treatment,
7 another treatment or that added MCE as a supplement to other interventions. Primary
8 outcomes were pain intensity and disability. They also considered function, quality of life,
9 return to work or recurrence as secondary outcomes. They considered the following time
10 points: short-term (less than 3 months after randomization); intermediate (at least three
11 months but less than 12 months after randomization); and long-term (12 months or more
12 after randomization) follow-up. 29 trials ($n = 2,431$) were included in this review. The
13 study sample sizes ranged from 20 to 323 participants. Results demonstrate that there is
14 low to high quality evidence that MCE is not clinically more effective than other exercises
15 for all follow-up periods and outcomes tested. When compared to minimal intervention,
16 there is low to moderate quality evidence that MCE is effective for improving pain at short,
17 intermediate and long-term follow-up with medium effect sizes. There was also a clinically
18 important difference for the outcomes function and global impression of recovery
19 compared with minimal intervention. There was moderate to high quality evidence that
20 there is no clinically important difference between MCE and manual therapy for all follow-
21 up periods and outcomes tested. Finally, there was very low to low quality evidence that
22 MCE is clinically more effective than exercise and electrophysical agents (EPA) for pain,
23 disability, global impression of recovery and quality of life with medium to large effect
24 sizes. Minor or no adverse events were reported in the included trials. Authors conclude
25 that given the evidence that MCE is not superior to other forms of exercise, the choice of
26 exercise for chronic LBP should probably depend on patient or therapist preferences,
27 therapist training, costs, and safety.

28
29 Macedo et al. (2016) completed a Cochrane Review on the effectiveness of motor control
30 exercise for acute non-specific low back pain. They only included RCTs examining the
31 effectiveness of MCE for patients with acute non-specific LBP. Authors considered trials
32 comparing MCE versus no treatment, versus another type of treatment or added as a
33 supplement to other interventions. Primary outcomes were pain intensity and disability.
34 Secondary outcomes were function, quality of life and recurrence. Authors considered the
35 following follow-up intervals: short term (less than three months after randomization);
36 intermediate term (at least three months but within 12 months after randomization); and
37 long term (12 months or longer after randomization). Only 3 trials were included with study
38 samples ranging from 33 to 123 participants. Evidence of very low to moderate quality
39 indicates that MCE showed no benefit over spinal manipulative therapy, other forms of
40 exercise or medical treatment in decreasing pain and disability among patients with acute
41 and subacute low back pain. Whether MCE can prevent recurrences of LBP remains

1 uncertain and no firm conclusions can be drawn regarding the effectiveness of MCE for
2 acute LBP.

3
4 Pilates was also examined in a Cochrane Review as a treatment for non-specific LBP
5 (Yamato et al., 2015). They included RCTs that examined the effectiveness of Pilates
6 intervention in adults with acute, subacute, or chronic non-specific low back pain. The
7 primary outcomes considered were pain, disability, global impression of recovery and
8 quality of life. A total of 6 trials compared Pilates to minimal intervention. They did not
9 find any high-quality evidence for any of the treatment comparisons, outcomes or follow-
10 up periods investigated. However, there is low to moderate quality evidence that Pilates is
11 more effective than minimal intervention for pain and disability. When Pilates was
12 compared with other exercises, the authors found a small effect for function at
13 intermediate-term follow-up. Thus, while there is some evidence for the effectiveness of
14 Pilates for low back pain, there is no conclusive evidence that it is superior to other forms
15 of exercises. The decision to use Pilates for low back pain may be based on the patient's or
16 care provider's preferences, and costs.

17
18 A systematic review and meta-analysis by Carey and Freburger (2016) assessed research
19 into the value of exercise as a way to treat and prevent LBP. The study found that exercise
20 alone was linked to a 35% reduction in risk, while the combination of exercise and
21 education was associated with a 45% risk reduction for up to one year. The use of exercise
22 was also found to result in a 78% reduction in sick leave for LBP. Authors found that while
23 education helped to further reduce that risk when combined with exercise, education alone
24 doesn't seem to have much effect, according to authors. They also suggest that for exercise
25 to remain protective against future LBP, it needs to be ongoing.

26
27 The Agency for Healthcare Research and Quality (AHRQ) published a Comparative
28 Effectiveness Review in 2016 on noninvasive treatments for LBP. They summarized the
29 research on exercise and LBP with the following key points:

- 30 1. For acute LBP, a systematic review found no differences between exercise therapy
31 versus no exercise in pain or function; for subacute LBP, there were no differences
32 in pain or function. Three other trials for acute to subacute LBP found inconsistent
33 results of exercise vs. usual care to improve pain and function.
- 34 2. For chronic LBP, a systematic review found exercise was associated with greater
35 pain relief versus no exercise and a more recent review using more restrictive
36 criteria and additional trials were consistent with these earlier findings.
- 37 3. More specifically, for chronic LBP, a review found motor control exercise was
38 associated with lower pain scores and better function in the short, intermediate and
39 long term vs. minimal intervention. Another systematic review found MCE
40 associated with lower pain intensity at the short term and intermediate term versus
41 general exercise. No significant findings were noted in the long term. Better
42 function was noted with MCE in the short and long term.

- 1 4. For radicular LBP, three trials not included in any systematic reviews found effects
2 that favored exercise versus usual care or no exercise in pain and function, though
3 effect sizes were small.
- 4 5. For comparisons of different exercise types, there were no clear differences for
5 patients with acute or chronic LBP.
- 6 6. Adverse events were not often reported and if they were, typically muscle soreness
7 and increased pain were reported. No serious harms were reported.

8

9 According to Qaseem et al. (2017), moderate-quality evidence showed that exercise
10 therapy resulted in small improvements in pain and function. Specific components
11 associated with greater effects on pain included individually designed programs,
12 supervised home exercise, and group exercise; regimens that included stretching and
13 strength training were most effective. In a systematic review, Vanti et al. (2019) found that
14 pain, disability, quality of life and fear-avoidance similarly improve by walking or exercise
15 in chronic low back pain. Walking may be considered as an alternative to other physical
16 activity. Adding walking to exercise does not induce greater improvement in the short-
17 term. Walking may be a less-expensive alternative to physical exercise in chronic low back
18 pain. Wewege et al. (2018) compared progressive aerobic training (PAT) to progressive
19 resistance training (PRT) for pain, disability, and quality of life (QoL) in people with
20 chronic non-specific low back pain (CNSLBP). Six studies were included, comprising 333
21 participants. Exercise significantly reduced pain intensity although neither mode proved
22 superior. PRT significantly improved the Short Form Health Survey-Mental Component
23 Score. Authors concluded that PAT and PRT decreased pain intensity in individuals with
24 CNSLBP although neither mode was superior. Resistance exercise improved psychological
25 wellbeing. High-quality RCTs comparing PAT, PRT, and PAT + PRT, are required. Shi et
26 al. (2018) analyzed all evidence available in the literature about effectiveness of the aquatic
27 exercise. Eight trials involving 331 patients were included in the meta-analysis, and the
28 results showed a relief of and physical function after aquatic exercise. However, there was
29 no significant effectiveness with regard to general mental health in aquatic group. Authors
30 concluded that aquatic exercise can statistically significantly reduce pain and increase
31 physical function in patients with low back pain. Shiri et al. (2018) assessed the effect of
32 exercise in population-based interventions to prevent low back pain (LBP) and associated
33 disability. Thirteen randomized controlled trials (RCTs) and 3 nonrandomized controlled
34 trials (NRCTs) qualified for the meta-analysis. Exercise alone reduced the risk of LBP by
35 33% and exercise combined with education reduced it by 27%. The severity of LBP and
36 disability from LBP were also lower in exercise groups than in control groups. Authors
37 concluded that exercise reduces the risk of LBP and associated disability, and a
38 combination of strengthening with either stretching or aerobic exercises performed 2-3
39 times per week can reasonably be recommended for prevention of LBP in the general
40 population. Suh et al. (2019) compared the efficiency between 2 exercises: the
41 individualized graded lumbar stabilization exercise (IGLSE) and walking exercise (WE).
42 A randomized controlled trial was conducted in 48 participants with chronic LBP. After

1 screening, participants were randomized to 1 of 4 groups: flexibility exercise (FE), WE,
2 stabilization exercise (SE), and stabilization with WE (SWE) groups. Participants
3 underwent each exercise for 6 weeks. The primary outcome was visual analog scale (VAS)
4 of LBP during rest and physical activity. Secondary outcomes were as follows: VAS of
5 radiating pain measured during rest and physical activity; frequency of medication use
6 (number of times/day); Oswestry disability index; Beck Depression Inventory; endurance
7 of specific posture; and strength of lumbar extensor muscles. The present study showed
8 that lumbar SE and WE significantly improved chronic LBP. Both WE and stabilization
9 with WE significantly improved muscular endurance of back muscles. Moreover, walking
10 and SEs also improved the core stability. It is also worth noting that patients in the WE and
11 SE groups were much more compliant than those in the other exercise groups. This study
12 suggests that lumbar SE and WE should be recommended to patients with chronic LBP
13 because they help not only to relieve back pain but also to prevent chronic back pain
14 through the improvement of muscle endurance.

15
16 Many clinical practice guidelines recommend similar approaches for the assessment and
17 management of low back pain. Recommendations include use of a biopsychosocial
18 framework to guide management with initial non-pharmacological treatment, including
19 education that supports self-management and resumption of normal activities and exercise,
20 and psychological programs for those with persistent symptoms (Foster et al., 2018). Jones
21 et al. (2020) discusses the use of pain education with therapeutic exercise to address the
22 psychosocial aspects that are associated with chronic low back pain. Pain education is the
23 umbrella term utilized to encompass any type of education to the patient about their chronic
24 pain. Therapeutic exercise in combination with pain education may allow for more well-
25 rounded and effective treatment for patients with chronic nonspecific low back pain (NS-
26 LBP). They summarized key findings: A thorough literature review yielded 8 studies
27 potentially relevant to the clinical question, and 3 studies that met the inclusion criteria
28 were included. The 3 studies included reports that exercise therapy reduced symptoms.
29 Two of the 3 included studies support the claim that exercise therapy reduces the symptoms
30 of chronic NS-LBP when combined with pain education, whereas one study found no
31 difference between pain education with therapeutic exercise. Authors concluded that there
32 is moderate evidence to support the use of pain education along with therapeutic exercise
33 when attempting to reduce symptoms of pain and disability in patients with chronic NS-
34 LBP. Educational interventions should be created to educate patients about the foundation
35 of pain, and pain education should be implemented as a part of the clinician's strategy for
36 the rehabilitation of patients with chronic NS-LBP.

37
38 Owen et al. (2020) examined the effectiveness of specific modes of exercise training in
39 non-specific chronic low back pain (NSCLBP). They included exercise training
40 randomized controlled/clinical trials in adults with NSCLBP. Among 9,543 records, 89
41 studies (patients=5,578) were eligible for qualitative synthesis and 70 (pain), 63 (physical
42 function), 16 (mental health) and 4 (trunk muscle strength) for Network Meta-analysis

1 (NMA). The NMA consistency model revealed that the following exercise training
 2 modalities had the highest probability of being best when compared with true control:
 3 Pilates for pain, resistance and stabilization/motor control for physical function, and
 4 resistance and aerobic for mental health. Stretching and McKenzie exercise effect sizes did
 5 not differ to true control for pain or function. NMA was not possible for trunk muscle
 6 endurance or analgesic medication. Authors concluded there is low quality evidence that
 7 Pilates, stabilization/motor control, resistance training and aerobic exercise training are the
 8 most effective treatments, pending outcome of interest, for adults with NSCLBP. Exercise
 9 training may also be more effective than therapist hands-on treatment. Heterogeneity
 10 among studies and the fact that there are few studies with low risk of bias are both
 11 limitations. Hayden et al. (2020) sought to determine which individuals might benefit the
 12 most from exercise for their low back pain. For studies included in this analysis, compared
 13 with no treatment/usual care, exercise therapy on average reduced pain, a result compatible
 14 with a clinically important 20% smallest worthwhile effect. Exercise therapy reduced
 15 functional limitations with a clinically important 23% improvement at short-term follow-
 16 up. Not having heavy physical demands at work and medication use for low back pain were
 17 potential treatment effect modifiers that were associated with superior exercise outcomes
 18 relative to non-exercise comparisons. Lower body mass index was also associated with
 19 better outcomes in exercise compared with no treatment/usual care. This study was limited
 20 by inconsistent availability and measurement of participant characteristics.

21
 22 Zhu et al. (2020) compared the effects of yoga for patients with chronic low back pain on
 23 pain, disability, quality of life with non-exercise (e.g., usual care, education), physical
 24 therapy exercise. A total of 18 randomized controlled trials were included in this meta-
 25 analysis. Yoga could significantly reduce pain at 4 to 8 weeks, 3 months, 6 to 7 months,
 26 and was not significant in 12 months compared with non-exercise. Yoga was better than
 27 non-exercise on disability at 4 to 8 weeks, 3 months, 6 months, 12 months. There was no
 28 significant difference on pain, disability compared with physical therapy exercise group.
 29 Furthermore, it suggested that there was a non-significant difference on physical and
 30 mental quality of life between yoga and any other interventions. Authors concluded that
 31 yoga might decrease pain from short term to intermediate term and improve functional
 32 disability status from short term to long term compared with non-exercise (e.g., usual care,
 33 education). Yoga had the same effect on pain and disability as any other exercise or
 34 physical therapy. Yoga might not improve the physical and mental quality of life based on
 35 the result of merging the 36-item short form health survey (SF-36) and the 12 item short
 36 form health survey (SF-12) data.

37
 38 Karlsson et al. (2020) assessed the overall certainty of evidence for the effects of exercise
 39 therapy, compared with other interventions, on pain, disability, recurrence, and adverse
 40 effects in adult patients with acute low back pain within a systematic review. Twenty-four
 41 reviews were included, in which 21 randomized controlled trials ($n = 2,685$) presented data
 42 for an acute population, related to 69 comparisons. Overlap was high, 76%, with a

1 corrected covered area of 0.14. Methodological quality varied from low to high. Exercise
2 therapy was categorized into general exercise therapy, stabilization exercise, and
3 McKenzie therapy. No important difference in pain or disability was evident when exercise
4 therapy was compared with sham ultrasound, nor for the comparators of usual care, spinal
5 manipulative therapy, advice to stay active, and educational booklet. Neither McKenzie
6 therapy nor stabilization exercise yielded any important difference in effects compared
7 with other types of exercise therapy. Certainty of evidence varied from very low to
8 moderate. Authors concluded that these findings suggest very low to moderate certainty of
9 evidence that exercise therapy may result in little or no important difference in pain or
10 disability, compared with other interventions, in adult patients with acute low back pain.

11
12 Skelly et al. (2020) updated the evidence from their 2018 report assessing persistent
13 improvement in outcomes following completion of therapy for noninvasive
14 nonpharmacological treatment for selected chronic pain conditions. They included 233
15 RCTs (31 new to this update). Many were small ($N < 70$), and evidence beyond 12 months
16 after treatment completion was sparse. The most common comparison was with usual care.
17 Evidence on harms was limited, with no evidence suggesting increased risk for serious
18 treatment-related harms for any intervention. Effect sizes were generally small for function
19 and pain. For chronic low back pain, function improved over short and/or intermediate
20 term for exercise (SOE moderate at short term for exercise). Improvements in pain at short
21 term were seen for exercise (SOE: low). At intermediate term, exercise (SOE: low) were
22 associated with improved pain. Compared with exercise, multidisciplinary rehabilitation
23 improved both function and pain at short and intermediate terms (small effects, SOE:
24 moderate).

25
26 Hayden et al. (2021a) assessed the impact of exercise treatment on pain and functional
27 limitations in adults with chronic non-specific low back pain compared to no treatment,
28 usual care, placebo and other conservative treatments in a Cochrane review. The review
29 includes data for trials identified in searches up to 27 April 2018. Authors included
30 randomized controlled trials that assessed exercise treatment compared to no treatment,
31 usual care, placebo or other conservative treatment on the outcomes of pain or functional
32 limitations for a population of adult participants with chronic non-specific low back pain
33 of more than 12 weeks' duration. They included 249 trials of exercise treatment, including
34 studies conducted in Europe (122 studies), Asia (38 studies), North America (33 studies),
35 and the Middle East (24 studies). Sixty-one per cent of studies (151 trials) examined the
36 effectiveness of two or more different types of exercise treatment, and 57% (142 trials)
37 compared exercise treatment to a non-exercise comparison treatment. Study participants
38 had a mean age of 43.7 years and, on average, 59% of study populations were female. Most
39 of the trials were judged to be at risk of bias, including 79% at risk of performance bias
40 due to difficulty blinding exercise treatments. Authors found moderate-certainty evidence
41 that exercise treatment is more effective for treatment of chronic low back pain compared
42 to no treatment, usual care, or placebo comparisons for pain outcomes at earliest follow-

1 up, a clinically important difference. Certainty of evidence was downgraded mainly due to
2 heterogeneity. For the same comparison, there was moderate-certainty evidence for
3 functional limitations outcomes; this finding did not meet the prespecified threshold for
4 minimal clinically important difference. Certainty of evidence was downgraded mainly due
5 to some evidence of publication bias. Compared to all other investigated conservative
6 treatments, exercise treatment was found to have improved pain and functional limitations
7 outcomes. These effects did not meet the prespecified threshold for clinically important
8 difference. Subgroup analysis of pain outcomes suggested that exercise treatment is
9 probably more effective than education alone or non-exercise physical therapy, but with
10 no differences observed for manual therapy. In studies that reported adverse effects (86
11 studies), one or more adverse effects were reported in 37 of 112 exercise groups (33%) and
12 12 of 42 comparison groups (29%). Twelve included studies reported measuring adverse
13 effects in a systematic way, with a median of 0.14 per participant in the exercise groups
14 (mostly minor harms, e.g., muscle soreness), and 0.12 in comparison groups. Authors
15 concluded that moderate-certainty evidence exists that exercise is probably effective for
16 treatment of chronic low back pain compared to no treatment, usual care or placebo for
17 pain. The observed treatment effect for the exercise compared to no treatment, usual care
18 or placebo comparisons is small for functional limitations, not meeting the threshold for
19 minimal clinically important difference. They also found exercise to have improved pain
20 (low-certainty evidence), and functional limitations outcomes (moderate-certainty
21 evidence) compared to other conservative treatments; however, these effects were small
22 and not clinically important when considering all comparisons together. Subgroup analysis
23 suggested that exercise treatment is probably more effective than advice or education alone,
24 or electrotherapy, but with no differences observed for manual therapy treatments. Hayden
25 et al. (2021b) wanted to investigate what the effects of specific types of exercise treatments
26 on pain intensity and functional limitation outcomes for adults with chronic low back pain
27 are in a systematic review with network meta-analysis of randomized controlled trials.
28 Exercise treatments prescribed or planned by a health professional that involved
29 conducting specific activities, postures and/or movements with a goal to improve low back
30 pain outcomes were included in the review. Outcome measures included pain intensity
31 (e.g., visual analogue scale or numerical rating scale) and back-related functional
32 limitations (e.g., Roland Morris Disability Questionnaire or Oswestry Disability Index),
33 each standardized to range from 0 to 100. This review included 217 randomized controlled
34 trials with 20,969 participants and 507 treatment groups. Most exercise types were more
35 effective than minimal treatment for pain and functional limitation outcomes. Network
36 meta-analysis results were compatible with moderate to clinically important treatment
37 effects for Pilates, McKenzie therapy, and functional restoration (pain only) and flexibility
38 exercises (function only) compared with minimal treatment, other effective treatments, and
39 other exercise types. This review found evidence that Pilates, McKenzie therapy and
40 functional restoration were more effective than other types of exercise treatment for
41 reducing pain intensity and functional limitations. Nevertheless, people with chronic low

1 back pain should be encouraged to perform the exercise that they enjoy to promote
2 adherence.

3
4 Thorton et al. (2021) summarized the evidence for non-pharmacological management of
5 low back pain (LBP) in athletes, a common problem in sport that can negatively impact
6 performance and contribute to early retirement. Among 1,629 references, 14 randomized
7 controlled trials (RCTs) involving 541 athletes were included. The trials had biases across
8 multiple domains including performance, attrition, and reporting. Treatments included
9 exercise, biomechanical modifications, and manual therapy. There were no trials
10 evaluating the efficacy of surgery or injections. Exercise was the most frequently
11 investigated treatment; no RTS data were reported for any exercise intervention. There was
12 a reduction in pain and disability reported after all treatments. Authors concluded that while
13 several treatments for LBP in athletes improved pain and function, it was unclear what the
14 most effective treatments were, and for whom. Exercise approaches generally reduced pain
15 and improved function in athletes with LBP, but the effect on RTS is unknown. No
16 conclusions regarding the value of manual therapy (massage, spinal manipulation) or
17 biomechanical modifications alone could be drawn because of insufficient evidence. High-
18 quality RCTs are urgently needed to determine the effect of commonly used interventions
19 in treating LBP in athletes. Quentin et al. (2021) conducted a systemic review and meta-
20 analysis on the effects of home-based exercise on pain and functional limitation in LBP.
21 They included 33 studies and 9,588 patients. They found that pain intensity decreased in
22 the exclusive home exercise group in the group which conducted exercise both at-home
23 and at another setting. Similarly, functional limitation also decreased in both groups.
24 Relaxation and postural exercise seemed to be ineffective in decreasing pain intensity,
25 whereas trunk, pelvic or leg stretching decreased pain intensity. Yoga improved functional
26 limitation.

27
28 Supervised training was the most effective method to improve pain intensity. Insufficient
29 data precluded robust conclusions around the duration and frequency of the sessions and
30 program. Authors concluded that home-based exercise training improved pain intensity
31 and functional limitation parameters in LBP. Van Dillen et al. (2021) sought to determine
32 whether an exercise-based treatment of person-specific motor skill training (MST) in
33 performance of functional activities is more effective in improving function than strength
34 and flexibility exercise (SFE) immediately, 6 months, and 12 months following treatment.
35 The effect of booster treatments 6 months following treatment also was examined. A total
36 of 154 people with at least 12 months of chronic, nonspecific LBP, aged 18 to 60 years,
37 with modified Oswestry Disability Questionnaire (MODQ) score of at least 20% were
38 randomized to either MST or SFE. Data were analyzed between December 1, 2017, and
39 October 6, 2020. Participants received 6 weekly 1-hour sessions of MST in functional
40 activity performance or SFE of the trunk and lower limbs. Half of the participants in each
41 group received up to 3 booster treatments 6 months following treatment. A total of 149
42 participants (91 women; mean [SD] age, 42.5 [11.7] years) received some treatment and

1 were included in the intention-to-treat analysis. Following treatment, MODQ scores were
2 lower for MST than SFE by 7.9 (95% CI, 4.7 to 11.0; $P < .001$). During the follow-up
3 phase, the MST group maintained lower MODQ scores than the SFE group, 5.6 lower at 6
4 months (95% CI, 2.1 to 9.1) and 5.7 lower at 12 months (95% CI, 2.2 to 9.1). Booster
5 sessions did not change MODQ scores in either treatment. Authors concluded that people
6 with chronic LBP who received MST had greater short-term and long-term improvements
7 in function than those who received SFE. Person-specific MST in functional activities
8 limited owing to LBP should be considered in the treatment of people with chronic LBP.

9
10 According to Chou (2021), low back pain is a common problem that is the leading cause
11 of disability and is associated with high costs. Evaluation focuses on identification of risk
12 factors indicating a serious underlying condition and increased risk for persistent disabling
13 symptoms in order to guide selective use of diagnostic testing (including imaging) and
14 treatments. Nonpharmacologic therapies, including exercise and psychosocial
15 management, are preferred for most patients with low back pain and may be supplemented
16 with adjunctive drug therapies. Surgery and interventional procedures are options in a
17 minority of patients who do not respond to standard treatments. Hlaing et al. (2021)
18 compared the effects of two different exercise regimes, Core stabilization exercises (CSE)
19 and strengthening exercise (STE), on proprioception, balance, muscle thickness and pain-
20 related outcomes in patients with subacute non-specific low back pain (NSLBP). Thirty-
21 six subacute NSLBP patients, [mean age, 34.78 ± 9.07 years; BMI, 24.03 ± 3.20 Kg/m²;
22 and duration of current pain, 8.22 ± 1.61 weeks], were included in this study. They were
23 randomly allocated into either CSE ($n = 18$) or STE groups ($n = 18$). Exercise training was
24 given for 30 min, three times per week, for up to 4 weeks. Proprioception, standing balance,
25 muscle thickness of transversus abdominis (TrA) and lumbar multifidus (LM), and pain-
26 related outcomes, comprising pain, functional disability and fear of movement, were
27 assessed at baseline and after 4 weeks of intervention. The CSE group demonstrated
28 significantly more improvement than the STE group after 4 weeks of intervention.
29 Improvements were in: proprioception, balance: single leg standing with eyes open and
30 eyes closed on both stable and unstable surfaces, and percentage change of muscle
31 thickness of TrA and LM. Although both exercise groups gained relief from pain, the CSE
32 group demonstrated greater reduction of functional disability and fear of movement. There
33 were no significant adverse effects in either type of exercise program. Authors concluded
34 that despite both core stabilization and strengthening exercises reducing pain, core
35 stabilization exercise is superior to strengthening exercise. It is effective in improving
36 proprioception, balance, and percentage change of muscle thickness of TrA and LM, and
37 reducing functional disability and fear of movement in patients with subacute NSLBP.

38
39 Rathnayake et al. (2021) systematically reviewed the evidence for the effect of self-
40 management interventions (SMIs) with an exercise component added on pain and disability
41 in people with CLBP. Authors concluded that there is low-quality evidence that SMIs with
42 exercises added have moderately positive effects on pain and disability in patients with

1 CLBP compared to control interventions involving usual care, typically consisting of
2 access to medication, exercise, advice, education, and manual therapy.

3
4 Drummond et al. (2021) assessed the effectiveness of sling exercise therapy (SET) in
5 individuals with chronic low back pain (LBP). The search identified 1,204 studies, with 12
6 studies meeting the inclusion criteria. Meta-analysis comparing SET with general exercise
7 revealed a nonsignificant effect for pain. Meta-analysis comparing SET with motor control
8 training/lumbar stabilization revealed a significant effect favoring SET for pain and
9 disability. Meta-analysis comparing SET with no treatment revealed a significant effect
10 favoring SET for pain. Meta-analysis comparing SET plus modalities with modalities
11 revealed a significant effect favoring the SET plus modalities group for pain and a
12 nonsignificant effect for disability. Sling exercise therapy was more effective than all
13 comparisons for various muscle attributes. The overall level of evidence ranged from very
14 low to moderate. Sling exercise therapy is effective in reducing pain, disability, and
15 improving core muscle activation, strength, thickness, and onset in patients with chronic
16 LBP. Because SET demonstrated comparable outcomes with common active interventions,
17 it provides an opportunity to implement pain-free exercises based on the patient's initial
18 functional level early in the plan of care.

19
20 Ferreira et al. (2021) assessed whether an exercise and education program was more
21 effective than an education booklet for preventing recurrence of low back pain (LBP).
22 Participants aged 18 years or older who had recovered from an episode of LBP within the
23 previous week were recruited from primary care practices and the community. Participants
24 were randomized to receive either 12 weeks of exercise and education (8 supervised
25 exercise sessions and 3 one-on-one sessions) or a control (education booklet). The primary
26 outcome was time to recurrence of LBP during the 1-year follow-up. Times to recurrence
27 of LBP leading to activity limitation, care seeking, and work absence were secondary
28 outcomes. Data were analyzed with Cox regression using intention-to-treat principles. The
29 same size was 111 (exercise and education, $n = 57$; educational booklet, $n = 54$). At the
30 end of the study period, data completeness was 84.2%. Thirty-six (63%) participants in the
31 exercise and education group and 31 (57%) participants in the control group had a
32 recurrence of LBP. There was no statistically significant difference in time to recurrence
33 of pain between groups (hazard ratio = 1.09; 95% confidence interval: 0.7, 1.8). There was
34 no statistically significant effect for any of the secondary outcomes. Authors concluded
35 that among people recently recovered from LBP, exercise and education may not
36 meaningfully reduce risk of recurrence compared to providing an educational booklet.

37
38 Burns et al. (2021) determined whether adding hip treatment to usual care for low back
39 pain (LBP) improved disability and pain in individuals with LBP and a concurrent hip
40 impairment. Seventy-six participants (age, 18 years or older; Oswestry Disability Index,
41 20% or greater; numeric pain-rating scale, 2 or more points) with LBP and a concurrent
42 hip impairment were randomly assigned to a group that received treatment to the lumbar

1 spine only (LBO group) ($n = 39$) or to one that received both lumbar spine and hip
 2 treatments (LBH group) ($n = 37$). The individual treating clinicians decided which specific
 3 low back treatments to administer to the LBO group. Treatments aimed at the hip (LBH
 4 group) included manual therapy, exercise, and education, selected by the therapist from a
 5 predetermined set of treatments. Primary outcomes were disability and pain, measured by
 6 the Oswestry Disability Index and the numeric pain-rating scale, respectively, at baseline,
 7 2 weeks, discharge, 6 months, and 12 months. The secondary outcomes were fear-
 8 avoidance beliefs (work and physical activity subscales of the Fear-Avoidance Beliefs
 9 Questionnaire), global rating of change, the Patient Acceptable Symptom State, and
 10 physical activity level. Investigators used mixed-model 2-by-3 analyses of variance to
 11 examine group-by-time interaction effects (intention-to-treat analysis). Data were available
 12 for 68 patients at discharge (LBH group, $n = 33$; LBO group, $n = 35$) and 48 at 12 months
 13 ($n = 24$ for both groups). There were no between-group differences in disability at
 14 discharge, 12 months, and all other time points. There were no between-group differences
 15 in pain at discharge, 12 months, and all other time points. There were no between-group
 16 differences in secondary outcomes, except for higher Fear-Avoidance Beliefs
 17 Questionnaire (work subscale) scores in the LBH group at 2 weeks and discharge. Authors
 18 concluded that adding treatments aimed at the hip to usual low back physical therapy did
 19 not provide additional short- or long-term benefits in reducing disability and pain in
 20 individuals with LBP and a concurrent hip impairment. Clinicians may not need to include
 21 hip treatments to achieve reductions in low back disability and pain in individuals with
 22 LBP and a concurrent hip impairment.

23
 24 Nava-Bringas et al. (2021) compared the effectiveness of lumbar stabilization exercises
 25 and flexion exercises for pain control and improvements of disability in individuals with
 26 chronic low back pain (CLBP) and degenerative spondylolisthesis (DS). A randomized
 27 controlled trial was conducted in a tertiary public hospital and included 92 individuals over
 28 the age of 50 years who were randomly allocated to lumbar stabilization exercises or
 29 flexion exercises. Participants received 6 sessions of physical therapy (monthly
 30 appointments) and were instructed to execute exercises daily at home during the 6 months
 31 of the study. The primary outcome (measured at baseline, 1 month, 3 months, and 6
 32 months) was pain intensity (visual analog scale, 0-100 mm) and disability (Oswestry
 33 Disability Index, from 0% to 100%). Secondary outcomes were disability (Roland-Morris
 34 Disability Questionnaire, from 0 to 24 points), changes in body mass index, and flexibility
 35 (fingertip to floor, in centimeters) at baseline and 6 months, and also the total of days of
 36 analgesic use at 6-month follow-up. Mean differences between groups were not significant
 37 for lumbar pain, radicular pain, for Oswestry scores, and for Roland Morris scores. Authors
 38 state that the findings from the present study reveal that flexion exercises are not inferior
 39 to and offer a similar response to stabilization exercises for the control of pain and
 40 improvements of disability in individuals with CLBP and DS.

1 De Campos et al. (2021) evaluate the evidence from randomized controlled trials (RCTs)
 2 on the effectiveness of prevention strategies to reduce future impact of low back pain
 3 (LBP), where impact is measured by LBP intensity and associated disability. 27 published
 4 reports of 25 different trials including a total of 8341 participants fulfilled the inclusion
 5 criteria. The pooled results, from three RCTs (612 participants), found moderate-quality
 6 evidence that an exercise program can prevent future LBP intensity, and from 4 RCTs (471
 7 participants) that an exercise and education program can prevent future disability due to
 8 LBP. It is uncertain whether prevention programs improve future quality of life (QoL) and
 9 workability due to the overall low-quality and very low-quality available evidence. Authors
 10 concluded that this review provides moderate-quality evidence that an exercise program,
 11 and a program combining exercise and education, are effective to reduce future LBP
 12 intensity and associated disability. It is uncertain whether prevention programs can
 13 improve future QoL and workability. Further high-quality RCTs evaluating prevention
 14 programs aiming to reduce future impact of LBP are needed.

15
 16 George et al. (2021) updated a clinical practice guideline for treatment of low back pain.
 17 Findings relative to exercise included the following:

- 18 • Exercise For Acute Low Back Pain
 - 19 ○ Physical therapists can use exercise training interventions, including
 - 20 specific trunk muscle activation, for patients with acute low back pain
 - 21 (LBP) (grade C).
- 22 • Exercise For Acute Low Back Pain With Leg Pain
 - 23 ○ Physical therapists may use exercise training interventions, including trunk
 - 24 muscle strengthening and endurance and specific trunk muscle activation,
 - 25 to reduce pain and disability for patients with acute LBP with leg pain
 - 26 (grade B).
- 27 • Exercise For Chronic Low Back Pain
 - 28 ○ Physical therapists should use exercise training interventions, including
 - 29 trunk muscle strengthening and endurance, multimodal exercise
 - 30 interventions, specific trunk muscle activation exercise, aerobic exercise,
 - 31 aquatic exercise, and general exercise, for patients with chronic LBP (grade
 - 32 A).
 - 33 ○ Physical therapists may provide movement control exercise or trunk
 - 34 mobility exercise for patients with chronic LBP (grade B).
- 35 • Exercise For Chronic Low Back Pain With Leg Pain
 - 36 ○ Physical therapists may use exercise training interventions, including
 - 37 specific trunk muscle activation and movement control, for patients with
 - 38 chronic LBP with leg pain (grade B).
- 39 • Exercise For Chronic Low Back Pain With Movement Control Impairment
 - 40 ○ Physical therapists should use specific trunk muscle activation and
 - 41 movement control exercise for patients with chronic LBP and movement
 - 42 control impairment (grade A).

- 1 • Exercise For Chronic Low Back Pain In Older Adults
- 2 ○ Physical therapists should use general exercise training to reduce pain and
- 3 disability in older adults with chronic LBP (grade A).
- 4 • Exercise For Postoperative Low Back Pain
- 5 ○ Physical therapists can use general exercise training for patients with LBP
- 6 following lumbar spine surgery (grade C).

7

8 Gianola et al. (2022) assessed the effectiveness of interventions for acute and subacute non-

9 specific low back pain (NS-LBP) based on pain and disability outcomes in a systematic

10 review with network meta-analysis. Forty-six RCTs ($n=8,765$) were included. At

11 immediate-term follow-up, for pain decrease, exercise was considered one of the most

12 efficacious treatments against an inert therapy. Similar findings were confirmed for

13 disability. Fernández-Rodríguez et al. (2022) sought to determine which type of exercise

14 is best for reducing pain and disability in adults with chronic low back pain (LBP) in a

15 systematic review with a network meta-analysis (NMA) of randomized controlled trials

16 (RCTs). Authors included 118 trials (9,710 participants). There were 28 head-to-head

17 comparisons, 7 indirect comparisons for pain, and 8 indirect comparisons for disability.

18 Compared with control, all types of physical exercises were effective for improving pain

19 and disability, except for stretching exercises (for reducing pain) and the McKenzie method

20 (for reducing disability). The most effective interventions for reducing pain were Pilates,

21 mind-body, and core-based exercises. The most effective interventions for reducing

22 disability were Pilates, strength, and core-based exercises. On SUCRA analysis, Pilates

23 had the highest likelihood for reducing pain (93%) and disability (98%). Authors concluded

24 that although most exercise interventions had benefits for managing pain and disability in

25 chronic LBP, the most beneficial programs were those that included (1) at least 1 to 2

26 sessions per week of Pilates or strength exercises; (2) sessions of less than 60 minutes of

27 core-based, strength, or mind-body exercises; and (3) training programs from 3 to 9 weeks

28 of Pilates and core-based exercises.

29

30 Grooten et al. (2022) aimed to identify systematic reviews of common exercise types used

31 in CLBP, to appraise their quality, and to summarize and compare their effect on pain and

32 disability. The included reviews were grouped into nine exercise types: aerobic training,

33 aquatic exercises, motor control exercises (MCE), resistance training, Pilates, sling

34 exercises, traditional Chinese exercises (TCE), walking, and yoga. Out of the 253 full texts

35 that were screened, we included 45 systematic reviews and meta-analyses. The quality of

36 the included reviews ranged from high to critically low. Due to large heterogeneity, no

37 meta-analyses were performed. Authors found low-to-moderate evidence of mainly short-

38 term and small beneficial effects on pain and disability for MCE, Pilates, resistance

39 training, TCE, and yoga compared to no or minimal intervention. Authors conclude that

40 findings show that the effect of various exercise types used in CLBP on pain and disability

41 varies with no major difference between exercise types. Essman and Lin (2022) highlighted

42 the role of exercise in preventing and managing acute and chronic axial low back pain

1 (LBP). They note that no single exercise method has been shown to be more effective than
2 another. Overall, their review summarizes the beneficial role of a personalized exercise
3 program and related counseling strategies in the prevention and management of LBP.

4
5 Bagg et al. (2022) estimated the effect of a graded sensorimotor retraining intervention
6 (RESOLVE) on pain intensity in people with chronic low back pain. This parallel, 2-group,
7 randomized clinical trial recruited participants with chronic (>3 months) nonspecific low
8 back pain from primary care and community settings. A total of 276 adults were
9 randomized (in a 1:1 ratio) to the intervention or sham procedure and attention control
10 groups delivered by clinicians at a medical research institute in Sydney, Australia.
11 Participants randomized to the intervention group ($n = 138$) were asked to participate in 12
12 weekly clinical sessions and home training designed to educate them about and assist them
13 with movement and physical activity while experiencing lower back pain. Participants
14 randomized to the control group ($n = 138$) were asked to participate in 12 weekly clinical
15 sessions and home training that required similar time as the intervention but did not focus
16 on education, movement, and physical activity. The control group included sham laser and
17 shortwave diathermy applied to the back and sham noninvasive brain stimulation. Among
18 276 randomized patients completed follow-up at 18 weeks. The mean pain intensity was
19 5.6 at baseline and 3.1 at 18 weeks in the intervention group and 5.8 at baseline and 4.0 at
20 18 weeks in the control group, with an estimated between-group mean difference at 18
21 weeks of -1.0 point, favoring the intervention group. In this randomized clinical trial
22 conducted at a single center among patients with chronic low back pain, graded
23 sensorimotor retraining, compared with a sham procedure and attention control,
24 significantly improved pain intensity at 18 weeks. The improvements in pain intensity were
25 small, and further research is needed to understand the generalizability of the findings.

26
27 Fleckenstein et al. (2022) investigated the effects of individualized interventions, based on
28 exercise alone or combined with psychological treatment, on pain intensity and disability
29 in patients with chronic non-specific low-back-pain. Fifty-eight studies ($n = 10,084$) were
30 included. At short-term follow-up (12 weeks), low-certainty evidence for pain intensity
31 and very low-certainty evidence for disability indicates effects of individualized versus
32 active exercises, and very low-certainty evidence for pain intensity, but not (low-certainty
33 evidence) for disability compared to passive controls. At long-term follow-up (1 year),
34 moderate-certainty evidence for pain intensity and disability indicates effects versus
35 passive controls. Sensitivity analyses indicates that the effects on pain, but not on disability
36 (always short-term and versus active treatments) were robust. Pain reduction caused by
37 individualized exercise treatments in combination with psychological interventions (in
38 particular behavioral-cognitive therapies) is of clinical importance. Certainty of evidence
39 was downgraded mainly due to evidence of risk of bias, publication bias and inconsistency
40 that could not be explained. Individualized exercise can treat pain and disability in chronic
41 non-specific low-back-pain. The effects at short term are of clinical importance (relative
42 differences versus active 38% and versus passive interventions 77%), especially in regard

1 to the little extra effort to individualize exercise. Sub-group analysis suggests a
2 combination of individualized exercise (especially motor-control based treatments) with
3 behavioral therapy interventions to booster effects.

4
5 Niederer et al. (2022) investigated how risk of bias and intervention type modify effect
6 sizes of exercise interventions that are intended to reduce chronic low back pain intensity.
7 Potential effect modifiers were risk of bias, exercise modes, study, and meta-analyses
8 characteristics. Data from 26 systematic reviews ($k = 349$ effect sizes, $n = 18,879$
9 participants) were analyzed. There was a clinically relevant effect overestimation in studies
10 with a high risk of bias due to missing outcomes and low sample size. There was a clinically
11 relevant underestimation of the effect when studies were at high risk of bias and outcome
12 measurement. Motor control and stabilization training had the largest effects; stretching
13 had the smallest effect. Authors concluded that the effects of exercise trials at high risk of
14 bias may be overestimated or underestimated. After accounting for risk of bias, motor
15 control and stabilization exercises may represent the most effective exercise therapies for
16 chronic low back pain. Cashin et al. (2022) aimed to synthesize and appraise the current
17 research to provide practical, evidence-based guidance concerning exercise prescription
18 for non-specific CLBP. Systematic reviews show exercise is effective for small, short-term
19 reductions in pain and disability, when compared with placebo, usual care, or waiting list
20 control, and serious adverse events are rare. A range of individualized or group-based
21 exercise modalities have been demonstrated as effective in reducing pain and disability.
22 Authors conclude that to promote recovery, sustainable outcomes and self-management,
23 exercise can be coupled with education strategies, as well as interventions that enhance
24 adherence, motivation, and patient self-efficacy.

25
26 García-Moreno et al. (2022) upgraded the evidence of the most effective preventive
27 physiotherapy interventions to improve back care in children and adolescents. Twenty
28 studies were finally included. The most common physiotherapy interventions were
29 exercise, postural hygiene, and physical activity. The mean age of the total sample was
30 11.79 years. Authors concluded that recent studies provide strong support for the use of
31 physiotherapy in the improvement of back care and prevention of non-specific low back
32 pain in children and adolescents. Based on GRADE methodology, they found that the
33 evidence was from very low to moderate quality and interventions involving physical
34 exercise, postural hygiene and physical activity should be preferred. Lindberg and Leggit
35 (2022) summarized that there is low- to moderate-quality evidence that exercise reduces
36 pain and improves function in patients with chronic low back pain compared with no
37 treatment, usual care, and other conservative interventions such as education, manual
38 therapy, and electrotherapy. This effect is clinically significant in the short term (six to 12
39 weeks) but less pronounced six months after treatment completion. The review does not
40 recommend a specific exercise regimen to treat chronic low back pain.

1 Prat-Luri et al. (2023) analyzed the effect of trunk-focused exercise programs (TEPs) and
2 moderator factors on chronic nonspecific low back pain (LBP). Forty randomized
3 controlled trials (n = 2,391) were included. TEPs showed positive effects for all outcomes
4 versus control. There were small effects in favor of TEPs versus general exercises for pain
5 and disability. Trunk and/or hip range-of-motion improvements were associated with
6 greater reductions in pain and disability. Low body mass was associated with higher pain
7 reduction. Authors concluded that trunk-focused exercise programs had positive effects on
8 pain, disability, quality of life, and trunk performance compared to control groups, and on
9 pain and disability compared to general exercises. Increasing trunk and/or hip range of
10 motion was associated with greater pain and disability reduction, and lower body mass
11 with higher pain reduction.

12
13 Ijzelenberg et al. (2023) evaluated the benefits and harms of exercise therapy for acute non-
14 specific low back pain in adults compared to sham/placebo treatment or no treatment at
15 short-term, intermediate-term, and long-term follow-up. This is an update of a Cochrane
16 Review first published in 2005. Authors included RCTs that examined the effects of
17 exercise therapy on non-specific LBP lasting six weeks or less in adults. Major outcomes
18 for this review were pain, functional status, and perceived recovery. Minor outcomes were
19 return to work, health-related quality of life, and adverse events. Main comparisons were
20 exercise therapy versus sham/placebo treatment and exercise therapy versus no treatment.
21 Outcomes were evaluated at short-term follow-up (time point within three months and
22 closest to six weeks after randomization; main follow-up), intermediate-term follow-up
23 (between nine months and closest to six months), and long-term follow-up (after nine
24 months and closest to 12 months). Authors included 23 studies (13 from the previous
25 review, 10 new studies) that involved 2,674 participants and provided data for 2,637
26 participants. Included studies were conducted in Europe ($N = 9$), the Asia-Pacific region
27 ($N = 9$), and North America ($N = 5$); and most took place in a primary care setting ($N =$
28 12), secondary care setting ($N = 6$), or both ($N = 1$). In most studies, the population was
29 middle-aged and included men and women. They judged 10 studies (43%) at low risk of
30 bias with regard to sequence generation and allocation concealment. There is very low-
31 certainty evidence that exercise therapy compared with sham/placebo treatment has no
32 clinically relevant effect on pain scores in the short term. There is very low-certainty
33 evidence that exercise therapy compared with sham/placebo treatment has no clinically
34 relevant effect on functional status scores in the short term. There is very low-certainty
35 evidence that exercise therapy compared with no treatment has no clinically relevant effect
36 on pain or functional status in the short term. Owing to insufficient reporting of adverse
37 events, authors were unable to reach any conclusions on the safety or harms related to
38 exercise therapy. Authors concluded that exercise therapy compared to sham/placebo
39 treatment may have no clinically relevant effect on pain or functional status in the short
40 term in people with acute non-specific LBP, but the evidence is very uncertain. Exercise
41 therapy compared to no treatment may have no clinically relevant effect on pain or

1 functional status in the short term in people with acute non-specific LBP, but the evidence
2 is very uncertain.

3
4 Li et al. (2023) evaluated the effects of different exercise therapies on chronic low back
5 pain and provided a reference for exercise regimens in CLBP patients. This study included
6 75 randomized controlled trials (RCTs) with 5,254 participants. Network meta-analysis
7 results showed that tai chi, yoga (SMD, -1.76; 95% CI -2.72 to -0.81), Pilates exercise, and
8 sling exercise showed a better pain improvement than conventional rehabilitation. Tai chi
9 and yoga showed a better pain improvement than no intervention provided. Yoga and core
10 or stabilization exercises showed a better physical function improvement than conventional
11 rehabilitation. Yoga and core or stabilization exercises showed a better physical function
12 improvement than no intervention provided. Authors concluded that compared with
13 conventional rehabilitation and no intervention provided, tai chi, yoga, Pilates exercise,
14 sling exercise, motor control exercise, and core or stabilization exercises significantly
15 improved CLBP in patients. Compared with conventional rehabilitation and no
16 intervention provided, yoga and core or stabilization exercises were statistically significant
17 in improving physical function in patients with CLBP. Due to the limitations of the quality
18 and quantity of the included studies, it is difficult to make a definitive recommendation
19 before more large-scale and high-quality RCTs are conducted.

20
21 Kazeminia et al. (2023) aimed to estimate the results of randomized clinical trials (RCT)
22 about the effect of pelvic floor muscle-strengthening exercises on reducing low back pain.
23 Nineteen RCTs with a sample size of 456 subjects in the intervention group and 470 in the
24 control group were included in the meta-analysis. Authors concluded that based on the
25 results of the present meta-analysis, pelvic floor muscle-strengthening exercises
26 significantly reduce the low back pain intensity. Therefore, these exercises can be regarded
27 as a part of a low back pain management plan.

28
29 Wong et al. (2023) compared the effects of Pilates exercise (PE) with other forms of
30 exercise on pain and disability in individuals with chronic non-specific low back pain
31 (CNSLBP) and to inform clinical practice and future research. Eleven RCTs were included.
32 A low certainty of evidence supported PE was more effective than general exercise (GE)
33 in pain reduction. Moreover, very low levels of certainty were revealed for effectiveness
34 of PE compared with direction-specific exercise (DSE) for pain reduction and equivalence
35 of PE and spinal stabilization exercise (SSE) for pain and disability. Authors concluded
36 that their review found no strong evidence for using one type of exercise intervention over
37 another when managing patients with CNSLBP. Existing evidence does not allow this
38 review to draw definitive recommendations. In the absence of a superior exercise form
39 clinicians should work collaboratively with the patient, using the individual's goals and
40 preferences to guide exercise selection. Further appropriately designed research is
41 warranted to explore this topic further.

1 Zaina et al. (2023) identified evidence-based rehabilitation interventions for persons with
2 non-specific low back pain (LBP) with and without radiculopathy and developed
3 recommendations from high-quality clinical practice guidelines (CPGs) to inform the
4 World Health Organization's (WHO) Package of Interventions for Rehabilitation (PIR).
5 Four high-quality CPGs were identified. Recommended interventions included (1)
6 education about recovery expectations, self-management strategies, and maintenance of
7 usual activities; (2) multimodal approaches incorporating education, exercise, and spinal
8 manipulation; (3) nonsteroidal anti-inflammatory drugs combined with education in the
9 acute stage; and (4) intensive interdisciplinary rehabilitation that includes exercise and
10 cognitive/behavioral interventions for persistent pain. No high-quality CPGs for people
11 younger than 16 years of age were found. Authors concluded that these recommendations
12 emphasize the potential benefits of education, exercise, manual therapy, and
13 cognitive/behavioral interventions.

14
15 Gilliam et al. (2023) assessed the effectiveness of mind-body (MB) exercise interventions
16 provided by physical therapists for reducing pain and disability in people with low back
17 pain (LBP). Randomized controlled trials evaluating the effects of Pilates, yoga, and tai chi
18 interventions performed by physical therapists on pain or disability outcomes in adults with
19 musculoskeletal LBP were included. Eight trials, 7 reporting on Pilates and 1 reporting on
20 yoga, were included. Short-term outcomes for pain and indicated MB exercise was more
21 effective than control intervention. Tests for subgroup differences between studies with
22 exercise vs non-exercise control groups revealed a moderating effect on short-term
23 outcomes where larger effects were observed in studies with non-exercise comparators.
24 Long-term outcomes for pain and disability suggested that MB exercise is not more
25 effective than control interventions for pain or disability. Quality of the evidence ranged
26 from very low to low. Authors concluded that physical therapist-delivered MB exercise
27 interventions, which overwhelmingly consisted of Pilates, were more effective than control
28 in the short and long-term for pain and in the short-term for disability, with differences in
29 the short-term effects lessened when compared with an active intervention. Pilates
30 interventions delivered by physical therapists represent a viable tool for the clinical
31 management of chronic LBP.

32
33 Ram et al. (2023) determined the effect of higher versus lower intensity exercise intensity
34 on pain, disability, quality of life and adverse events in people with CLBP. Four trials (n =
35 214 participants, 84% male) reported across five studies were included. Higher intensity
36 exercise reduced disability more than lower intensity exercise at end-treatment but not at
37 6-month follow-up. Higher intensity exercise did not reliably improve pain and quality of
38 life more than lower intensity exercise. Adverse events did not differ between exercise
39 intensities. All studies were at high risk of bias. Based on very low certainty evidence from
40 a limited number of studies, exercise intensity does not appear to meaningfully influence
41 clinical outcomes in people with CLBP.

1 Almeida et al. (2023) evaluated the effectiveness of the McKenzie method in people with
 2 (sub)acute non-specific low back pain in a Cochrane review. This review included five
 3 RCTs with a total of 563 participants recruited from primary or tertiary care. Three trials
 4 were conducted in the USA, one in Australia, and one in Scotland. Three trials received
 5 financial support from non-commercial funders and two did not provide information on
 6 funding sources. All trials were at high risk of performance and detection bias. None of the
 7 included trials measured adverse events. McKenzie method versus minimal intervention
 8 (educational booklet; McKenzie method as a supplement to other intervention - main
 9 comparison): There is low-certainty evidence that the McKenzie method may result in a
 10 slight reduction in pain in the short term but not in the intermediate term. There is low-
 11 certainty evidence that the McKenzie method may not reduce disability in the short term
 12 nor in the intermediate term. McKenzie method versus manual therapy: There is low-
 13 certainty evidence that the McKenzie method may not reduce pain in the short term and
 14 may result in a slight increase in pain in the intermediate term. There is low-certainty
 15 evidence that the McKenzie method may not reduce disability in the short term nor in the
 16 intermediate term. McKenzie method versus other interventions (massage and advice):
 17 There is very low-certainty evidence that the McKenzie method may not reduce disability
 18 in the short term nor in the intermediate term. Authors concluded that based on low- to
 19 very low-certainty evidence, the treatment effects for pain and disability found in our
 20 review were not clinically important. Thus, they can conclude that the McKenzie method
 21 is not an effective treatment for (sub)acute NSLBP.

22
 23 Gilanyi et al. (2023) determined the effect of exercise on pain self-efficacy in adults with
 24 nonspecific chronic low back pain (NSCLBP). Authors included randomized controlled
 25 trials that compared the effect of exercise on pain self-efficacy to control, in adults with
 26 NSCLBP. Seventeen trials were included, of which eight ($n = 1,121$ participants; 60.6%
 27 female; mean age: 49.6 years) were included in the meta-analysis. Exercise increased pain
 28 self-efficacy by 3.02 points on the 60-point Pain Self-Efficacy Questionnaire. The certainty
 29 of evidence was moderate; all trials were at high risk of bias. Authors concluded that there
 30 was moderate-certainty evidence that exercise increased pain self-efficacy in adults with
 31 NSCLBP.

32
 33 Santos et al. (2023) evaluated the efficacy of Pilates on pain, functional disorders, and
 34 quality of life in patients with chronic low back pain (CLBP). Nineteen randomized
 35 controlled trials with a total of 1108 patients were included. Compared with the controls,
 36 this meta-analysis revealed that Pilates may have positive efficacy for pain relief and the
 37 improvement of functional disorders in CLBP patients, but the improvement in quality of
 38 life seems to be less obvious.

39
 40 Verville et al. (2023) evaluated benefits and harms of structured exercise programs for
 41 chronic primary low back pain (CPLBP) in adults to inform a World Health Organization
 42 (WHO) standard clinical guideline. Thirteen RCTs rated with overall low or unclear risk

1 of bias were synthesized. Assessing individual exercise types (predominantly very low
2 certainty evidence), pain reduction was associated with aerobic exercise and Pilates vs. no
3 intervention, and motor control exercise vs. sham. Improved function was associated with
4 mixed exercise vs. usual care, and Pilates vs. no intervention. Temporary increased minor
5 pain was associated with mixed exercise vs. no intervention, and yoga vs. usual care. Little
6 to no difference was found for other comparisons and outcomes. When pooling exercise
7 types, exercise vs. no intervention probably reduces pain in adults and functional
8 limitations in adults and older adults (moderate certainty evidence). Authors concluded
9 with moderate certainty that structured exercise programs probably reduce pain and
10 functional limitations in adults and older people with CLBP.

11
12 Zhang et al. (2023) compared the efficacy of different exercises therapy on CLBP,
13 dysfunction, quality of life, and mobility in the elderly. Sixteen articles (18 RCTs) were
14 included, comprising a total of 989 participants. The quality of included studies was
15 relatively high. Meta-analysis results indicated that exercise therapy could improve visual
16 analog scale (VAS), Oswestry disability index (ODI), short-form 36-item health survey
17 physical composite summary (SF-36PCS), short-form 36-item health survey mental
18 composite summary (SF-36MCS), and timed up and go test (TUG). Exercise therapy
19 effectively improved VAS, ODI, and SF-36 indexes in the elderly. Based on the subgroup,
20 when designing the exercise therapy regimen, aerobics, strength, and mind-body exercise
21 (≥ 12 weeks, ≥ 3 times/week, ≥ 60 min) should be considered carefully, to ensure the safety
22 and effectiveness for the rehabilitation of CLBP patients.

23
24 Roren et al. (2023) critically reviewed available evidence regarding the efficacy of physical
25 activity for people with LBP. They reported that in acute and subacute LBP, exercise did
26 not reduce pain compared to no exercise. In chronic low back pain (CLBP), exercise
27 reduced pain at the earliest follow-up compared with no exercise. In a recent systematic
28 review, exercise improved function both at the end of treatment and in the long-term
29 compared with usual care. Exercise also reduced work disability in the long-term. Authors
30 were unable to establish a clear hierarchy between different exercise modalities.
31 Multidisciplinary functional programs consistently improved pain and function in the
32 short- and long-term compared with usual care and physiotherapy and improved the long-
33 term likelihood of returning to work compared to non-multidisciplinary programs.

34
35 Heidari et al. (2023) aimed to systematically analyze the efficacy of aquatic exercise on
36 pain intensity, disability, and quality of life among adults with low back pain. Out of 856
37 articles, 14 RCTs ($n = 484$ participants; 257 in the experimental groups and 227 in the
38 control groups) met inclusion criteria. Pooled results illustrated that aquatic exercises
39 significantly reduced pain, improved disability, and improved quality of life in both the
40 physical component score and the mental component score when compared with a control
41 group. Authors concluded that the current review showed that aquatic exercise regimens

1 were effective among adults with low back pain. High-quality clinical investigations are
 2 still needed to support the use of therapeutic aquatic exercise in a clinical setting.

3
 4 Babiloni-Lopez et al. (2023) aimed to systematically review and synthesize evidence (i.e.,
 5 active [land-based training] and nonactive controls [e.g., receiving usual care]) regarding
 6 the effects of water-based training on patients with nonspecific chronic low-back pain
 7 (NSCLBP). The included studies satisfied the following criteria: (a) NSCLBP (≥ 12 weeks)
 8 patients, (b) water-based intervention, (c) control group (land-based trained; nonactive
 9 group), and (d) outcomes related to pain, disability, quality of life, or flexibility. The main
 10 outcome analyzed in the meta-analysis was pain intensity. Secondary outcomes included
 11 disability, body mass index, and flexibility. After intervention, pain intensity was reduced
 12 compared with nonactive controls and a similar reduction was noted when compared with
 13 land-based trained group. Greater decrease in disability and greater increase in sit-and-
 14 reach were noted after intervention compared with the nonactive group. In conclusion,
 15 water-based exercise therapy reduces pain intensity, disability, and increases flexibility in
 16 NSCLBP compared with nonactive subjects and was equally effective compared with land-
 17 based exercise to reduce pain. Favorable effects may be expected at ≤ 8 weeks. However,
 18 due to several methodological issues (e.g., high heterogeneity), for the improvement of
 19 most outcomes, authors were unable to provide other than a weak recommendation in favor
 20 of intervention compared with control treatment.

21
 22 Ceballos-Laito et al. (2023) evaluated the effectiveness of hip interventions on pain and
 23 disability in patients with LBP in the short-, medium-, and long-term. A total of 2,581
 24 studies were screened. Eight were included in the meta-analysis involving 508 patients
 25 with LBP. The results provided very low certainty that both hip strengthening and hip
 26 stretching improved pain and disability in the short-term, respectively. No benefits were
 27 found in the medium- or long-term. The risk of bias, heterogeneity, and imprecision of the
 28 results downgraded the level of evidence. Very low certainty evidence suggests a positive
 29 effect of hip strengthening in isolation or combined with specific low back exercise and
 30 hip stretching combined with specific low back exercise for decreasing pain intensity and
 31 disability in the short-term, in patients with LBP.

32
 33 Tikhile and Patil (2024) evaluated the efficacy of various physiotherapy strategies in
 34 alleviating LBP, considering a range of interventions and their associated outcomes.
 35 Through a thorough examination of existing literature from January 2017 to October 2023,
 36 this review synthesizes evidence on the effectiveness of interventions such as manual
 37 therapy, exercise therapy, electrotherapy modalities, and education-based approaches. The
 38 review also scrutinizes the comparative effectiveness of different physiotherapy modalities
 39 and their suitability for specific patient populations, considering factors such as chronicity,
 40 severity, and underlying pathology. This review aims to provide insights into the most
 41 effective physiotherapy strategies for alleviating LBP, chronic low back pain (CLBP) and
 42 chronic nonspecific low back pain (CNLBP) and guiding clinical practice toward evidence-

1 based interventions. Twenty-one studies that fulfilled the criteria for inclusion (aged 20 to
2 50 years and of both genders) were added to the review. Exercises for core stability,
3 strengthening, orthosis, transcutaneous electrical nerve stimulation, heat massage therapy,
4 interferential current, Mulligan's mobilization, low-level laser therapy, and McGill
5 stabilization exercises (core exercises) were among the therapeutic strategies. The
6 McKenzie method (back exercises), ultrasound, sensory-motor training, Swiss ball
7 exercises, and other techniques reduced pain and enhanced strength, balance, and ease of
8 daily activities. Every therapeutic approach has an impact on recovery rates ranging from
9 minimal to maximal. Conventional physical therapy is less effective than most recent
10 advanced techniques like mobilization and exercises. In summary, the integration of
11 manual techniques, orthoses and alternative intervention strategies with conservative
12 therapeutic approaches can effectively alleviate pain, enhance function and yield better
13 overall outcomes. To get more information about the optimal dosage, therapeutic
14 modalities and long-term effects of these treatments, more research is required.

15
16 El Melhat et al. (2024) explored the effectiveness and patient-related outcomes of various
17 conservative approaches, including physical therapy modalities and alternative therapies
18 in the treatment of lumbar disc herniation associated with radiculopathy (LDHR). The
19 objective of this article was to introduce advanced and new treatment techniques,
20 supplementing existing knowledge on various conservative treatments. Authors identified
21 the following interventions to yield moderate evidence (Level B) of effectiveness for the
22 conservative treatment of LDHR: patient education and self-management, McKenzie
23 method, mobilization and manipulation, exercise therapy, traction (short-term outcomes),
24 neural mobilization, and epidural injections. Two interventions were identified to have
25 weak evidence of effectiveness (Level C): traction for long-term outcomes and dry
26 needling. Three interventions were identified to have conflicting or no evidence (Level D)
27 of effectiveness: electro-diagnostic-based management, laser and ultrasound, and
28 electrotherapy.

29
30 Maharty et al. (2024) summarized evaluation and management of chronic LBP in adults.
31 Patients with chronic low back pain should have a history and physical examination to
32 identify red flags that may indicate serious conditions that warrant immediate intervention
33 or yellow flags (i.e., psychological, environmental, and social factors) that indicate risk of
34 disability. The examination should include an evaluation for radicular symptoms. Routine
35 imaging is not recommended but is indicated when red flags are present, there is a
36 neuromuscular deficit, or if pain does not resolve with conservative therapy. Patients
37 should avoid bed rest. Nonpharmacologic treatment is first-line management and may
38 include therapies with varying evidence of support, such as counseling, exercise therapy,
39 spinal manipulation, massage, heat, dry needling, acupuncture, transcutaneous electrical
40 nerve stimulation, and physical therapy. Pharmacologic interventions are second-line
41 treatment. Nonsteroidal anti-inflammatory drugs are the initial medication of choice;
42 duloxetine may also be beneficial. Evidence is inconclusive to recommend the use of

1 benzodiazepines, muscle relaxants, antidepressants, corticosteroids, insomnia agents,
2 anticonvulsants, cannabis, acetaminophen, or long-term opioids. Epidural corticosteroid
3 injections are not recommended except for short-term symptom relief in patients with
4 radicular pain. Most patients with chronic low back pain will not require surgery;
5 evaluation for surgery may be considered in those with persistent functional disabilities
6 and pain from progressive spinal stenosis, worsening spondylolisthesis, or herniated disk.

7
8 Zhou et al. (2024) aimed to identify and compare the recommendations of recent clinical
9 practice guidelines (CPGs) for the management of LBP across the world. Analysis
10 identified a total of 22 CPGs that met the inclusion criteria and were of middle and high
11 methodological quality. The guidelines exhibited heterogeneity in their recommendations,
12 particularly in the approach to different stages of LBP. For acute LBP, the guidelines
13 recommended the use of non-steroidal anti-inflammatory drugs (NSAIDs), therapeutic
14 exercise, staying active, and spinal manipulation. For subacute LBP, the guidelines
15 recommended the use of NSAIDs, therapeutic exercise, staying active, and spinal
16 manipulation. For chronic LBP, the guidelines recommended therapeutic exercise, the use
17 of NSAIDs, spinal manipulation, and acupuncture. Authors state that current CPGs provide
18 recommendations for almost all major aspects of the management of LBP, but there is
19 marked heterogeneity between them. Some recommendations lack clarity and overlap with
20 other treatments within the guidelines.

21
22 Patti et al. (2024) investigated the effectiveness of Pilates exercise on pain intensity and
23 functional disability caused by LBP. Randomized controlled trials (RCTs) evaluating LBP
24 in which the primary treatment was based on Pilates exercise compared with no exercise,
25 or non-specific exercise were included in the literature search. The search returned 1566
26 records of which 36 articles were included in the systematic review and 19 in the meta-
27 analysis. Twenty-two studies compared the effects of Pilates exercise vs no exercise and
28 13 studies examined the effects of Pilates exercise vs non-specific exercise. Analysis
29 showed that Pilates had a positive effect on the perception of LBP vs no exercise. A similar
30 trend occurred with non-specific exercise. Authors concluded that Pilates exercise can
31 decrease LBP compared to no exercise and non-specific exercise. General practitioners
32 should consider Pilates exercise as an effective strategy to manage LBP and counteract the
33 growing health.

34
35 Liang et al. (2024) quantified the dose-response relationship between overall and specific
36 exercise modalities and pain, in patients with nonspecific chronic low back pain. Authors
37 included randomized controlled trials of exercise interventions in adults with nonspecific
38 chronic LBP and at least 1 pain outcome reported at the main trial end point. Eighty-two
39 trials were included (n = 5033 participants). They found a nonlinear dose-response
40 relationship between total exercise and pain in patients with nonspecific chronic LBP. The
41 maximum significant response was observed at 920 MET minutes. The minimal clinically
42 important difference for achieving meaningful pain improvement was 520 MET minutes

1 per week. The dose to achieve minimal clinically important difference varied by type of
2 exercise; Pilates was the most effective. The certainty of the evidence was very low to
3 moderate for all outcomes. Authors concluded that the dose-response relationship of
4 different exercise modalities to improve pain in patients with nonspecific chronic LBP had
5 a U-shaped trajectory and low- to moderate-certainty evidence. The clinical effect was
6 most pronounced with Pilates exercise.

7
8 IJzelenberg et al. (2024) assessed the effectiveness of exercise for acute non-specific low
9 back pain (LBP) vs: (1) sham treatment and (2) no treatment at short term (main follow-up
10 time). The primary outcomes were back pain, back-specific functional status, and recovery.
11 Authors identified 23 randomized controlled trials (2674 participants). There is very low-
12 certainty evidence that exercise therapy compared with sham/placebo treatment has no
13 clinically relevant effect on pain and on functional status in the short term. There is very
14 low-quality evidence which suggests no difference in effect on pain and functional status
15 for exercise vs no treatment at short-term follow-up. Similar results were found for the
16 other follow-up moments. The certainty of the evidence was downgraded because many
17 randomized controlled trials had a high risk of bias, were small in size, and/or there was
18 substantial heterogeneity. Authors concluded that exercise therapy compared with
19 sham/placebo and no treatment may have no clinically relevant effect on pain or functional
20 status in the short term in people with acute non-specific LBP, but the evidence is very
21 uncertain. Owing to insufficient reporting of adverse events, authors were unable to reach
22 any conclusions on the safety or harms related to exercise therapy.

23
24 de Roode et al. (2024) investigated the proportion of improvement in pain and disability
25 that can be attributed to contextual effects in the outcome of exercise therapy for patients
26 with low back pain in a systematic review. Low back pain is the leading cause of global
27 disability for which exercise therapy is a widely recommended treatment. Research
28 indicates that contextual factors may also influence treatment outcomes in low back pain.
29 Examples include the patient-therapist relationship and other treatment-related
30 circumstances that affect patient expectations. By focusing on the specific treatment effect,
31 clinical trials often ignore the effect of contextual factors, thereby contributing to the so-
32 called efficacy paradox. This means that treatment effects observed in clinical practice are
33 often greater than those reported in clinical trials. Authors conducted a meta-analysis with
34 eligible articles reporting randomized controlled trials that compared exercise therapy to
35 placebo interventions. Outcomes of interest were pain and disability. Meta-analysis was
36 carried out to calculate the proportion attributable to contextual effects for both pain and
37 disability. Eight studies met the inclusion criteria and were included in the meta-analysis.
38 Five studies were rated as having a moderate risk of bias and two studies had a low risk of
39 bias. Proportion attributable to contextual effects was 0.60 for pain and 0.69 for disability.
40 Authors concluded that a large extent of pain and disability improvement after exercise
41 therapy in low back pain is attributable to contextual effects although this conclusion is
42 based on low certainty evidence.

1 Diez-Buil et al. (2024) compared whether the combination of exercise with education is
2 more effective for the treatment of low back and/or pelvic pain (PP) than each of these
3 interventions separately in pregnant women. A total of 13 articles were selected. There is
4 a significant decrease in pain in the combination of exercise and education compared with
5 education alone. With respect to disability, there is a significant decrease in the exercise
6 and education group compared with the group that only addressed education). One article
7 analyzed kinesophobia, reporting no significant changes. Authors concluded that the
8 combination of exercise and education seems to be more effective in reducing pain and
9 disability in pregnant women with low back and/or PP than the use of education alone. In
10 kinesophobia, the results found are not significant.

11
12 Herrero et al. (2024) analyzed whether interventions aimed at modifying lifestyle can be
13 effective in improving pain intensity and functional disability in CNSLBP. A total of 20
14 studies were included for qualitative analysis, of which 16 were randomized clinical trials
15 with a moderate-high methodological quality and were part of the quantitative analysis.
16 The interventions that had the greatest effect in reducing pain intensity were cognitive
17 therapy combined with functional exercise programs, lumbar stabilization exercise and
18 resistance exercise; meanwhile, for functional disability, they were functional exercise
19 programs, aerobic exercise and standard care. In conclusion, a multimodal intervention
20 aimed at changing one's lifestyle that encompasses cognitive, behavioral, and physical
21 aspects seems to be highly effective in improving pain intensity and functional disability
22 caused by CNSLBP; however, it is not yet known if these improvements are maintained in
23 the long term.

24
25 Salehi et al. (2024) assessed the efficacy of specific exercises in general population with
26 non-specific low back pain (LBP). Fifty-four trials met the inclusion criteria for this study.
27 Additionally, 46 of these trials were randomized controlled trials and were further
28 evaluated for the meta-analysis. Authors included trials investigating the effectiveness of
29 exercise therapy, including isometric activation of deep trunk muscles, strengthening
30 exercises, stabilization exercises, stretching exercises, and proprioceptive neuromuscular
31 facilitation exercises (PNF) in LBP patients. The primary outcome was pain intensity,
32 measured using tools such as the visual analogue scale (VAS) and numeric pain rating scale
33 (NPRS). The secondary outcome was disability, assessed through instruments such as the
34 Roland Morris Disability Questionnaire (RMDQ) and Oswestry Disability Index (ODI).
35 The meta-analysis indicated a small efficacy in favor of isometric activation of deep trunk
36 muscles, a moderate efficacy in favor of stabilization exercises, and a large efficacy in
37 favor of PNF exercises for reducing pain intensity as assessed by VAS or NPRS tools.
38 Moreover, the meta-analysis revealed a moderate efficacy for isometric activation of deep
39 trunk muscles, and a large efficacy for PNF exercises in improving disability, assessed
40 using RMDQ or ODI questionnaires. The level of certainty in the evidence was very low
41 to low. Authors concluded that these findings emphasize the importance of incorporating
42 localized therapeutic exercises as a fundamental aspect of managing non-specific LBP.

1 Clinicians should consider utilizing localized therapeutic exercise tailored to individual
2 patient needs. Furthermore, further research investigating optimal exercise therapy,
3 optimal dose of the exercises, durations, and long-term adherence is warranted to enhance
4 the precision and efficacy of exercise-based interventions for non-specific LBP.

5
6 Gou et al. (2024) synthesized the available data on the effectiveness of hamstring stretching
7 exercises in relieving pain intensity and improving function for patients with low back pain.
8 Searches retrieved 344 trials, of which 14 met the inclusion criteria for this review (n = 735
9 participants). The combined meta-analysis showed hamstring stretching resulted in lower
10 pain scores in different categories of low back pain. Subgroup analysis showed that
11 hamstring stretching led to a larger range of motion for cases of back pain with radiating
12 pain. The combined meta-analysis revealed that hamstring stretching resulted in lower
13 Oswestry Disability Index scores in comparison to regular treatment, particularly in
14 individuals suffering from low back pain across all subtypes. Authors concluded that this
15 meta-analysis demonstrated the effectiveness of hamstring stretching exercises in reducing
16 pain intensity in various categories of low back pain and improving the straight leg raise in
17 patients experiencing back pain with radiating pain. Additionally, it highlights the
18 improvement in function for patients with back pain across all subtypes.

19
20 Lim et al. (2024) assessed the efficacy of pelvic floor muscle training and physical therapy
21 interventions in patients with low back pain. Studies providing pelvic floor muscle training
22 in individuals with low back pain were included. Nineteen studies were included in this
23 review. Pelvic floor muscle training showed low certainty evidence for improving pain and
24 reflected a clinically meaningful reduction in pain. The evidence for disability
25 improvement had a low certainty due to high heterogeneity. Substantial improvements in
26 pain and disability were observed when pelvic floor muscle training was added to standard
27 physical therapy, with low certainty of evidence supporting these findings. Whereas pelvic
28 floor muscle training substantially improved pain compared to other interventions, there
29 was no marked improvement in disability. Authors concluded that pelvic floor muscle
30 training is potentially beneficial in addition to physical therapy for reducing low back pain,
31 particularly in pregnancy-related cases. However, the evidence should be interpreted
32 considering the quality and risk of bias.

33
34 Alonso-Sal et al. (2024) evaluated the effectiveness of exercise interventions for managing
35 nonspecific low back pain (NSLBP) and explores their impact on related biopsychosocial
36 factors, physical health variables, and inflammatory biomarkers. Materials and Methods:
37 Fifteen randomized controlled trials involving 1338 participants aged 18 to 65 years with
38 NSLBP were included. Exercise significantly reduced pain intensity and improved
39 biopsychosocial factors such as depression, disability, functionality, quality of life, and
40 kinesiophobia. Additionally, it enhanced physical parameters like proprioception, muscle
41 thickness, and physical performance. However, the review found insufficient evidence
42 regarding the effects of exercise on inflammatory biomarkers in NSLBP patients. Authors

1 concluded that findings suggest that physical exercise is an effective intervention for pain
2 reduction and the improvement of overall health in NSLBP, though further research is
3 needed to clarify its impact on inflammation.

4
5 Lim et al. (2025) summarized the content and critically appraised the quality and
6 applicability of recent clinical practice guidelines (CPGs) for nonpharmacological,
7 nonsurgical management of spine pain. Authors included 30 CPGs, primarily (90%)
8 developed in Western countries, which contained 404 recommendations. High-quality
9 CPGs consistently recommended exercise therapy and multimodal care, encompassing a
10 combination of exercises, mobilization/manipulation, education, alternative medicine, and
11 cognitive-behavioral treatments. Generally, CPGs did not recommend assistive (eg, corsets
12 and orthosis) devices or electro/thermotherapies (eg, therapeutic ultrasound and
13 transcutaneous electrical nerve stimulation). Approximately half of the CPGs demonstrated
14 good methodological quality according to AGREE II, whereas the rest were of poor quality.
15 On the AGREE-REX assessment, one third of the recommendations were of excellent
16 quality. Although recent guidelines frequently recommended exercise therapy and
17 multimodal care for the management of spine pain, their recommendations often
18 overlooked demographics and comorbidities. Despite methodological improvements, most
19 CPGs lacked simple clinical applicability and considerations of knowledge users' values.

20
21 Comachio et al. (2025) synthesized the evidence from systematic reviews on the benefits
22 and harms of exercise therapy and physical activity (PA) for the secondary prevention and
23 management of low back pain (LBP). Eligible studies were systematic reviews of
24 randomized controlled trials and observational studies, with or without meta-analyses. The
25 primary outcome for secondary prevention was LBP recurrence, while for management,
26 primary outcomes included pain intensity and disability, with adverse events as secondary
27 outcomes. Data were extracted across immediate, short-term, intermediate, and long-term
28 follow-up periods. A total of 70 systematic reviews were included, 43 with meta-analyses,
29 7 with network meta-analyses, and 20 without meta-analyses. Six (out of 10) reviews with
30 meta-analyses for secondary prevention indicated a small benefit from general exercises
31 and leisure-time PA (low-to-moderate certainty). For LBP management, 35 (out of 36)
32 reviews reported that exercise therapies such as Pilates, motor control, mixed exercise, Tai
33 Chi, water-based exercises, and yoga showed small beneficial effects on pain and disability
34 compared to minimal intervention mainly in the short-term (low-to-moderate certainty).
35 Seven network meta-analyses favored motor control and Pilates over other forms of
36 exercise to reduce pain (low certainty). Adverse events were reported in less than 31% of
37 the reviews, predominantly involving post-exercise soreness and temporary increases in
38 pain, mainly in yoga-related studies. Adverse events were considered minor, and no serious
39 adverse events were reported. There is low-to-moderate certainty that exercise therapy and
40 leisure-time PA are beneficial for improving pain and preventing the recurrence of LBP.
41 However, evidence on the potential harms of these interventions is limited, and adverse
42 events related to exercise and PA remain under-investigated.

1 González-Gómez et al. (2025) compared the efficacy of ET with MT in terms of pain
2 intensity, disability and physical function in people with CLBP in a systematic review with
3 meta-analysis and meta-regression of randomized controlled trials (RCTs). Six RCTs (743
4 patients) were included. Meta-analyses showed, albeit non-clinically relevant, a significant
5 difference for long-term in favor of ET for disability. Meta-regression showed that the
6 female-male ratio, treatment duration and mean age explain variability in pain intensity
7 and disability. Authors conclude that ET had a small beneficial effect on long-term
8 disability in people with CLBP. Nevertheless, evidence does not provide conclusive
9 differences between both the treatments overall, influenced by heterogeneity and the
10 number of studies included. Biopsychosocial factors may moderate the differences in
11 outcomes. The GRADE assessment revealed very low certainty across all outcomes,
12 highlighting the lack of high-quality research.

13
14 Cheng et al. (2025) validated the effectiveness of six exercise therapies in treating low back
15 pain using Meta-analysis methods, and proposed optimal exercise duration, frequency, and
16 cycle. The intervention measures for the experimental group included core stability, Tai Ji,
17 aquatic therapy, yoga, sling exercise, and combined training. The intervention programs
18 included in the various studies exhibited significant differences, with each exercise session
19 ranging from 15 to 75 min in duration, the frequency of exercise per week varying from
20 once to seven times, and the shortest exercise period lasted for 2 weeks, while the longest
21 lasted for a year. The outcome indicators used the VAS/NRS to assess pain intensity and
22 the ODI and RMDQ to evaluate lumbar dysfunction. The overall effect size for six exercise
23 therapies for low back pain was significant. Subgroup analyses showed yoga had the largest
24 effect. Exercise duration ≤ 30 min, frequencies > 4 times/week, and cycles ≤ 4 weeks were
25 most effective. Sample sizes of 30~60 cases and studies with moderate bias risk also
26 showed large effects. The Oswestry Disability Index scores demonstrated the most
27 significant effect size. The effect size of the physical factors in the control group was the
28 largest. Authors concluded that all six exercise therapies effectively alleviated low back
29 pain, with yoga showing the best results. The optimal exercise intervention protocol
30 involved exercise duration not exceeding 30 min per session, frequency of more than 4
31 times per week, and cycle not exceeding 4 weeks. Additionally, exercise interventions
32 exhibited the most significant improvements in Oswestry Disability Index scores for low
33 back pain.

34
35 Rizzo et al. (2025) provided accessible, high-quality evidence on the effects of non-
36 pharmacological and non-surgical interventions for people with LBP and to highlight areas
37 of remaining uncertainty and gaps in the evidence regarding the effects of these
38 interventions for people with LBP. Authors included 31 Cochrane reviews of 644 trials that
39 randomized 97,183 adults with LBP. They have high confidence in the findings of 19
40 reviews, moderate confidence in the findings of two reviews, and low confidence in the
41 findings of 10 reviews. Results are presented for non-pharmacological/non-surgical
42 interventions compared to placebo/sham or no treatment/usual care at short-term (\leq three

1 months) follow-up. Placebo/sham comparisons: Acute/subacute LBP compared to placebo,
2 there is probably no difference in function (at one-week follow-up) for spinal manipulation
3 (moderate-certainty evidence). Data for safety were reported only for heated back wrap.
4 Compared to placebo, heated back wrap may result in skin pinkness (low-certainty
5 evidence). Chronic LBP compared to sham acupuncture, acupuncture probably provides a
6 small improvement in function (moderate-certainty evidence). Compared to sham traction,
7 there is probably no difference in pain intensity for traction (moderate-certainty evidence).
8 Data for safety were reported only for acupuncture. There may be no difference between
9 acupuncture and sham acupuncture for safety outcomes (low-certainty evidence). No
10 treatment/usual care comparisons: Acute/subacute LBP compared to advice to rest, advice
11 to stay active probably provides a small reduction in pain intensity (moderate-certainty
12 evidence). Compared to advice to rest, advice to stay active probably provides a small
13 improvement in function (moderate-certainty evidence). Data for safety were reported only
14 for massage. There may be no difference between massage and usual care for safety (low-
15 certainty evidence). Chronic LBP compared to no treatment, acupuncture probably
16 provides a medium reduction in pain intensity (moderate-certainty evidence), and a small
17 improvement in function (moderate-certainty evidence). Compared to usual care,
18 acupuncture probably provides a small improvement in function (moderate-certainty
19 evidence). Compared to no treatment/usual care, exercise therapies probably provide a
20 small to medium reduction in pain intensity (moderate-certainty evidence), and probably
21 provide a small improvement in function (moderate-certainty evidence). Compared to
22 usual care, multidisciplinary therapies probably provide a medium reduction in pain
23 intensity (moderate-certainty evidence), and probably provide a small improvement in
24 function (moderate-certainty evidence). Compared to no treatment, psychological therapies
25 using operant approaches probably provide a small reduction in pain intensity (moderate-
26 certainty evidence). Compared to usual care, psychological therapies (including
27 progressive muscle relaxation and behavioral approaches) probably provide a small
28 reduction in pain intensity (moderate-certainty evidence), but there is probably no
29 difference in function (moderate-certainty evidence). It is uncertain whether there is a
30 difference between non-pharmacological/non-surgical interventions and no
31 treatment/usual care for safety (very low-certainty evidence). Authors' concluded that
32 spinal manipulation probably makes no difference to function compared to placebo for
33 people with acute/subacute LBP. Acupuncture probably improves function slightly for
34 people with chronic LBP, compared to sham acupuncture. There is probably no difference
35 between traction and sham traction for pain intensity in people with chronic LBP.
36 Compared to advice to rest, advice to stay active probably reduces pain intensity slightly
37 and improves function slightly for people with acute LBP. Acupuncture probably reduces
38 pain intensity, and improves function slightly for people with chronic LBP, compared to
39 no treatment. Acupuncture probably improves function slightly for people with chronic
40 LBP, compared to usual care. Exercise therapies probably reduce pain intensity, and
41 improve function slightly for people with chronic LBP, compared to no treatment/usual
42 care. Multidisciplinary therapies probably reduce pain intensity, and improve function

1 slightly for people with chronic LBP, compared to usual care. Compared to usual care,
2 psychological therapies probably reduce pain intensity slightly, but probably make no
3 difference to function for people with chronic LBP.

4
5 Jin et al. (2025) evaluated the effects of postoperative exercise on pain, disability, and
6 quality of life after lumbar disc herniation surgery. Compared with the control group, the
7 exercise intervention group had significantly lower visual analog scale scores both in the
8 short-term and long-term follow-up. The exercise intervention group also showed a
9 significant reduction in Oswestry Disability Index in the short-term follow-up, but not in
10 the long-term follow-up. Among the Short-Form Health Survey factors, physical function,
11 pain, general health, energy and vitality, and social function significantly improved in the
12 exercise intervention group compared with the control group. Authors concluded that
13 exercise after lumbar disc herniation surgery was effective in reducing pain and disability
14 and improving quality of life.

15
16 Jenkins et al. (2025) assessed the long-term effectiveness of non-surgical interventions for
17 reducing pain intensity and disability in adults with chronic low back pain. 75 trials (15
18 395 participants) were included. Risk of bias was rated high for the majority of studies (51
19 [68%] of 75). In people with non-specific chronic low back pain at long-term follow-up,
20 there was moderate certainty evidence that cognitive behavioral therapy and mindfulness
21 probably result in reductions in pain intensity for cognitive behavioral therapy and for
22 mindfulness) and disability. Goal setting and needling probably reduce disability at long-
23 term follow-up. There was low certainty evidence that multidisciplinary care could reduce
24 pain intensity and exercise might reduce disability at very long-term follow-up.
25 Heterogeneity was evident in several of the meta-analyses, and results should be interpreted
26 with caution. Some interventions, including cognitive behavioral therapy, mindfulness,
27 exercise, and multidisciplinary care could produce the long-term benefits required to
28 reduce the global burden due to non-specific chronic low back pain; however, the effects
29 are mostly small, and the strength of evidence is relatively uncertain. Greater attention is
30 needed on developing and testing interventions with long-term effects for chronic low back
31 pain.

32
33 Hennemann et al. (2025) sought to determine the effectiveness of the McKenzie Method
34 compared to any conservative interventions on pain and disability in patients with chronic
35 low back pain (LBP) with directional preference (DP). Five trials (n = 743) were included.
36 There was low-certainty evidence that the McKenzie Method, compared to all other
37 interventions combined, produced clinically important reductions in short-term pain and in
38 intermediate-term disability. Low-to-moderate certainty evidence showed that the
39 McKenzie Method also resulted in clinically important improvements in short-term pain
40 and disability when compared specifically to other exercise approaches, and in
41 intermediate-term pain and disability as well as long-term disability when compared to
42 minimal intervention. Low-certainty evidence showed usually small, clinically

1 unimportant effects in comparison to manual therapy. Authors found low-to-moderate
2 certainty evidence that the McKenzie Method was superior to all other interventions
3 combined for up to 6 months for pain and up to 12 months for disability, with clinically
4 important differences versus exercise in the short term and versus minimal interventions in
5 the intermediate term. The only clinically important long-term effect was on disability
6 compared to minimal intervention.

7
8 Zhao et al. (2025) investigated the impact of various combinations of exercise prescription
9 variables-namely type, duration, frequency, and period-on improving chronic low back
10 pain (CLBP) in adults. The goal is to provide evidence to inform the development of
11 exercise prescriptions for CLBP interventions. Among the 26 studies included, the exercise
12 interventions in the experimental groups primarily involved core stability training,
13 combined training, Pilates, yoga, qigong, suspension training, Swiss ball exercises,
14 strength training, perturbation therapy, sit-up exercises, and tai chi. The intervention
15 durations varied, ranging from 4, 6, 8, 12, 13, and ≥ 16 weeks. The frequency of
16 interventions ranged from 1–2 sessions, 3 sessions, 5 sessions, to 7 sessions per week.
17 Intervention durations ranged from 15–30 min, 40 min, 45 min, 50 min, to ≥ 60 min.
18 Significant effects were observed for durations of 15-30 min and ≥ 60 min when compared
19 to the control group. Intervention periods of 4 weeks, 12 weeks and ≥ 16 weeks also resulted
20 in significantly better outcomes compared to the control group. The intervention effect for
21 durations of ≥ 16 weeks was significantly greater than that for 12 weeks and 6 weeks. A
22 frequency of three sessions per week demonstrated significantly superior outcomes
23 compared to the control group. An intervention duration of 15 to 30 min, three sessions per
24 week, an intervention period of ≥ 16 weeks, and Tai Chi exercise may be the most effective
25 approaches for improving chronic low back pain in adults. Authors concluded that Tai Chi
26 exercise, lasting 15 to 30 min per session, performed three times a week over an
27 intervention period of at least 16 weeks, may represent the most effective intervention for
28 alleviating chronic low back pain in adults. However, due to the limited number of studies
29 included, further research is necessary to provide stronger evidence.

30
31 Jones et al. (2025) sought to determine the association between exercise adherence and the
32 effects of exercise on pain intensity and functional limitations in adults with chronic non-
33 specific low back pain (CNSLBP) in a systematic review with meta-analysis. Interventions
34 included in RCTS reviewed were trials of exercise compared with no exercise (e.g., usual
35 care, placebo/sham or another conservative treatment). Adherence to exercise must have
36 been reported. This study included 46 trials with 56 exercise groups. High exercise
37 adherence (80 to 100%) was associated with reduced pain intensity (0 to 100 scale) (low
38 certainty) and functional limitations (0 to 100 scale) (low certainty). Moderate exercise
39 adherence (60 to 79%) was not associated with reduced pain intensity (very low certainty)
40 or functional limitations (very low certainty). Low exercise adherence ($< 59\%$) was
41 associated with reduced pain intensity (low certainty) and functional limitations (moderate
42 certainty). Compared with low adherence, additional differences in outcomes for moderate

1 and high adherence were mostly negligible. Higher exercise adherence is associated with
2 larger improvements in clinical outcomes in adults with CNSLBP, although overall
3 differences are small compared with lower adherence. Other factors besides adherence
4 between the trials and exercise programs could explain these results. Further research is
5 needed to determine the causal effect of exercise adherence on outcomes in adults with
6 CNSLBP.

7
8 García-Moreno et al. (2025) compared the different physiotherapy treatments and
9 determine the most effective treatment to reduce the nonspecific low back pain (NSLBP)
10 intensity in children and adolescents. A meta-analysis of 11 controlled trials with 827
11 participants found that physiotherapy treatments effectively reduced NSLBP intensity on
12 posttest measurement and 6-month follow-up. Network meta-analysis showed both
13 therapeutic exercise and a combination of therapeutic exercise and manual therapy were
14 effective compared to no treatment. There were no significant differences between
15 therapeutic exercise and the combination of therapeutic exercise and manual therapy.
16 Authors concluded that physical exercise has proven to be the most effective treatment for
17 addressing the intensity of NSLBP in children and adolescents. While combining it with
18 manual therapy may yield even better results, it is crucial to emphasize that physical
19 exercise should serve as the cornerstone in the physiotherapeutic approach to managing
20 NSLBP intensity in this age group.

21
22 Niederer et al. (2025) estimated the comparative effectiveness of tailored exercise therapies
23 in people with chronic, nonspecific low back pain in a systematic review with network
24 meta-analysis. Only randomized controlled trials (RCTs) studying the effects of any type
25 of tailored exercise therapies in persons with chronic, nonspecific low back pain were
26 included. Outcomes of interest were pain intensity and disability. Authors included 58
27 randomized trials (n = 10,510 participants) with 29 different treatment or control
28 categories. Risk of bias for pain intensity and disability was high. Cognitive functional
29 therapy alone or combined with biofeedback were, with moderate-certainty evidence, the
30 most effective treatments for pain intensity and disability reduction when compared to
31 usual care. Authors concluded that in comparison to usual care, cognitive functional
32 therapy, alone or with biofeedback, likely results in a large disability reduction.

33
34 Vega-Retuerta et al. (2025) determined the effectiveness of multidisciplinary interventions
35 including exercise in improving health outcomes for patients with non-specific chronic low
36 back pain (NSCLBP) across different regions. Multidisciplinary interventions including
37 exercise aimed at relieving symptoms of NSCLBP were included. Outcomes analyzed were
38 disability, pain, fear avoidance, quality of life, perceived change, depression, anxiety and
39 stress. Thirty-one studies were included in the qualitative analysis and 29 in the meta-
40 analysis. The qualitative analysis showed improvements in favor of the intervention group
41 (IG) in pain, fear avoidance, quality of life and perceived change. The meta-analysis
42 reported significant pooled effect size (ES) in favor of the IGs in disability, pain, fear

1 avoidance, quality of life, and perceived change among patients with NSCLBP.
 2 Multidisciplinary approaches including exercise appear to be effective in reducing
 3 disability, pain, fear avoidance, quality of life and perceived change, but not depression,
 4 anxiety or stress in patients with NSCLBP. The findings should be interpreted cautiously
 5 in light of substantial heterogeneity and the limited quality of the available evidence.

6
 7 Baroncini et al. (2025) compared active versus passive physiotherapy versus their
 8 combination in terms of pain and disability in patients with mechanical and/or aspecific
 9 cLBP in a Bayesian network meta-analysis. Data from 2768 patients (mean age 46.9 ± 10.9
 10 years, mean BMI 25.8 ± 2.9 kg/m²) were collected. The mean length of follow-up was 6.2
 11 ± 6.1 months. Between groups, comparability was found at baseline in terms of mean age,
 12 proportion of women, mean BMI, symptom duration and patient-reported outcome
 13 measures (PROMs). By the end of the follow-up period, the active group evidenced the
 14 lowest pain scores. The active group evidenced the lowest RMQ score. The active group
 15 evidenced the lowest ODI score. Authors concluded that active physiotherapy showed
 16 better results than passive physiotherapy and a combination of both for the management of
 17 mechanical and/or non-specific cLBP.

18
 19 Xia et al. (2025) evaluated four mind-body exercises (MBEs), including Tai Chi (TC),
 20 Qigong (QG), Daoyin, and Yoga to determine their efficacy in managing CNSLBP. The
 21 results of the network meta-analysis showed that in terms of pain scores (PS), Qigong had
 22 the best effect; in terms of physical functioning (PF), Qigong remained the best
 23 intervention; whereas, in terms of quality of life (QOL), Tai Chi had the most significant
 24 effect. Qigong was superior in terms of pain and functional improvement, while Tai Chi
 25 performed optimally in terms of quality-of-life improvement. QG is the most effective for
 26 pain relief and improving physical function, while TC best enhances QoL. Different MBEs
 27 have distinct effects on functional outcomes, suggesting that interventions should be
 28 tailored to individual patient needs. Combining multiple approaches may optimize results
 29 for CNSLBP management. This study highlights the efficacy of mind-body exercises
 30 (Qigong, Yoga, Tai Chi) in improving pain, function, and quality of life for chronic low
 31 back pain. Qigong and Yoga excel in pain relief, while Tai Chi enhances well-being. By
 32 integrating physical movement with mental focus, these low-intensity, adaptable therapies
 33 offer a holistic, non-pharmacological approach, emphasizing personalized care for long-
 34 term management.

35
 36 Grande et al. (2025) investigated the barriers and facilitators to physical activity and
 37 exercise among people with chronic low back pain (CLBP). Qualitative or mixed-methods
 38 studies with a qualitative approach were included. Studies must have recruited adults, of
 39 both sexes, aged 18 years or older with CLBP. Fifty-seven studies were included. Barriers
 40 to physical activity were mainly related to pain intensity, fear of movement, intervention
 41 type, lack of information, motivation, and support. Facilitators of physical activity were
 42 adequate information, professional and social support, perceived benefits, and favorable

1 conditions to engage in physical activity. Authors concluded that the barriers to people
2 with chronic LBP engaging in physical activity included pain intensity and fear of reinjury,
3 type of intervention, lack of information, motivation and support, and occupational and
4 socioenvironmental factors. The main facilitators were receiving information and support
5 from health professionals, motivational activities, knowledge about benefits of the
6 intervention, and external factors.

7
8 Joyce et al. (2025) sought to determine whether, in active individuals with sacroiliac joint
9 dysfunction (SIJD), an exercise intervention, manual manipulation, or a combination of the
10 two effective in reducing pain and improving function. Three studies were included in this
11 critically appraised topic. One study found that both exercise therapy and manipulation
12 therapy decreased pain and improved function, but found no between-group differences.
13 The other 2 studies found that exercise therapy, manipulation therapy, and the combination
14 of the 2 were effective in decreasing pain and improving function in the participants. The
15 results of the studies suggested that exercise therapy, manual therapy, and a combination
16 of the 2 therapies can be effective in reducing pain and dysfunction in patients with SIJD.
17 While manual therapy techniques alone may be effective in reducing SIJD pain short term,
18 exercise therapy or a combination of the 2 may be more effective in reducing pain long
19 term.

20
21 Jangra et al. (2025) performed a systematic review and meta-analysis on efficacy of
22 physical therapy interventions in sacroiliac joint dysfunction. Randomized controlled trials
23 on sacroiliac joint dysfunction, using physiotherapy as an intervention were evaluated for
24 changes in pain and disability. The result of the study showed a statistically significant
25 improvement of muscle energy technique on pain and mobilization on pain, exercise on
26 pain, and exercise on disability. However, there was no significant result of taping on pain
27 and muscle energy technique, mobilization and taping on disability. It can be concluded
28 that physiotherapy interventions are effective in managing the pain and restoring the
29 functional disability of the patients suffering from sacroiliac joint dysfunction. MET and
30 exercises were found to be more effective in reducing pain than mobilization in treating
31 sacroiliac joint dysfunction and only exercises are effective in improving the disability in
32 SIJD. Therefore a combined approach may be more beneficial.

33 34 **PRACTITIONER SCOPE AND TRAINING**

35 Practitioners should practice only in the areas in which they are competent based on their
36 education training and experience. Levels of education, experience, and proficiency may
37 vary among individual practitioners. It is ethically and legally incumbent on a practitioner
38 to determine where they have the knowledge and skills necessary to perform such services.

39
40 It is best practice for the practitioner to appropriately render services to a patient only if
41 they are trained, equally skilled, and adequately competent to deliver a service compared
42 to others trained to perform the same procedure. If the service would be most competently

1 delivered by another health care practitioner who has more skill and expert training, it
2 would be best practice to refer the patient to the more expert practitioner.

3
4 Best practice can be defined as a clinical, scientific, or professional technique, method, or
5 process that is typically evidence-based and consensus driven and is recognized by a
6 majority of professionals in a particular field as more effective at delivering a particular
7 outcome than any other practice (Joint Commission International Accreditation Standards
8 for Hospitals, 2020).

9
10 Depending on the practitioner’s scope of practice, training, and experience, a member’s
11 condition and/or symptoms during examination or the course of treatment may indicate the
12 need for referral to another practitioner or even emergency care. In such cases it is prudent
13 for the practitioner to refer the member for appropriate co-management (e.g., to their
14 primary care physician) or if immediate emergency care is warranted, to contact 911 as
15 appropriate. See the *Managing Medical Emergencies (CPG 159 – S)* clinical practice
16 guideline for information.

17 **References**

18
19 Abenhaim, L., Rossignol, M., Valat, J. P., Nordin, M., Avouac, B., Blotman, F., Charlot,
20 J., Dreiser, R. L., Legrand, E., Rozenberg, S., & Vautravers, P. (2000). The role of
21 activity in the therapeutic management of back pain. Report of the International Paris
22 Task Force on Back Pain. *Spine*, 25(4 Suppl), 1S–33S

23
24 Almeida MO, Narciso Garcia A, Menezes Costa LC, van Tulder MW, Lin CC, Machado
25 LA. The McKenzie method for (sub)acute non-specific low back pain. *Cochrane*
26 *Database Syst Rev*. 2023 Apr 5;4(4):CD009711

27
28 Alonso-Sal A, Alonso-Perez JL, Sosa-Reina MD, et al. Effectiveness of Physical Activity
29 in the Management of Nonspecific Low Back Pain: A Systematic Review. *Medicina*
30 (Kaunas). 2024;60(12):2065. Published 2024 Dec 16

31
32 Andersson GB. Epidemiological features of chronic low-back pain. *Lancet*.
33 1999;354(9178):581-585

34
35 Babiloni-Lopez C, Fritz N, Ramirez-Campillo R, Colado JC. Water-Based Exercise in
36 Patients With Nonspecific Chronic Low-Back Pain: A Systematic Review With Meta-
37 Analysis. *J Strength Cond Res*. 2024 Jan 1;38(1):206-219

38
39 Bagg MK, Wand BM, Cashin AG, et al. Effect of Graded Sensorimotor Retraining on Pain
40 Intensity in Patients With Chronic Low Back Pain: A Randomized Clinical Trial.
41 *JAMA*. 2022;328(5):430-439. Doi:10.1001/jama.2022.9930

- 1 Baroncini A, Maffulli N, Manocchio N, et al. Active and passive physical therapy in
 2 patients with chronic low-back pain: a level I Bayesian network meta-analysis. *J*
 3 *Orthop Traumatol.* 2025;26(1):66. Published 2025 Oct 3. doi:10.1186/s10195-025-
 4 00885-4
- 5
- 6 Bogduk N, Jull G. The theoretical pathology of acute locked back. A basis for manipulative
 7 therapy. *23(4):77-81.* 1985
- 8
- 9 Bronfort, G., Haas, M., Evans, R., Kawchuk, G., & Dagenais, S. (2008). Evidence-
 10 informed management of chronic low back pain with spinal manipulation and
 11 mobilization. *The spine journal: official journal of the North American Spine*
 12 *Society, 8(1), 213–225*
- 13
- 14 Broszko C, Golden K, Holmes CR, Fulleborn S, Biglow C, Guthmann R. Is exercise
 15 therapy effective treatment for low back pain? *J Fam Pract.* 2021;70(8):E2-E3. Doi
 16
- 17 Browder DA, Childs JD, Cleland JA, Fritz JM. Effectiveness of an extension-oriented
 18 treatment approach in a subgroup of subjects with low back pain: a randomized clinical
 19 trial. *Phys Ther.* 2007;87(12):1608-1579
- 20
- 21 Brox JI, Storheim K, Grotle M, Tveito TH, Indahl A, Eriksen HR. Systematic review of
 22 back schools, brief education, and fear-avoidance training for chronic low back pain.
 23 *Spine J.* 2008;8(6):948-958
- 24
- 25 Brumitt J, Matheson JW, Meira EP. Core stabilization exercise prescription, part I: current
 26 concepts in assessment and intervention. *Sports Health.* 2013;5(6):504-509
- 27
- 28 Brumitt J, Matheson JW, Meira EP. Core stabilization exercise prescription, part 2: a
 29 systematic review of motor control and general (global) exercise rehabilitation
 30 approaches for patients with low back pain. *Sports Health.* 2013;5(6):510-513
- 31
- 32 Carey TS, Freburger JK. Exercise and the Prevention of Low Back Pain: Ready for
 33 Implementation. *JAMA Intern Med.* 2016;176(2):208-209
- 34
- 35 Cashin AG, Booth J, McAuley JH, et al. Making exercise count: Considerations for the
 36 role of exercise in back pain treatment. *Musculoskeletal Care.* 2022;20(2):259-270.
 37 Doi:10.1002/msc.1597
- 38
- 39 Ceballos-Laita L, Estébanez-de-Miguel E, Jiménez-Rejano JJ, Bueno-Gracia E, Jiménez-
 40 Del-Barrio S. The effectiveness of hip interventions in patients with low-back pain: A
 41 systematic review and meta-analysis. *Braz J Phys Ther.* 2023 Mar-Apr;27(2):100502

- 1 Cheng M, Tian Y, Ye Q, Li J, Xie L, Ding F. Evaluating the effectiveness of six exercise
2 interventions for low back pain: a systematic review and meta-analysis. *BMC*
3 *Musculoskelet Disord.* 2025;26(1):433. Published 2025 May 1. doi:10.1186/s12891-
4 025-08658-0
- 5
- 6 Childs JD, Flynn TW, Fritz JM. A perspective for considering the risks and benefits of
7 spinal manipulation in patients with low back pain. *Man Ther.* 2006;11(4):316-320
8
- 9 Childs JD, Fritz JM, Flynn TW, Irrgang JJ, Johnson KK, Majkowski GR, Delitto A. A
10 clinical prediction rule to identify patients with low back pain most likely to benefit
11 from spinal manipulation: a validation study. *Ann Intern Med.* 2004;141(12):920-928
12
- 13 Chou R, Deyo R, Friedly J, Skelly A, Hashimoto R, Weimer M, Fu R, Dana T, Kraegel P,
14 Griffin J, Grusing S, Brodt E. *Noninvasive Treatments for Low Back Pain.* Agency for
15 Healthcare Research and Quality (US); 2016
16
- 17 Chou R. Low Back Pain. *Ann Intern Med.* 2021;174(8):ITC113-ITC128
18
- 19 Comachio J, Beckenkamp PR, Ho EK, et al. Benefits and harms of exercise therapy and
20 physical activity for low back pain: An umbrella review. *J Sport Health Sci.*
21 2025;14:101038. doi:10.1016/j.jshs.2025.101038
22
- 23 Costa LO, Maher CG, Latimer J, Hodges PW, Herbert RD, Refshauge KM, McAuley JH,
24 Jennings MD. Motor control exercise for chronic low back pain: a randomized placebo-
25 controlled trial. *Phys Ther.* 2009;89(12):1275-1286
26
- 27 De Campos TF, Maher CG, Fuller JT, Steffens D, Attwell S, Hancock MJ. Prevention
28 strategies to reduce future impact of low back pain: a systematic review and meta-
29 analysis. *Br J Sports Med.* 2021;55(9):468-476. Doi:10.1136/bjsports-2019-101436
30
- 31 de Roode A, Heymans MW, van Lankveld W, Staal JB. The impact of contextual effects
32 in exercise therapy for low back pain: a systematic review and meta-analysis. *BMC*
33 *Med.* 2024;22(1):484. Published 2024 Oct 23
34
- 35 Delitto A, George SZ, Van Dillen L, Whitman JM, Sowa G, Shekelle P, Denninger TR,
36 Godges JJ; Orthopaedic Section of the American Physical Therapy Association. Low
37 back pain. *J Orthop Sports Phys Ther.* 2012;42(4):A1-A57
38
- 39 Diez-Buil H, Hernandez-Lucas P, Leirós-Rodríguez R, Echeverría-García O. Effects of the
40 combination of exercise and education in the treatment of low back and/or pelvic pain
41 in pregnant women: Systematic review and meta-analysis. *Int J Gynaecol Obstet.*
42 2024;164(3):811-822

- 1 Drummond C, Lebedeva V, Kirker K, Masaracchio M. Sling Exercise in the Management
2 of Chronic Low Back Pain: A Systematic Review and Meta-Analysis [published online
3 ahead of print, 2021 Sep 23]. *J Strength Cond Res.*
4 2021;10.1519/JSC.0000000000004135
5
- 6 El Melhat AM, Youssef ASA, Zebdawi MR, Hafez MA, Khalil LH, Harrison DE. Non-
7 Surgical Approaches to the Management of Lumbar Disc Herniation Associated with
8 Radiculopathy: A Narrative Review. *J Clin Med.* 2024;13(4):974. Published 2024
9 Feb 8
10
- 11 Essman M, Lin CY. The Role of Exercise in Treating Low Back Pain. *Curr Sports Med*
12 *Rep.* 2022;21(8):267-271. Published 2022 Aug 1
13
- 14 Fernández-Rodríguez R, Álvarez-Bueno C, Cavero-Redondo I, et al. Best Exercise Options
15 for Reducing Pain and Disability in Adults With Chronic Low Back Pain: Pilates,
16 Strength, Core-Based, and Mind-Body. A Network Meta-analysis. *J Orthop Sports*
17 *Phys Ther.* 2022;52(8):505-521
18
- 19 Ferreira, M. L., Ferreira, P. H., Latimer, J., Herbert, R. D., Hodges, P. W., Jennings, M.
20 D., Maher, C. G., & Refshauge, K. M. (2007). Comparison of general exercise, motor
21 control exercise and spinal manipulative therapy for chronic low back pain: A
22 randomized trial. *Pain, 131*(1-2), 31–37
23
- 24 Ferreira ML, Ferreira PH, Latimer J, Herbert RD, Hodges PW, Jennings MD, Maher CG,
25 Refshauge KM. Comparison of general exercise, motor control exercise and spinal
26 manipulative therapy for chronic low back pain: A randomized trial. *Pain.* 2007;131(1-
27 2):31–37
28
- 29 Fleckenstein J, Floessel P, Engel T, et al. Individualized Exercise in Chronic Non-Specific
30 Low Back Pain: A Systematic Review with Meta-Analysis on the Effects of Exercise
31 Alone or in Combination with Psychological Interventions on Pain and Disability. *J*
32 *Pain.* 2022;23(11):1856-1873
33
- 34 Flynn T, Fritz J, Whitman J, Wainner R, Magel J, Rendeiro D, Butler B, Garber M, Allison
35 S. A clinical prediction rule for classifying patients with low back pain who
36 demonstrate short-term improvement with spinal manipulation. *Spine.*
37 2002;27(24):2835–2843
38
- 39 Foster NE, Anema JR, Cherkin D, Chou R, Cohen SP, Gross DP, Ferreira PH, Fritz JM,
40 Koes BW, Peul W, Turner JA, Maher CG; Lancet Low Back Pain Series Working
41 Group. Prevention and treatment of low back pain: evidence, challenges, and promising
42 directions. *Lancet.* 2018;391(10137):2368–2383

- 1 Fritz JM, Childs JD, Flynn TW. Pragmatic application of a clinical prediction rule in
 2 primary care to identify patients with low back pain with a good prognosis following a
 3 brief spinal manipulation intervention. *BMC Fam Pract.* 2005;6(1):29
 4
- 5 García-Moreno JM, Calvo-Muñoz I, Gómez-Conesa A, López-López JA. Effectiveness of
 6 physiotherapy interventions for back care and the prevention of non-specific low back
 7 pain in children and adolescents: a systematic review and meta-analysis. *BMC*
 8 *Musculoskelet Disord.* 2022;23(1):314. Published 2022 Apr 2
 9
- 10 García-Moreno JM, Calvo-Muñoz I, Gómez-Conesa A, López-López JA. Therapeutic
 11 Exercise is Effective in Reducing the Intensity of Nonspecific Low Back Pain in
 12 Children and Adolescents: A Systematic Review and Network Meta-analysis. *Arch*
 13 *Phys Med Rehabil.* 2025;106(5):771-780. doi:10.1016/j.apmr.2024.11.002
 14
- 15 Gatchel RJ, Mayer TG. Evidence-informed management of chronic low back pain with
 16 functional restoration. *Spine J.* 2008;8(1):65-69
 17
- 18 Gatchel RJ, Rollings KH. Evidence-informed management of chronic low back pain with
 19 cognitive behavioral therapy. *Spine J.* 2008;8(1):40-44
 20
- 21 George SZ, Fritz JM, Silfies SP, et al. Interventions for the Management of Acute and
 22 Chronic Low Back Pain: Revision 2021. *J Orthop Sports Phys Ther.*
 23 2021;51(11):CPG1-CPG60
 24
- 25 Gianola S, Barger S, Del Castillo G, et al. Effectiveness of treatments for acute and
 26 subacute mechanical non-specific low back pain: a systematic review with network
 27 meta-analysis. *Br J Sports Med.* 2022;56(1):41-50
 28
- 29 Gilanyi YL, Wewege MA, Shah B, Cashin AG, Williams CM, Davidson SRE, McAuley
 30 JH, Jones MD. Exercise Increases Pain Self-efficacy in Adults With Nonspecific
 31 Chronic Low Back Pain: A Systematic Review and Meta-analysis. *J Orthop Sports*
 32 *Phys Ther.* 2023 Jun;53(6):335–342
 33
- 34 Gilliam JR, George SZ, Norman KS, Hendren S, Sahu PK, Silfies SP. Mind-Body Exercise
 35 Performed by Physical Therapists for Reducing Pain and Disability in Low Back Pain:
 36 A Systematic Review With Meta-analysis. *Arch Phys Med Rehabil.* 2023
 37 May;104(5):776-789
 38
- 39 González-Gómez L, Moral-Munoz JA, Rosales-Tristancho A, Cuevas-Moreno A,
 40 Cardellat-González M, Rodríguez-Domínguez ÁJ. Exercise Therapy Versus Manual
 41 Therapy for the Management of Pain Intensity, Disability, and Physical Function in

- 1 People With Chronic Low Back Pain: A Systematic Review With Meta-Analysis and
2 Meta-Regression. *Eur J Pain*. 2025;29(8):e70090. doi:10.1002/ejp.70090
3
- 4 Gou Y, Lei H, Chen X, Wang X. The effects of hamstring stretching exercises on pain
5 intensity and function in low back pain patients: A systematic review with meta-
6 analysis of randomized controlled trials. *SAGE Open Med*.
7 2024;12:20503121241252251. Published 2024 Jul 29
8
- 9 Grande GHD, Vidal RVC, Salini MCR, Christofaro DGD, Oliveira CB. Barriers and
10 Facilitators to Physical Activity and Exercise Among People With Chronic Low Back
11 Pain: A Qualitative Evidence Synthesis. *J Orthop Sports Phys Ther*. 2025;55(5):312-
12 330. doi:10.2519/jospt.2025.12905
13
- 14 Grooten WJA, Boström C, Dederig Å, et al. Summarizing the effects of different exercise
15 types in chronic low back pain – a systematic review of systematic reviews. *BMC*
16 *Musculoskelet Disord*. 2022;23(1):801. Published 2022 Aug 22
17
- 18 Guzmán J, Esmail R, Karjalainen K, Malmivaara A, Irvin E, Bombardier C.
19 Multidisciplinary rehabilitation for chronic low back pain: systematic review. *BMJ*.
20 2001;322(7301):1511-1516
21
- 22 Haldeman S, Dagenais S. What have we learned about the evidence-informed management
23 of chronic low back pain? *Spine J*. 2008;8(1):266-277
24
- 25 Hayden JA, van Tulder MW, Malmivaara A, Koes BW. Exercise therapy for treatment of
26 non-specific low back pain. *Cochrane Database Syst Rev*. 2005 Jul 20;(3):CD000335
27
- 28 Hayden JA, Wilson MN, Stewart S, Cartwright JL, Smith AO, Riley RD, van Tulder M,
29 Bendix T, Cecchi F, Costa LOP, Dufour N, Ferreira ML, Foster NE, Gudavalli MR,
30 Hartvigsen J, Helmhout P, Kool J, Koumantakis GA, Kovacs FM, Kuukkanen T, Long
31 A, Macedo LG, Machado LAC, Maher CG, Mehling W, Morone G, Peterson T,
32 Rasmussen-Barr E, Ryan CG, Sjögren T, Smeets R, Staal JB, Unsgaard-Tøndel M,
33 Wajswelner H, Yeung EW; Chronic Low Back Pain IPD Meta-Analysis Group.
34 Exercise treatment effect modifiers in persistent low back pain: an individual
35 participant data meta-analysis of 3514 participants from 27 randomised controlled
36 trials. *Br J Sports Med*. 2020 Nov;54(21):1277-1278
37
- 38 Hayden JA, Ellis J, Ogilvie R, Malmivaara A, van Tulder MW. Exercise therapy for
39 chronic low back pain. *Cochrane Database Syst Rev*. 2021a;9(9):CD009790. Published
40 2021 Sep 28

- 1 Hayden JA, Ellis J, Ogilvie R, et al. Some types of exercise are more effective than others
2 in people with chronic low back pain: a network meta-analysis. *J Physiother.*
3 2021b;67(4):252-262
4
- 5 Heidari F, Mohammad Rahimi N, Aminzadeh R. Aquatic Exercise Impact on Pain
6 Intensity, Disability and Quality of Life in Adults with Low Back Pain: A Systematic
7 Review and Meta-analysis. *Biol Res Nurs.* 2023 Oct;25(4):527-541
8
- 9 Hennemann V, Ziegelmann PK, Marcolino MAZ, Duncan BB. The McKenzie Method
10 delivered by credentialed therapists for chronic low back pain with directional
11 preference: systematic review with meta-analysis. *J Man Manip Ther.* 2025;33(2):96-
12 111. doi:10.1080/10669817.2024.2408084
13
- 14 Herrero P, Val P, Lapuente-Hernández D, Cuenca-Zaldívar JN, Calvo S, Gómez-Trullén
15 EM. Effects of Lifestyle Interventions on the Improvement of Chronic Non-Specific
16 Low Back Pain: A Systematic Review and Network Meta-Analysis. *Healthcare*
17 (Basel). 2024;12(5):505. Published 2024 Feb 20
18
- 19 Hicks GE, Fritz JM, Delitto A, McGill SM. Preliminary development of a clinical
20 prediction rule for determining which patients with low back pain will respond to a
21 stabilization exercise program. *Arch Phys Med Rehabil.* 2005;86(9):1753-1762
22
- 23 Hides JA, Jull GA, Richardson CA. Long-term effects of specific stabilizing exercises for
24 first-episode low back pain. *Spine.* 2001;26(11):E243-E248
25
- 26 Hlaing SS, Puntumetakul R, Khine EE, Boucaut R. Effects of core stabilization exercise
27 and strengthening exercise on proprioception, balance, muscle thickness and pain
28 related outcomes in patients with subacute nonspecific low back pain: a randomized
29 controlled trial. *BMC Musculoskelet Disord.* 2021;22(1):998. Published 2021 Nov 30
30
- 31 Hoffman BM, Papas RK, Chatkoff DK, Kerns RD. Meta-analysis of psychological
32 interventions for chronic low back pain. *Health Psychol.* 2007;26(1):1-9
33
- 34 IJzelenberg W, Oosterhuis T, Hayden JA, Koes BW, van Tulder MW, Rubinstein SM, de
35 Zoete A. Exercise therapy for treatment of acute non-specific low back pain. *Cochrane*
36 *Database Syst Rev.* 2023 Aug 30;8(8):CD009365
37
- 38 IJzelenberg W, Oosterhuis T, Hayden JA, et al. Exercise Therapy for Treatment of Acute
39 Non-specific Low Back Pain: A Cochrane Systematic Review and Meta-analysis of
40 Randomized Controlled Trials. *Arch Phys Med Rehabil.* 2024;105(8):1571-1584

- 1 Jangra P, Kaur J, Malik M, Rani M. A systematic review and meta-analysis of randomized
2 controlled trials on the effect of various therapeutic interventions on sacroiliac joint
3 dysfunction. *J Bodyw Mov Ther.* 2025;45:627-638. doi:10.1016/j.jbmt.2025.09.019
4
- 5 Jenkins HJ, Corrêa L, Brown BT, et al. Long-term effectiveness of non-surgical
6 interventions for chronic low back pain: a systematic review and meta-analysis. *Lancet*
7 *Rheumatol.* 2025;7(9):e607-e617. doi:10.1016/S2665-9913(25)00064-5
8
- 9 Jin H, Kwon J, Lee J, Lee DJ, Lee DH, Jeon JY. Effects of Exercise on Pain and Disability
10 After Lumbar Disc Herniation Surgery: A Systematic Review and Meta-analysis of
11 Randomized Controlled Trials. *Am J Phys Med Rehabil.* 2025;104(6):511-518.
12 doi:10.1097/PHM.0000000000002658
13
- 14 Johannsen F, Remvig L, Kryger P, et al. Exercises for chronic low back pain: a clinical
15 trial. *J Orthop Sports Phys Ther.* 1995;22(2):52-59
16
- 17 Jones KC, Tocco EC, Marshall AN, Valovich McLeod TC, Welch Bacon CE. Pain
18 education with therapeutic exercise in chronic nonspecific low back pain rehabilitation:
19 a critically appraised topic. *J Sport Rehabil.* 2020;29(8):1204-1209
20
- 21 Jones MD, Hansford HJ, Bastianon A, et al. Exercise adherence is associated with
22 improvements in pain intensity and functional limitations in adults with chronic non-
23 specific low back pain: a secondary analysis of a Cochrane review. *J Physiother.*
24 2025;71(2):91-99. doi:10.1016/j.jphys.2025.03.004
25
- 26 Jordan A, Bendix T, Nielsen H, Hansen FR, Høst D, Winkel A. Intensive training,
27 physiotherapy, or manipulation for patients with chronic neck pain: a prospective,
28 single-blinded, randomized clinical trial. *Spine.* 1998;23(3):311-319
29
- 30 Joyce CJ, Nowell K, Boling MC, Buckley BD. The Effects of Exercise and Manual
31 Manipulation on Sacroiliac Joint Dysfunction: A Critically Appraised Topic. *J Sport*
32 *Rehabil.* 2025;34(6):682-689. Published 2025 Mar 25. doi:10.1123/jsr.2024-0205
33
- 34 Karlsson M, Bergenheim A, Larsson MEH, Nordeman L, van Tulder M, Bernhardsson S.
35 Effects of exercise therapy in patients with acute low back pain: a systematic review of
36 systematic reviews. *Syst Rev.* 2020 Aug 14;9(1):182
37
- 38 Kazeminia M, Rajati F, Rajati M. The effect of pelvic floor muscle-strengthening exercises
39 on low back pain: a systematic review and meta-analysis on randomized clinical trials.
40 *Neurol Sci.* 2023 Mar;44(3):859-872

- 1 Kofotolis N, Kellis E. Effects of two 4-week proprioceptive neuromuscular facilitation
2 programs on muscle endurance, flexibility, and functional performance in women with
3 chronic low back pain. *Physical Therapy*. 2006;86(7):1001–1012
4
- 5 Koumantakis GA, Watson PJ, Oldham JA. Supplementation of general endurance exercise
6 with randomized training versus general exercise only. Physiological and functional
7 outcomes of a randomized controlled trial of patients with recurrent low back pain.
8 *Clinical Biomechanics (Bristol, Avon)*. 2005;20(5):474–482
9
- 10 Kuukkanen TM, Mälkiä EA. An experimental controlled study on postural sway and
11 therapeutic exercise in subjects with low back pain. *Clinical Rehabilitation*.
12 2000;14(2):192–202
13
- 14 Lehtola V, Luomajoki H, Leinonen V, Gibbons S, Airaksinen O. Sub-classification based
15 specific movement control exercises are superior to general exercise in sub-acute low
16 back pain when both are combined with manual therapy: A randomized controlled trial.
17 *BMC Musculoskeletal Disorders*. 2016;17:135
18
- 19 Li Y, Yan L, Hou L, Zhang X, Zhao H, Yan C, Li X, Li Y, Chen X, Ding X. Exercise
20 intervention for patients with chronic low back pain: a systematic review and network
21 meta-analysis. *Front Public Health*. 2023 Nov 17;11:1155225
22
- 23 Liang Z, Tian S, Wang C, et al. The Best Exercise Modality and Dose for Reducing Pain
24 in Adults With Low Back Pain: A Systematic Review With Model-Based Bayesian
25 Network Meta-analysis. *J Orthop Sports Phys Ther*. 2024;54(5):315-327
26
- 27 Liddle SD, Baxter GD, Gracey JH. Exercise and chronic low back pain: what works? *Pain*.
28 2004;107(1-2):176–190
29
- 30 Lim Y, Do Y, Lee SH, Lee H. Efficacy of pelvic floor muscle training with physical therapy
31 for low back pain: A systematic review and meta-analysis. *Clin Rehabil*.
32 2024;38(12):1590-1608
33
- 34 Lim TH, Mak HY, Man Ngai SM, et al. Nonpharmacological Spine Pain Management in
35 Clinical Practice Guidelines: A Systematic Review Using AGREE II and AGREE-
36 REX Tools. *J Orthop Sports Phys Ther*. 2025;55(1):12-25
37
- 38 Lindberg B, Leggit JC. Effectiveness of Exercise Therapy in Patients With Chronic Low
39 Back Pain. *Am Fam Physician*. 2022;106(4):380-381

- 1 Macedo LG, Saragiotto BT, Yamato TP, Costa LO, Menezes Costa LC, Ostelo RW, Maher
 2 CG. Motor control exercise for acute non-specific low back pain. *Cochrane Database*
 3 *Syst Rev.* 2016;2:CD012085
 4
- 5 Manniche C, Jordan A. The value of exercise therapy. *Spine.* 1995;20(11):1221–1222.
 6
- 7 May S, Donelson R. Evidence-informed management of chronic low back pain with the
 8 McKenzie method. *Spine J.* 2008;8(1):134–141
 9
- 10 Mayer J, Mooney V, Dagenais S. Evidence-informed management of chronic low back
 11 pain with lumbar extensor strengthening exercises. *Spine J.* 2008;8(1):96–113
 12
- 13 Nava-Bringas TI, Romero-Fierro LO, Trani-Chagoya YP, et al. Stabilization Exercises
 14 Versus Flexion Exercises in Degenerative Spondylolisthesis: A Randomized
 15 Controlled Trial. *Phys Ther.* 2021;101(8):pzab108
 16
- 17 Niederer D, Weippert M, Behrens M. What Modifies the Effect of an Exercise Treatment
 18 for Chronic Low Back Pain? A Meta-epidemiologic Regression Analysis of Risk of
 19 Bias and Comparative Effectiveness. *J Orthop Sports Phys Ther.* 2022;52(12):792-802
 20
- 21 Niederer D, Fleckenstein J, Floessel P, et al. Comparative Effectiveness of Tailored
 22 Exercise Therapies Alone or Combined With Psychological Interventions for Chronic,
 23 Nonspecific Low Back Pain: A Systematic Review With Network Meta-analysis. *J*
 24 *Orthop Sports Phys Ther.* 2026;56(1):16-27. doi:10.2519/jospt.2025.13281
 25
- 26 Ostelo RW, van Tulder MW, Vlaeyen JW, Linton SJ, Morley SJ, Assendelft WJ.
 27 Behavioural treatment for chronic low-back pain. *Cochrane Database Syst Rev.*
 28 2005;(1):CD002014
 29
- 30 Owen PJ, Miller CT, Mundell NL, Verswijveren SJJM, Tagliaferri SD, Brisby H, Bowe
 31 SJ, Belavy DL. Which specific modes of exercise training are most effective for
 32 treating low back pain? Network meta-analysis. *Br J Sports Med.* 2020
 33 Nov;54(21):1279-1287
 34
- 35 Patti A, Thornton JS, Giustino V, et al. Effectiveness of Pilates exercise on low back pain:
 36 a systematic review with meta-analysis. *Disabil Rehabil.* 2024;46(16):3535-3548
 37
- 38 Philadelphia Panel. Philadelphia Panel evidence-based clinical practice guidelines on
 39 selected rehabilitation interventions for low back pain. *Phys Ther.* 2001;81(10):1641-
 40 1674

- 1 Prat-Luri A, de Los Rios-Calonge J, Moreno-Navarro P, Manresa-Rocamora A, Vera-
 2 Garcia FJ, Barbado D. Effect of Trunk-Focused Exercises on Pain, Disability, Quality
 3 of Life, and Trunk Physical Fitness in Low Back Pain and How Potential Effect
 4 Modifiers Modulate Their Effects: A Systematic Review With Meta-analyses. *J Orthop
 5 Sports Phys Ther.* 2023;53(2):64-93
 6
- 7 Qaseem A, Wilt TJ, McLean RM, Forcica MA; Clinical Guidelines Committee of the
 8 American College of Physicians. Noninvasive Treatments for Acute, Subacute, and
 9 Chronic Low Back Pain: A Clinical Practice Guideline From the American College of
 10 Physicians. *Ann Intern Med.* 2017;166(7):514-530
 11
- 12 Quentin C, Bagheri R, Ugbolue UC, et al. Effect of Home Exercise Training in Patients
 13 with Nonspecific Low-Back Pain: A Systematic Review and Meta-Analysis. *Int J
 14 Environ Res Public Health.* 2021;18(16):8430. Published 2021 Aug 10
 15
- 16 Ram AK, Summers SJ, Booth J, Gibbs MT, Jones MD. Higher intensity exercise reduces
 17 disability more than lower intensity exercise in adults with chronic low back pain: A
 18 systematic review and meta-analysis. *Musculoskeletal Care.* 2023 Sep;21(3):611-622
 19
- 20 Rathnayake APS, Sparkes V, Sheeran L. What is the effect of low back pain self-
 21 management interventions with exercise components added? A systematic review with
 22 meta-analysis. *Musculoskelet Sci Pract.* 2021;56:102469
 23
- 24 Rizzo RR, Cashin AG, Wand BM, et al. Non-pharmacological and non-surgical treatments
 25 for low back pain in adults: an overview of Cochrane reviews. *Cochrane Database Syst
 26 Rev.* 2025;3(3):CD014691. Published 2025 Mar 27.
 27 doi:10.1002/14651858.CD014691.pub2
 28
- 29 Roren A, Daste C, Coleman M, Rannou F, Freyssenet D, Moro C, Lefèvre-Colau MM,
 30 Nguyen C. Physical activity and low back pain: A critical narrative review. *Ann Phys
 31 Rehabil Med.* 2023 Mar;66(2):101650
 32
- 33 Saragiotto BT, Maher CG, Yamato TP, Costa LO, Menezes Costa LC, Ostelo RW, Macedo
 34 LG. Motor control exercise for chronic non-specific low-back pain. *Cochrane Database
 35 Syst Rev.* 2016;(1):CD012004
 36
- 37 Salehi S, Sobhani V, Mir SM, Keivanfar N, Shamsoddini A, Hashemi SE. Efficacy of
 38 specific exercises in general population with non-specific low back pain: A systematic
 39 review and meta-analysis of randomized controlled trials. *J Bodyw Mov Ther.*
 40 2024;39:673-705

- 1 Sherman KJ, Cherkin DC, Erro J, Miglioretti DL, Deyo RA. Comparing yoga, exercise,
2 and a self-care book for chronic low back pain: a randomized, controlled trial. *Ann*
3 *Intern Med.* 2005;143(12):849-856
4
- 5 Shi Z, Zhou H, Lu L, Pan B, Wei Z, Yao X, Kang Y, Liu L, Feng S. Aquatic Exercises in
6 the Treatment of Low Back Pain: A Systematic Review of the Literature and Meta-
7 Analysis of Eight Studies. *Am J Phys Med Rehabil.* 2018;97(2):116-122
8
- 9 Shiri R, Coggon D, Falah-Hassani K. Exercise for the Prevention of Low Back Pain:
10 Systematic Review and Meta-Analysis of Controlled Trials. *Am J Epidemiol.*
11 2018;187(5):1093-1101
12
- 13 Skelly AC, Chou R, Dettori JR, Turner JA, Friedly JL, Rundell SD, Fu R, Brodt ED,
14 Wasson N, Kantner S, Ferguson AJR. Noninvasive Nonpharmacological Treatment for
15 Chronic Pain: A Systematic Review Update [Internet]. Rockville (MD): Agency for
16 Healthcare Research and Quality (US); 2020 Apr. Report No.: 20-EHC009. PMID:
17 32338846
18
- 19 Slade SC, Keating JL. Unloaded movement facilitation exercise compared to no exercise
20 or alternative therapy on outcomes for people with nonspecific chronic low back pain:
21 a systematic review. *J Manipulative Physiol Ther.* 2007;30(4):301-311
22
- 23 Standaert CJ, Weinstein SM, Rumpeltes J. Evidence-informed management of chronic low
24 back pain with lumbar stabilization exercises. *Spine J.* 2008;8(1):114-120
25
- 26 Stuber KJ, Bruno P, Sajko S, Hayden JA. Core stability exercises for low back pain in
27 athletes: a systematic review of the literature. *Clin J Sport Med.* 2014;24(6):448-456
28
- 29 Suh JH, Kim H, Jung GP, Ko JY, Ryu JS. The effect of lumbar stabilization and walking
30 exercises on chronic low back pain: A randomized controlled trial. *Medicine*
31 *(Baltimore).* 2019;98(26):e16173
32
- 33 Thornton JS, Caneiro JP, Hartvigsen J, et al. Treating low back pain in athletes: a
34 systematic review with meta-analysis. *Br J Sports Med.* 2021;55(12):656-662
35
- 36 Tikhile P, Patil DS. Unveiling the Efficacy of Physiotherapy Strategies in Alleviating Low
37 Back Pain: A Comprehensive Review of Interventions and Outcomes. *Cureus.*
38 2024;16(3):e56013. Published 2024 Mar 12
39
- 40 Traeger AC, Qaseem A, McAuley JH. Low Back Pain. *JAMA.* 2021;326(3):286

- 1 van Dillen LR, Lanier VM, Steger-May K, et al. Effect of Motor Skill Training in
 2 Functional Activities vs Strength and Flexibility Exercise on Function in People With
 3 Chronic Low Back Pain: A Randomized Clinical Trial [published correction appears
 4 in JAMA Neurol. 2021 Jan 19;:]. JAMA Neurol. 2021;78(4):385-395
 5
- 6 van Tulder MW, Ostelo R, Vlaeyen JW, Linton SJ, Morley SJ, Assendelft WJ. Behavioral
 7 treatment for chronic low back pain: a systematic review within the framework of the
 8 Cochrane Back Review Group. Spine. 2000;25(20):2688-2699
 9
- 10 Vanti C, Andreatta S, Borghi S, Guccione AA, Pillastrini P, Bertozzi L. The effectiveness
 11 of walking versus exercise on pain and function in chronic low back pain: a systematic
 12 review and meta-analysis of randomized trials. Disability and Rehabilitation.
 13 2019;41(6):622-632
 14
- 15 Vega-Retuerta N, Sánchez-Parente S, Segura-Jiménez V. Effectiveness of
 16 multidisciplinary approaches including exercise to treat non-specific chronic low back
 17 pain: A systematic review and meta-analysis across multiple regions. *Prev Med*.
 18 2025;199:108381. doi:10.1016/j.ypmed.2025.108381
 19
- 20 Verville L, Ogilvie R, Hincapié CA, Southerst D, Yu H, Bussièrès A, Gross DP, Pereira P,
 21 Mior S, Tricco AC, Cedraschi C, Brunton G, Nordin M, Connell G, Wong JJ, Shearer
 22 HM, Lee JGB, Wang D, Hayden JA, Cancelliere C. Systematic Review to Inform a
 23 World Health Organization (WHO) Clinical Practice Guideline: Benefits and Harms of
 24 Structured Exercise Programs for Chronic Primary Low Back Pain in Adults. *J Occup
 25 Rehabil*. 2023 Dec;33(4):636-650
 26
- 27 Wai EK, Rodriguez S, Dagenais S, Hall H. Evidence-informed management of chronic low
 28 back pain with physical activity, smoking cessation, and weight loss. *Spine J*.
 29 2008;8(1):195-202
 30
- 31 Wang XQ, Zheng JJ, Yu ZW, Bi X, Lou SJ, Liu J, Cai B, Hua YH, Wu M, Wei ML, Shen
 32 HM, Chen Y, Pan YJ, Xu GH, Chen PJ. A meta-analysis of core stability exercise
 33 versus general exercise for chronic low back pain. *PLoS One*. 2012;7(12):e52082
 34
- 35 Wewege MA, Booth J, Parmenter BJ. Aerobic vs resistance exercise for chronic non-
 36 specific low back pain: a systematic review and meta-analysis. *J Back Musculoskelet
 37 Rehabil*. 2018;31(5):889-899
 38
- 39 Whitman JM, Flynn TW, Childs JD, Wainner RS, Gill HE, Ryder MG, Garber MB,
 40 Bennett AC, Fritz JM. A comparison between two physical therapy treatment programs
 41 for patients with lumbar spinal stenosis: a randomized clinical trial. *Spine*.
 42 2006;31(22):2541-2549

- 1 Wong CM, Rugg B, Geere JA. The effects of Pilates exercise in comparison to other forms
 2 of exercise on pain and disability in individuals with chronic non-specific low back
 3 pain: A systematic review with meta-analysis. *Musculoskeletal Care*. 2023
 4 Mar;21(1):78-96
- 5
 6 World Health Organization. Low back pain. Published 2023. Accessed January 20, 2026.
 7 <https://www.who.int/news-room/fact-sheets/detail/low-back-pain>
 8
- 9 Yamato TP, Maher CG, Saragiotto BT, Hancock MJ, Ostelo RW, Cabral CM, Menezes
 10 Costa LC, Costa LO. Pilates for low back pain. *Cochrane Database Syst Rev*.
 11 2015;(7):CD010265
- 12
 13 Xia Y, Hu Y, Ouyang J, Huang R. Four MBE Therapies to Improve Pain Scores, Physical
 14 Functioning, and Quality of Life in Patients With Chronic Nonspecific Low Back Pain:
 15 Network Meta-Analysis of Randomised Controlled Trials. *Eur J Pain*.
 16 2025;29(8):e70092. doi:10.1002/ejp.70092
- 17
 18 Yu Z, Yin Y, Wang J, Zhang X, Cai H, Peng F. Efficacy of Pilates on Pain, Functional
 19 Disorders and Quality of Life in Patients with Chronic Low Back Pain: A Systematic
 20 Review and Meta-Analysis. *Int J Environ Res Public Health*. 2023 Feb 6;20(4):2850
 21
- 22 Zaina F, Côté P, Cancelliere C, Di Felice F, Donzelli S, Rauch A, Verville L, Negrini S,
 23 Nordin M. A Systematic Review of Clinical Practice Guidelines for Persons With Non-
 24 specific Low Back Pain With and Without Radiculopathy: Identification of Best
 25 Evidence for Rehabilitation to Develop the WHO's Package of Interventions for
 26 Rehabilitation. *Arch Phys Med Rehabil*. 2023 Nov;104(11):1913-1927
 27
- 28 Zhang SK, Gu ML, Zhang T, Xu H, Mao SJ, Zhou WS. Effects of exercise therapy on
 29 disability, mobility, and quality of life in the elderly with chronic low back pain: a
 30 systematic review and meta-analysis of randomized controlled trials. *J Orthop Surg*
 31 *Res*. 2023 Jul 19;18(1):513
- 32
 33 Zhao K, Zhang P, Li H, Li L. Exercise prescription for improving chronic low back pain
 34 in adults: a network meta-analysis. *Front Public Health*. 2025;13:1512450. Published
 35 2025 May 30. doi:10.3389/fpubh.2025.1512450
- 36
 37 Zhou T, Salman D, McGregor AH. Recent clinical practice guidelines for the management
 38 of low back pain: a global comparison. *BMC Musculoskelet Disord*. 2024;25(1):344.
 39 Published 2024 May 1
- 40
 41 Zhu F, Zhang M, Wang D, Hong Q, Zeng C, Chen W. Yoga compared to non-exercise or
 42 physical therapy exercise on pain, disability, and quality of life for patients with chronic

- 1 low back pain: A systematic review and meta-analysis of randomized controlled trials.
- 2 PLoS One. 2020 Sep 1;15(9):e0238544