Clinical Practice Guideline:	Spinal Manipulative Therapy for Treatment of Children and Infants
Date of Implementation:	July 16, 2009
Product:	Specialty
	Related Policies: CPG 119: Spinal Manipulative Therapy for Non-Musculoskeletal and Related Disorders for more specific information. CPG 135: Physical Therapy Medical Policy/Guideline CPG 155: Occupational Therapy Medical Policy/Guideline CPG 278: Chiropractic Services CPG 285: Spinal Manipulative Therapy (SMT) for Musculoskeletal and Related Disorders

17 Table of Contents

18	GUIDELINES	. 1
19	EVIDENCE REVIEW	. 2
20	SAFETY	. 9
21	PRACTITIONER SCOPE AND TRAINING	11
22	References	11

2324 GUIDELINES

American Specialty Health – Specialty (ASH) considers Spinal Manipulation or Mobilization for the treatment of children and infants (age 14 and under) to be medically necessary when the documentation establishes a valid diagnosis and symptomatic status and satisfies the criteria outlined in the *Spinal Manipulative Therapy (SMT) for Musculoskeletal and Related Disorders (CPG 285-S)* clinical practice guideline.

30

While the literature is insufficient to conclude spinal manipulative therapy in children with 31 spinal pain is either clinically effective or ineffective as the evidence is generally very 32 limited and low quality; it may be reasonable to infer from the literature supporting SMT 33 for low back pain and neck pain in adults that there is a similar benefit in children. To the 34 extent to which children are physiologically and bio-mechanically different from adults, 35 there may be an impact of those differences on the benefit: risk profile. It is likely that these 36 differences are greater in infants and children than in adolescents, thus additional caution 37 should be considered prior to performing SMT on infants and children. 38

Page 1 of 16

Spinal manipulation is considered not medically necessary when the above criteria are not 1 2 met.

3

Spinal manipulation is considered not medically necessary for non-musculoskeletal and 4 related disorders (e.g., asthma, infantile colic, nocturnal enuresis, or otitis media) in 5 children. Moreover, ASH clinical committees have determined that SMT for non-6 musculoskeletal and related disorders in children poses a health and safety risk through 7 substitution harm. See the Spinal Manipulative Therapy for Non-Musculoskeletal 8 Conditions and Related Disorders (CPG 119-S) clinical practice guideline for more 9 specific information. 10

11

EVIDENCE REVIEW 12

The literature on spinal manipulation in children shows potential beneficial effects in a few 13 conditions, though the evidence is generally very limited and of low quality. The evidence 14 base for the provision of SMT in children for spine pain is largest in the treatment of 15 adolescent idiopathic scoliosis, and low back pain. However, for both of these conditions, 16 there is insufficient evidence to support a conclusion regarding the effectiveness of SMT. 17

18

SMT for low back pain was the focus of a study by Hayden et al. (2003). The study 19 20 involved a prospective cohort of 54 consecutive 4- to 18-year-old low-back-pain patients from 15 randomly selected chiropractors in Calgary, Alberta, and Toronto, Ontario, 21 Canada. Patients presented primarily with uncomplicated mechanical low-back pain of less 22 than three months duration, had a median of five visits (interquartile range 3 to 8) over a 23 median treatment period of 22 days (interquartile range 7 to 56) and were treated most 24 commonly with SMT (95.2%) and/or passive manual therapy (42.9%). Over the course of 25 treatment, 90.7% of patients improved; 81.4% improved by more than 20% on the pediatric 26 visual analogue scale (VAS), and 53.7% had "important" improvement (defined as the 27 median change on the VAS in 78.9% of patients who reported that they were "much 28 improved.") 92.3% of patients reported improvement of some kind. Those with pain for 29 more than 12 weeks at the beginning of treatment were less likely to improve within the 30 first five visits (RR = 2.1; 95% CI = 1.1, 4.3), whereas those with restricted range of motion 31 at baseline were more likely to improve (RR = 0.39; 95% CI = 0.21, 0.75). No 32 33 complications or adverse events were reported. Because the study lacked a comparison group, no conclusions can be drawn about the efficacy or relative effectiveness of SMT for 34 pediatric patients with low-back pain. A systematic review by Vaughn et al. (2012) only 35 identified two RCTs and two prospective cohort studies in their literature search. They did 36 not include any research studying SMT and excessive spinal curvatures. Authors concluded 37 that given the paucity of data in the literature to support or refute using SMT for pediatric 38 39 patients with spinal conditions, further research is necessary to recommend the use of this intervention in children. For a discussion of the research supporting SMT in adults for the 40 treatment of low back pain and neck pain, see the Spinal Manipulative Therapy (SMT) for 41 Musculoskeletal and Related Disorders (CPG 285-S) clinical practice guideline. 42

Page 2 of 16

CPG 120 Revision 16 - S Spinal Manipulative Therapy for Treatment of Children and Infants Revised – March 20, 2025 To CQT for review 02/10/2025 CQT reviewed 02/10/2025 To QIC for review and approval 03/04/2025 QIC reviewed and approved 03/04/2025 To QOC for review and approval 03/20/2025 QOC reviewed and approved 03/20/2025

Evans et al. (2018) conducted a multicenter randomized trial comparing 12 weeks of spinal 1 manipulative therapy (SMT) combined with exercise therapy (ET) to ET alone for low 2 back pain. Participants were 185 adolescents aged 12 to 18 years with chronic LBP. The 3 primary outcome was LBP severity at 12, 26, and 52 weeks. Secondary outcomes included 4 disability, quality of life, medication use, patient- and caregiver-rated improvement, and 5 satisfaction. Of the 185 enrolled patients, 179 (97%) provided data at 12 weeks and 174 6 (94%) at 26 and 52 weeks. Adding SMT to ET resulted in a significantly larger reduction 7 in LBP severity over the course of 1 year. The group difference in LBP severity (0-10 8 scale) was small at the end of treatment but was larger at weeks 26 and 52. At 26 weeks, 9 SMT with ET performed better than ET alone for disability and improvement. The SMT 10 11 with ET group reported significantly greater satisfaction with care at all time points. There were no serious treatment-related adverse events. For adolescents with chronic LBP, spinal 12 manipulation combined with exercise was more effective than exercise alone over a 1-year 13 period, with the largest differences occurring at 6 months. These findings warrant 14 replication and evaluation of cost effectiveness. 15

16

Dissing et al. (2018) investigated the effectiveness of adding manipulative therapy to other 17 conservative care for spinal pain in a school-based cohort of Danish children aged 9-15 18 years. A text message system and clinical examinations were used for data collection. 19 20 Interventions included either (1) advice, exercises, and soft-tissue treatment; or (2) advice, exercises and soft-tissue treatment plus manipulative therapy. The primary outcome was 21 number of recurrences of spinal pain. Secondary outcomes were duration of spinal pain. 22 change in pain intensity and Global Perceived Effect. Authors found no significant 23 difference between groups in the primary outcome and intervention group 2. Children in 24 the group receiving manipulative therapy reported a higher Global Perceived Effect. No 25 adverse events were reported. Main limitations are the potential discrepancy between 26 parental and child reporting and that the study population may not be comparable to a 27 normal care-seeking population. Authors concluded that adding manipulative therapy to 28 other conservative care in school children with spinal pain did not result in fewer recurrent 29 episodes. The choice of treatment-if any-for spinal pain in children therefore relies on 30 personal preferences and could include conservative care with and without manipulative 31 therapy. Participants in this trial may differ from a normal care-seeking population. 32

33

Dissing et al. (2019) acknowledged that interventions may be more effective for subgroups 34 of those affected with low back pain and completed a secondary analysis to investigate 35 this. In this secondary analysis of data from a randomized clinical trial, they tested whether 36 five indicators of a potential increased need for treatment might act as effect modifiers for 37 manipulative therapy in the treatment of spinal pain in children. Investigators hypothesized 38 39 that the most severely affected children would benefit more from manipulative therapy. To explore potential effect modification, various types of regression models were used 40 depending on the type of outcome, including interaction tests. Authors found that children 41 with long duration of spinal pain or co-occurring musculoskeletal pain prior to inclusion 42

CPG 120 Revision 16 – S Spinal Manipulative Therapy for Treatment of Children and Infants **Revised – March 20, 2025** To CQT for review 02/10/2025 CQT reviewed 02/10/2025 To QIC for review and approval 03/04/2025 QIC reviewed and approval 03/04/2025 To QOC for review and approval 03/20/2025 OOC reviewed and approval 03/20/2025 Page 3 of 16

as well as low quality of life at baseline tended to benefit from manipulative therapy over 1 non-manipulative therapy, whereas the opposite was seen for children reporting high 2 intensity of pain. However, most results were statistically insignificant. Authors concluded 3 that this secondary analysis indicated that children more effected by certain baseline 4 characteristics, but not pain intensity, have a greater chance to benefit from treatment that 5 include manipulative therapy. However, these analyses were both secondary and 6 underpowered, and therefore merely exploratory. The results underline the need for a 7 careful choice of inclusion criteria in future investigations of manipulative therapy in 8 children. 9

10

11 Driehuis et al. (2019) conducted a systematic review of the evidence for effectiveness and harms of specific SMT techniques for infants, children, and adolescents. Of the 1,236 12 identified studies, 26 studies were eligible. Infants and children/adolescents were treated 13 for various (non-)musculoskeletal indications, hypothesized to be related to spinal joint 14 dysfunction. Studies examining the same population, indication and treatment comparison 15 were scarce. Due to very low-quality evidence, it is uncertain whether gentle, low-velocity 16 mobilizations reduce complaints in infants with colic or torticollis, and whether high-17 velocity, low-amplitude manipulations reduce complaints in children/adolescents with 18 autism, asthma, nocturnal enuresis, headache, or idiopathic scoliosis. Five case reports 19 20 described severe harms after HVLA manipulations in four infants and one child. Authors found the evidence was of very low-quality that prevented drawing any conclusions about 21 the effectiveness of specific SMT techniques in infants, children, and adolescents. 22

23

Parnell Prevost et al. (2019) evaluated the use of manual therapy (MT) for clinical 24 conditions in the pediatric population, assessed the methodological quality of the studies 25 found, and synthesized findings based on health condition within a systematic review. They 26 also assessed the reporting of adverse events within the included studies and compared the 27 conclusions to those of the UK Update report. Six databases were searched using the 28 following inclusion criteria: children under the age of 18 years old; treatment using manual 29 therapy; any type of healthcare profession; published between 2001 and March 31, 2018; 30 and English. Case reports were excluded. Of the 3,563 articles identified, 165 full articles 31 were screened, and 50 studies met the inclusion criteria. Twenty-six articles were included 32 33 in prior reviews with 24 new studies identified. Eighteen studies were judged to be of high quality. Conditions evaluated were: attention deficit hyperactivity disorder (ADHD), 34 autism, asthma, cerebral palsy, clubfoot, constipation, cranial asymmetry, cuboid 35 syndrome, headache, infantile colic, low back pain, obstructive apnea, otitis media, 36 pediatric dysfunctional voiding, pediatric nocturnal enuresis, postural asymmetry, preterm 37 infants, pulled elbow, suboptimal infant breastfeeding, scoliosis, suboptimal infant 38 39 breastfeeding, temporomandibular dysfunction, torticollis, and upper cervical dysfunction. Musculoskeletal conditions, including low back pain and headache, were evaluated in 40 seven studies. Twenty studies reported adverse events, which were transient and mild to 41 moderate in severity. Authors concluded that moderate-positive overall assessment was 42

found for 3 conditions: low back pain and chiropractic manipulation, pulled elbow (MT), and premature infants (osteopathic manipulation and craniosacral techniques). Inconclusive unfavorable outcomes were found for 2 conditions: scoliosis (OMT) and torticollis (MT). All other condition's overall assessments were either inconclusive favorable or unclear for all manual therapies including SMT. Adverse events were uncommonly reported. More robust clinical trials in this area of healthcare are needed.

7

Lynge et al. (2021) investigated the effectiveness of chiropractic spinal manipulation 8 versus sham manipulation in children aged 7-14 with recurrent headaches. A total of 199 9 children aged 7 to 14 years, with at least one episode of headache per week for the previous 10 11 6 months and at least one musculoskeletal dysfunction were identified. All participants received standard oral and written advice to reduce headaches. In addition, children in the 12 active treatment group received chiropractic spinal manipulation and children in the control 13 group received sham manipulation for a period of 4 months. Number and frequency of 14 treatments were based on the chiropractor's individual evaluation in the active treatment 15 group; the children in the control group received approximately eight visits during the 16 treatment period. 'Number of days with headache', 'pain intensity' and 'medication' were 17 reported weekly by text messages, and global perceived effect by text message after 4 18 months. 'Number of days with headache' and 'pain intensity' were chosen as equally 19 20 important outcomes of highest priority, followed by global perceived effect and medication. Results demonstrated that chiropractic spinal manipulation resulted in 21 significantly fewer days with headaches and better global perceived effect compared with 22 a sham manipulation procedure. There was no difference between groups for pain intensity 23 during headache episodes. Due to methodological shortcomings, no conclusions could be 24 drawn about medication use. Authors concluded that chiropractic spinal manipulation 25 resulted in fewer headaches and higher global perceived effect, with only minor side 26 effects. It did not lower the intensity of the headaches. Since the treatment is easily 27 applicable, of low cost and minor side effects, chiropractic spinal manipulation might be 28 considered as a valuable treatment option for children with recurrent headaches. 29

30

Dice et al. (2021) sought to identify the following among physical therapists holding 31 advanced credentials in pediatrics, neurodevelopmental treatment, or manual therapy: (1) 32 33 consensus regarding effective techniques in the preadolescent population, (2) differences in opinion, and (3) perceived decision-making barriers and factors regarding use of manual 34 therapy techniques. Credentialed physical therapists in the United States were recruited for 35 a 3-round Delphi investigation. An electronic survey in Round 1 identified musculoskeletal 36 and neurological impairments and the manual techniques considered effective to treat such 37 conditions, in addition to factors and barriers. Responses were used to create the second 38 39 round, during which a 4-point Likert scale was used to score each survey item. A third round of scoring established consensus. Descriptive statistics and composite scores were 40 calculated for each manual technique by impairment. Consensus was determined for 41 several concepts. First, neuromuscular techniques were considered effective across all 42

Page 5 of 16

impairments, and joint mobilizations (grades I-IV) were believed to be effective to treat 1 joint and muscle and myofascial impairments. Second, visceral manipulation and 2 craniosacral therapy were considered ineffective in treating most impairments. There was 3 lack of consensus and clear differences of opinion regarding the use of grade V 4 mobilizations (SMT) and dry needling. Significant barriers to use of manual therapy were 5 lack of knowledge, lack of evidence, and fear of litigation and harming patients. Authors 6 summarized by stating that this study is an initial step for developing manual therapy 7 guidelines, research, and educational opportunities regarding manual therapy in pediatric 8 physical therapy. 9

10

11 Milne et al. (2022) sought to identify and map the available evidence regarding effectiveness and harms of spinal manipulation and mobilization for infants, children, and 12 adolescents with a broad range of conditions; and identify and synthesize policies, 13 regulations, position statements and practice guidelines informing their clinical use. 14 Infants, children, and adolescents (birth to < 18 years) with any childhood 15 disorder/condition who received an intervention of spinal manipulation and mobilization 16 were included as participants. Eighty-seven articles were included. Methodological quality 17 of articles varied. Spinal manipulation and mobilization may be utilized clinically to 18 manage pediatric populations with adolescent idiopathic scoliosis, asthma, attention deficit 19 20 hyperactivity disorder, autism spectrum disorder, back/neck pain, breastfeeding difficulties, cerebral palsy, dysfunctional voiding, excessive crying, headaches, infantile 21 colic, kinetic imbalances due to suboccipital strain, nocturnal enuresis, otitis media, 22 torticollis and plagiocephaly. This descriptive synthesis revealed: no evidence to explicitly 23 support the effectiveness of spinal manipulation or mobilization for any condition in 24 pediatric populations. Mild transient symptoms were commonly described in randomized 25 controlled trials and on occasion, moderate-to-severe adverse events were reported in 26 systematic reviews of randomized controlled trials and other lower quality studies. There 27 was strong to very strong evidence for 'no significant effect' of spinal manipulation for 28 managing asthma (pulmonary function), headache and nocturnal enuresis, and 29 inconclusive or insufficient evidence for all other conditions explored. There is insufficient 30 evidence to draw conclusions regarding spinal mobilization to treat pediatric populations 31 with any condition. Authors concluded that their descriptive synthesis of the collective 32 33 findings does not provide support for spinal manipulation or mobilization in pediatric populations for any condition. Increased reporting of adverse events is required to 34 determine true risks. Randomized controlled trials examining effectiveness of spinal 35 manipulation and mobilization in pediatric populations are warranted. 36

37

Franke et al. (2022) reviewed the literature to determine the effectiveness of osteopathic manipulative treatment (OMT) for all pediatric complaints. Forty-seven RCTs examining pediatric conditions were reviewed. Twenty-three studies reported significant favorable outcomes for OMT relative to the control intervention, and 14 additional studies reported non-significant outcomes, which suggested potential favorable effects of OMT. Authors

Page 6 of 16

1 concluded that although a number of studies indicated positive results with use of OMT,

2 few pediatric conditions have been investigated in more than one study, which results in

no high-quality evidence for any condition. Additional research may change estimates of
 effect, and larger, high-quality RCTs focusing on a smaller range of conditions are

- 5 recommended.
- 6

Gross et al. (2024) developed evidence-based practice position statements directing 7 physiotherapists clinical reasoning for the safe and effective use of spinal manipulation and 8 mobilization for pediatric populations (<18 years) with varied musculoskeletal or non-9 musculoskeletal conditions. A three-stage guideline process using validated methodology 10 was completed: 1. Literature review stage (one scoping review, two reviews exploring 11 psychometric properties); 2. Delphi stage (one 3-Round expert Delphi survey); and 3. 12 Refinement stage (evidence-to-decision summative analysis, position statement 13 development, evidence gap map analyses, and multilayer review processes). Evidence-14 based practice position statements were developed to guide the appropriate use of spinal 15 manipulation and mobilization for pediatric populations. All were predicated on clinicians 16 using biopsychosocial clinical reasoning to determine when the intervention was 17 appropriate. 18

- 19 1. It is not recommended to perform:
 - a. Spinal manipulation and mobilization on infants.
 - b. Cervical and lumbar spine manipulation on children.
- c. Spinal manipulation and mobilization on infants, children, and adolescents
 for non-musculoskeletal pediatric conditions including asthma, attention
 deficit hyperactivity disorder, autism spectrum disorder, breastfeeding
 difficulties, cerebral palsy, infantile colic, nocturnal enuresis, and otitis
 media.
- It may be appropriate to treat musculoskeletal conditions including spinal mobility
 impairments associated with neck-back pain and neck pain with headache utilizing:
 - a. Spinal mobilization and manipulation on adolescents;
 - b. Spinal mobilization on children; or
 - c. Thoracic manipulation on children for neck-back pain only.
- 32 3. No high certainty evidence to recommend these interventions was available. 33 Reports of mild to severe harms exist; however, risk rates could not be determined.
- 33 34

29

30

31

20

21

Dice et al. (2024) established international consensus regarding the use of spinal 35 manipulation and mobilization among infants, children, and adolescents among expert 36 international physiotherapists. Twenty-six international expert physiotherapists in manual 37 therapy and pediatrics voluntarily participated in a 3-Round Delphi survey to reach a 38 39 consensus via direct electronic mail solicitation using Qualtrics[®]. Consensus was defined a-priori as >75% agreement on all items with the same ranking of agreement or 40 disagreement. Round 1 identified impairments and conditions where spinal mobilization 41 and manipulation might be utilized. In Rounds 2 and 3, panelists agreed or disagreed using 42

Page 7 of 16

a 4-point Likert scale. Eleven physiotherapists from seven countries representing five 1 continents completed all three Delphi rounds. Consensus regarding spinal mobilization or 2 manipulation included: Manipulation is not recommended: (1) for infants across all 3 conditions, impairments, and spinal levels; and (2) for children and adolescents across most 4 conditions and spinal levels. Manipulation may be recommended for adolescents to treat 5 spinal region-specific joint hypomobility (thoracic, lumbar), and pain (thoracic). 6 Mobilization may be recommended for children and adolescents with hypomobility, joint 7 pain, muscle/myofascial pain, or stiffness at all spinal levels. Consensus revealed spinal 8 manipulation should not be performed on infants regardless of condition, impairment, or 9 spinal level. Additionally, the panel agreed that manipulation may be recommended only 10 for adolescents to treat joint pain and joint hypomobility (limited to thoracic and/or lumbar 11 levels). Spinal mobilization may be recommended for joint hypomobility, joint pain, 12 muscle/myofascial pain, and muscle/myofascial stiffness at all spinal levels among 13 children and adolescents. 14

15

Yu et al. (2024) evaluated benefits and harms of rehabilitation interventions for non-16 specific low back pain (LBP) or thoracic spine pain in the pediatric population. Ten 17 quantitative studies (i.e., 8 RCTs, 2 non-randomized clinical trials) and one qualitative 18 study were included. With very low to moderate certainty evidence, in adolescents with 19 20 LBP, spinal manipulation (1-2 sessions/week over 12 weeks, 1 RCT) plus exercise may be associated with a greater likelihood of experiencing clinically important pain reduction 21 versus exercise alone; and group-based exercise over 8 weeks (2 RCTs and 1 non-22 randomized trial) may reduce pain intensity. The qualitative study found information 23 provided via education/advice and compliance of treatment were related to effective 24 treatment. No economic studies or studies examining thoracic spine pain were identified. 25 Authors concluded that spinal manipulation and group-based exercise may be beneficial in 26 reducing LBP intensity in adolescents. Education should be provided as part of a care 27 program. The overall evidence is sparse. Methodologically rigorous studies are needed. 28

29

Keating et al. (2024) conducted a formal consensus process and best evidence synthesis to 30 build upon existing recommendations on best practices for chiropractic management of 31 children. Authors did a synthesis of results of a literature search to inform the development 32 33 of recommendations from a multidisciplinary steering committee, including experts in pediatrics, followed by a formal Delphi panel consensus process. All 60 panelists 34 completed the process and reached at least 80% consensus on all recommendations after 35 three Delphi rounds. Recommendations for best practices for chiropractic care for children 36 addressed these aspects of the clinical encounter: patient communication, including 37 informed consent; appropriate clinical history, including health habits; appropriate physical 38 39 examination procedures; red flags/contraindications to chiropractic care and/or spinal manipulation; aspects of chiropractic management of pediatric patients, including infants; 40 modifications of spinal manipulation and other manual procedures for pediatric patients; 41 appropriate referral and co-management; and appropriate health promotion and disease 42

Page 8 of 16

prevention practices. Specific to chiropractic management of pediatric patients, authors 1 report there are four basic chiropractic management approaches to the care of a child: (1) 2 sole management by a chiropractor, (2) independent concurrent care by a chiropractor and 3 other provider(s), (3) co-management with other appropriate health care providers, and (4) 4 referral to another registered/licensed or certified health care provider/specialist. They 5 recommend that chiropractors follow the principles of evidence-based practice, which are 6 to make clinical judgments based on the best available evidence combined with clinical 7 experience and the patient's preferences. Children may present to chiropractic practices 8 with various conditions and developmental concerns not directly related to the 9 neuromusculoskeletal system. They note there is a paucity of high-level research evidence 10 11 for the effectiveness of spinal manipulation and/or chiropractic care for such conditions. They also note there are special considerations for use of spinal manipulation and other 12 manual procedures with children and practitioners must modify manipulative and/or 13 mobilization and soft tissue techniques as appropriate for the child's age and 14 developmental stage. 15

16

Piqueras-Toharias et al. (2024) sought to determine whether high-velocity low-amplitude 17 (HVLA) spinal manipulation is more effective than other treatments for children with 18 idiopathic scoliosis (IS) in a systematic review. Five studies were selected for review. The 19 20 results indicated moderate improvements in pain and the Cobb angle and limited improvements in quality of life. Authors concluded that HVLA spinal manipulation does 21 not seem to have significant effects on reducing spinal deformity in IS patients, nor does it 22 significantly impact quality of life. However, this therapy may have significant effects on 23 24 reducing pain in these patients.

2526 SAFETY

27 The potential risk of a major complication due to spinal manipulation is rare (Hurwitz et al., 1996; Todd et al., 2014). These rare, serious adverse events attributed to SMT in 28 children included quadriplegia and death. Evidence of complications associated with SMT 29 in children comes primarily from case reports and case series. While serious adverse events 30 may be associated with pediatric spinal manipulation, neither causation nor incidence rates 31 can be inferred from observational data (Vohra et al., 2007). No serious complications from 32 33 SMT have been reported from any of the published randomized clinical trials or observational studies involving SMT in children. Several minor transient adverse reactions 34 have been reported. Based on a review of the literature, both the possible harms and 35 possible benefits of SMT in children appear to be minimal. 36

37

Cervical mobilization and manipulation have been suspected of creating a cervical artery dissection (CAD) as an adverse event. However, these assumptions are based on case studies which are unable to establish direct causality. Chaibi and Bjørn Russel (2019) conducted a literature review to provide clinicians with an updated step-by-step risk– benefit assessment strategy tool to (a) facilitate clinicians understanding of CAD,

Page 9 of 16

(b) appraise the risk and applicability of cervical manual-therapy, and (c) provide clinicians 1 with adequate tools to better detect and exclude CAD in clinical settings. Cervical artery 2 dissection refers to a tear in the internal carotid or the vertebral artery that results in an 3 intramural hematoma and/or aneurysmal dilatation. Although cervical artery dissection is 4 thought to occur spontaneously and is rare, physical trauma to the neck, especially 5 hyperextension and rotation, has been reported as a trigger. Headache and/or neck pain is 6 the most common initial symptom of cervical artery dissection. Other symptoms include 7 Horner's syndrome and lower cranial nerve palsy. Both headache and/or neck pain are 8 common symptoms and leading causes of disability. Because manual-therapy interventions 9 can alleviate headache and/or neck pain, many patients seek manual therapists, such as 10 11 chiropractors and physiotherapists to help them manage symptoms. There is debate as to whether CAD symptoms lead the patient to seek cervical manual-therapy or whether the 12 cervical manual therapy provoked CAD along with the non-CAD presenting complaints. 13 Thus, practitioners need to be diligent with subjective and objective evaluations of patients 14 to understand the risk for CAD and whether to address its potential existence. 15

16

Corso et al. (2020) conducted a rapid review of the safety of SMT in children (< 10 years). 17 Their aim was to 1) describe adverse events; 2) report the incidence of adverse events; and 18 3) determine whether SMT increases the risk of adverse events compared to other 19 20 interventions. Authors found that most adverse events are mild (e.g., increased crying, soreness). One case report describes a severe adverse event (rib fracture in a 21-day-old) 21 and another an indirect harm in a 4-month-old. The incidence of mild adverse events ranges 22 from 0.3% to 22.22%. Whether SMT increases the risk of adverse events in children is 23 unknown. Authors concluded that the risk of moderate and severe adverse events is 24 unknown in children treated with SMT. It is unclear whether SMT increases the risk of 25 adverse events in children < 10 years. Vos et al. (2021) carried out a 3- year survey on 26 pediatric use of complementary and alternative medicine (CAM) in the Netherlands 27 Pediatricians were asked to register cases of adverse events associated with pediatric CAM 28 usage. In 3 years, 32 unique adverse events were registered. Twenty-two of these adverse 29 events were indirect and not related to the specific CAM therapy but due to delaying, 30 changing, or stopping of regular treatment, a deficient or very restrictive diet or an incorrect 31 diagnosis by a CAM therapist. These events were associated with many different CAM 32 33 therapies. Nine events were deemed direct adverse events like bodily harm or toxicity and one-third of them occurred in infants. Only supplements, manual therapies, and (Chinese) 34 herbs were involved in these nine events. For SMT, 2 adverse events occurred: torticollis 35 and transient nerve palsy. Relatively few cases of adverse events associated with pediatric 36 CAM usage were found, mostly due to delaying or stopping conventional treatment. 37 Nevertheless, parents, pediatricians and CAM providers should be vigilant for both direct 38 39 and indirect adverse events in children using CAM, especially in infants.

CPG 120 Revision 16 – S Spinal Manipulative Therapy for Treatment of Children and Infants Revised – March 20, 2025 To CQT for review 02/10/2025 CQT reviewed 02/10/2025 To QIC for review and approval 03/04/2025 QIC reviewed and approval 03/04/2025 To QOC for review and approval 03/20/2025 OOC reviewed and approval 03/20/2025 Page 10 of 16

1 Clinicians need to provide pediatric patients and their parents or guardians with

- 2 information regarding benefits, harms, and alternatives relevant to making an informed3 treatment decision.
- 4

5 **PRACTITIONER SCOPE AND TRAINING**

Practitioners should practice only in the areas in which they are competent based on their
education, training, and experience. Levels of education, experience, and proficiency may
vary among individual practitioners. It is ethically and legally incumbent on a practitioner
to determine where they have the knowledge and skills necessary to perform such services.

It is best practice for the practitioner to appropriately render services to a patient only if they are trained, equally skilled, and adequately competent to deliver a service compared to others trained to perform the same procedure. If the service would be most competently delivered by another health care practitioner who has more skill and expert training, it would be best practice to refer the patient to the more expert practitioner.

16

Best practice can be defined as a clinical, scientific, or professional technique, method, or process that is typically evidence-based and consensus driven and is recognized by a majority of professionals in a particular field as more effective at delivering a particular outcome than any other practice (Joint Commission International Accreditation Standards for Hospitals, 2020).

22

Depending on the practitioner's scope of practice, training, and experience, a member's condition and/or symptoms during examination or the course of treatment may indicate the need for referral to another practitioner or even emergency care. In such cases it is prudent for the practitioner to refer the member for appropriate co-management (e.g., to their primary care physician) or if immediate emergency care is warranted, to contact 911 as appropriate. See the *Managing Medical Emergencies (CPG 159 – S)* clinical practice guideline for information.

30

31 *References*

Balon J, Aker PD, Crowther ER, Danielson C, et al. A comparison of active and simulated
 chiropractic manipulation as adjunctive treatment for childhood asthma. N Engl J Med
 1998; 339:1013-1020

35

Bronfort G, Evans RL, Kubic P, Filkin P. Chronic pediatric asthma and chiropractic spinal manipulation: A prospective clinical series and randomized clinical pilot study. J Manipulative Physiol Ther 2001; 24:369-377

39

Bronfort G, Haas M, Evans R, Leininger B, Triano J. Effectiveness of manual therapies:
 the UK evidence report. Chiropr Osteopat. 2010;18:3

1 2 3 4	Chaibi A, Russell MB. A risk-benefit assessment strategy to exclude cervical artery dissection in spinal manual-therapy: a comprehensive review. Ann Med. 2019;51(2):118-127. doi:10.1080/07853890.2019.1590627
5 6 7 8 9	Clar C, Tsertsvadze A, Court R, Hundt GL, Clarke A, Sutcliffe P. Clinical effectiveness of manual therapy for the management of musculoskeletal and non-musculoskeletal conditions: systematic review and update of UK evidence report. Chiropr Man Therap. 2014 Mar 28;22(1):12
10 11 12	Cohen MH, Kemper KJ. Complementary therapies in pediatrics: A legal perspective. Pediatrics 2005; 115:774-780
13 14 15	Corso M, Cancelliere C, Mior S, Taylor-Vaisey A, Côté P. The safety of spinal manipulative therapy in children under 10 years: a rapid review. Chiropr Man Therap. 2020 Feb 25;28(1):12
10 17 18 19 20	Dice JL, Dendy D, Sizer PS, Cook CE, Feuling S, Brismée JM. Manual Therapy in Preadolescent Children: A Delphi Investigation of Physical Therapists in the United States. Phys Ther. 2021;101(4):pzab027
21 22 23 24	Dice JL, Brismée JM, Froment FP, et al. Spinal manipulation and mobilisation among infants, children, and adolescents: an international Delphi survey of expert physiotherapists. J Man Manip Ther. 2024;32(3):284-294
25 26 27 28 29	Dissing KB, Hartvigsen J, Wedderkopp N, Hestbæk L. Conservative care with or without manipulative therapy in the management of back and/or neck pain in Danish children aged 9-15: a Randomized controlled trial nested in a school-based cohort. BMJ Open. 2018;8(9):e021358. Published 2018 Sep 10
30 31 32 33 34	Dissing KB, Vach W, Hartvigsen J, Wedderkopp N, Hestbæk L. Potential treatment effect modifiers for manipulative therapy for children complaining of spinal pain. Secondary analyses of a Randomized controlled trial. Chiropr Man Therap. 2019;27:59. Published 2019 Dec 10
34 35 36 37	Dobson D, Lucassen PL, Miller JJ, Vlieger AM, Prescott P, Lewith G. Manipulative therapies for infantile colic. Cochrane Database Syst Rev. 2012 Dec 12;12:CD004796
38 39 40 41	Driehuis F, Hoogeboom TJ, Nijhuis-van der Sanden MWG, de Bie RA, Staal JB. Spinal manual therapy in infants, children and adolescents: A systematic review and meta- analysis on treatment indication, technique and outcomes. PloS One. 2019 Jun 25;14(6):e0218940

Page 12 of 16

1 2 3 4	Evans R, Haas M, Schulz C, Leininger B, Hanson L, Bronfort G. Spinal manipulation and exercise for low back pain in adolescents: a randomized trial. Pain. 2018;159(7):1297-1307
5 6 7	Franke H, Franke JD, Fryer G. Effectiveness of osteopathic manipulative treatment for pediatric conditions: A systematic review. J Bodyw Mov Ther. 2022;31:113-133. doi:10.1016/j.jbmt.2022.03.013
8	
9	Ferrance, RJ, Miller, J. Chiropractic diagnosis and management of non-musculoskeletal
10	conditions in children and adolescents. Chiropractic & Osteopathy. 2010;18:14
11	
12	Glazener CM, Evans JH, Cheuk DK. Complementary and miscellaneous interventions for
13	nocturnal enuresis in children. Cochrane Database Syst Rev 2005; April
14	8(2):CD005230
15	
16	Gleberzon BJ, Arts J, Mei A, McManus EL. The use of spinal manipulative therapy for
1/	pediatric nearin conditions: a systematic review of the interature. J Can Uniropr Assoc. 2012 Jum 56(2):128–41
10	2012 Juli, 50(2). 128-41
19 20	Gotlib A Rupert R Assessing the evidence for the use of chiropractic manipulation in
20	nediatric health conditions – a systematic review Pediatr Child Health 2005: 10:157-
21	161
22	101
24	Gotlib A. Rupert R. Chiropractic manipulation in pediatric health conditions – an updated
25	systematic review. Chiropr Osteopat 2008; 16:11 (12 September 2008)
26	
27	Gross AR, Olson KA, Pool J, et al. Spinal manipulation and mobilisation in paediatrics -
28	an international evidence-based position statement for physiotherapists. J Man Manip
29	Ther. 2024;32(3):211-233.
30	
31	Guiney PA, Chou R, Vianna A, Lovenheim J. Effects of osteopathic manipulative
32	treatment on pediatric patients with asthma: A randomized controlled trial. J Am
33	Osteopath Assoc 2005; 105:7-12
34	
35	Hawk C, Khorsan R, Lisi AJ, Ferrance RJ, Evans MW. Chiropractic care for
36	nonmusculoskeletal conditions: A systematic review with implications for whole
37	systems research. J Altern Complement Med 2007; 13:491-512
38	Hands C. Calandida MI, Vallana C. Handitt EC. D. (D. C.
39 40	Hawk U, Schneider MJ, Vallone S, Hewitt EG. Best Practices for Uniropractic Care of Children A Concensus Undete. I Monimulative Divisiol Theor. 2016;20(2):159-169
40	Cindien. A Consensus Opuale. J Manipulative Physiol Ther. 2010;39(3):138-108

Page 13 of 16

1 2 3 4	Hayden JA, Mior SA, Verhoef MJ. Evaluation of chiropractic management of pediatric patients with low back pain: a prospective cohort study. J Manipulative Physiol Ther 2003; 26:1-8
5 6 7	Hondras MA, Linde K, Jones AP. Manual therapy for asthma. Cochrane Database Syst Rev 2005; April 18(2):CD001002
8 9 10	Hughes S, Bolton J. Is chiropractic an effective treatment in infantile colic? Arch Dis Child 2002; 86:517-522
10 11 12 13	Humphreys, BK. Possible adverse events in children treated by manual therapy: a review. Chiropractic & Osteopathy 2010;18:12
14 15 16	Hurwitz, EL, Aker, PD, Adams, AH, Meeker, WC, Shekelle, PG. Manipulation and Mobilization of the Cervical Spine: A Systematic Review of the Literature. Spine. 1 August 1996;21(15): 1746-1759
17 18 19 20 21	Husereau D, Clifford T, Aker P, Leduc D, Medninkai S. Spinal manipulation for infantile colic. Ottawa: Canadian Coordinating Office for Health Technology Assessment 2003; Technology report no.42
21 22 23 24 25	Keating G, Hawk C, Amorin-Woods L, et al. Clinical Practice Guideline for Best Practice Management of Pediatric Patients by Chiropractors: Results of a Delphi Consensus Process. J Integr Complement Med. 2024;30(3):216-232
23 26 27 28 29	Khorshid KA, Sweat RW, Zemba DA, Zemba BN. Clinical efficacy of upper cervical versus full spine chiropractic care on children with autism: a randomized clinical trial. J Vert Sublux Res 2006; March 9, online:1-7
30 31 32	Lantz CA, Chen J. Effect of chiropractic intervention on small scoliotic curves in younger subjects: A time-series cohort design. J Manipulative Physiol Ther 2001; 24:385-393
33 34 35	Leach RA. Differential compliance instrument in the treatment of infantile colic: A report of two cases. J Manipulative Physiol Ther 2002; 25:58-62
36 37 38	Lee AC, Li DH, Kemper KJ. Chiropractic care for children. Arch Pediatr Adolesc Med 2000; 154:401-407
39 40 41 42	Lynge S, Dissing KB, Vach W, Christensen HW, Hestbaek L. Effectiveness of chiropractic manipulation versus sham manipulation for recurrent headaches in children aged 7-14 years – a Randomized clinical trial. Chiropr Man Therap. 2021;29(1):1. Published 2021 Jan 7

Page 14 of 16

1 2 3	Miller JE, Benfield K. Adverse effects of spinal manipulative therapy in children younger than 3 years: A retrospective study in a chiropractic teaching clinic. J Manipulative Physiol Ther 2008; 31:419-423
5 6 7 8	Mills MV, Henley CE, Barnes LL, et al. The use of osteopathic manipulative treatment as adjuvant therapy in children with recurrent acute otitis media. Arch Pediatr Adolesc Med 2003; 157:861-866
9 10 11 12	Milne N, Longeri L, Patel A, et al. Spinal manipulation and mobilisation in the treatment of infants, children, and adolescents: a systematic scoping review. BMC Pediatr. 2022;22(1):721. Published 2022 Dec 19. doi:10.1186/s12887-022-03781-6
13 14 15	Olafsdottir E, Forshei S, Fluge G, Markestad T. Randomised controlled trial of infantile colic treated with chiropractic spinal manipulation. Arch Dis Child 2001; 84:138-141
16 17 18	Olson KA, Clewley D, Milne N, et al. Spinal manipulation and mobilisation for paediatric conditions: time to stop the madness. J Man Manip Ther. 2024;32(3):207-210
19 20 21	Parnell Prevost C, Gleberzon B, Carleo B, Anderson K, Cark M, Pohlman KA. Manual therapy for the pediatric population: a systematic review. BMC Complement Altern Med. 2019 Mar 13;19(1):60
22 23 24 25 26	Piqueras-Toharias M, Ibáñez-Vera AJ, Peinado-Rubia AB, Rodríguez-Almagro D, Lomas- Vega R, Sedeño-Vidal A. Effects of High-Velocity Spinal Manipulation on Quality of Life, Pain and Spinal Curvature in Children with Idiopathic Scoliosis: A Systematic Review. Children (Basel). 2024;11(10):1167. Published 2024 Sep 26
27 28 29 20	Pohlman KA, Holton-Brown MS. Otitis media and spinal manipulative therapy: a literature review. J Chiropr Med. 2012 Sep;11(3):160-9
30 31 32 33 34	Rowe DE, Feise RJ, Crowther ER, Grod JP, Menke JM, Goldsmith CH, Stoline MR, Souza TA, Kambach B. Chiropractic manipulation in adolescent idiopathic scoliosis: a pilot study. Chiropr Osteopat 2006; 14:15
35 36 37 38	Sawyer CE, Evans RL, Boline PD, et al. A feasibility study of chiropractic spinal manipulation versus sham spinal manipulation for chronic otitis media with effusion in children. J Manipulative Physiol Ther 1999; 22:292-298
39 40	Shafrir Y, Kaufman BA. Quadriplegia after chiropractic manipulation in an infant with congenital torticollis caused by a spinal cord astrocytoma. J Pediatr 1992; 120:266-268

Page 15 of 16

1	Straub WF, Spino MP, Alattar MM, Pfleger B, Downes JW, Belizaire MA, Heinonen OJ,
2	Vasankari T. The effect of chiropractic care on jet lag of Finnish junior elite athletes. J
3	Manipulative Physiol Ther 2001; 24:191-198
4	
5	Terrett AGJ, Kleynhans AM. Complications from manipulation of the low back.
6	Chiropractic J Aust. 1992;22:129
7	
8	Todd AJ, Carroll MT, Robinson A, Mitchell EK. Adverse Events Due to Chiropractic and
9	Other Manual Therapies for Infants and Children: A Review of the Literature. J
10	Manipulative Physiol Ther. 2014 Oct 30. pii: S0161-4754
11	
12	Vaughn DW, Kenyon LK, Sobeck CM, Smith RE. Spinal manual therapy interventions for
13	pediatric patients: a systematic review. J Man Manip Ther. 2012 Aug;20(3):153-9
14	
15	Vohra S, Johnston BC, Cramer K, Humphries K. Adverse events associated with pediatric
16	spinal manipulation: A systematic review. Pediatrics 2007; 119:e275-e283
17	
18	Vos B, Rake JP, Vlieger A. Adverse events associated with pediatric complementary and
19	alternative medicine in the Netherlands: a national surveillance study. Eur J Pediatr.
20	2021;180(7):2165-2171
21	
22	Yu H, Southerst D, Wong JJ, et al. Rehabilitation of back pain in the pediatric population:
23	a mixed studies systematic review. Chiropr Man Therap. 2024;32(1):14. Published
24	2024 May 8