

1	Clinical Practice Guideline:	Tobacco Cessation Counseling
2		
3	Date of Implementation:	April 19, 2012
4		
5	Effective Date:	April 16, 2026
6		
7	Product:	Specialty

8 **Table of Contents**

9 GUIDELINES 1

10 INTRODUCTION..... 1

11 SCREENING RECOMMENDATIONS 3

12 EFFECTIVE INTERVENTIONS 5

13 Counseling..... 7

14 Evidence-Based Behavior Modification Techniques..... 7

15 Medications 8

16 PRACTITIONER SCOPE AND TRAINING 9

17 PRACTITIONER RESOURCES 9

18 MEMBER RESOURCES.....10

19 *References*10

20

21 **GUIDELINES**

22 American Specialty Health – Specialty (ASH) clinical committees determined that in the

23 context of best practices and for the population of all patients, the evaluation of

24 tobacco/nicotine use is necessary. In this same context, a brief intervention for the

25 population of tobacco users is recommended. An example of a brief intervention would be

26 what is recommended by the United States Preventive Services Task Force (USPSTF) for

27 tobacco users. Lastly, ASH clinical committees concluded in the same context of best

28 practices for the population of tobacco users, a direct intervention or referral, depending

29 upon the expertise and scope of the practitioner, for appropriate tobacco/nicotine cessation

30 intervention is necessary.

31

32 **INTRODUCTION**

33 An office visit with a health care practitioner can provide an opportunity to talk with

34 patients about their tobacco/nicotine use. Given the health effects associated with chronic

35 tobacco use, the office visit provides a "teachable moment" during which a qualified

36 healthcare professional can relate current health problems to tobacco use, provide brief

37 counseling, or set an appropriate referral for patients who use tobacco products.

1 Tobacco contains nicotine, an addictive substance. Addiction to nicotine can happen after
2 the first exposure so prevention is a key intervention to reducing smoking in the population.

3
4 Tobacco use is the leading preventable cause of death in the U.S. This is more than the
5 combined total from AIDS, alcohol, cocaine, heroin, homicide, suicide, motor vehicle
6 crashes, and fires. The association between tobacco use and premature death is one of the
7 best documented in the epidemiological literature, beginning with Doll’s study of over
8 40,000 male physicians in 1951; this study then continued, following participants for 50
9 years. These studies showed that cigarette smokers had twice the death rate ratio as
10 nonsmokers (42% to 24%) for premature death (at ages 35-69). Cigarette smoking was
11 found to be highly correlated with all causes of death, as was number of cigarettes smoked,
12 which demonstrates a strong dose-response effect.

13
14 Smoking is known to cause cancer, heart disease, stroke, lung diseases such as COPD, and
15 diabetes. Tobacco use in any form can also lead to health issues including various cancers,
16 pregnancy complications, lung diseases, gum disease, and vision problems. Some studies
17 suggest that tobacco use may be a risk factor for low back pain and may contribute to
18 poorer outcomes in people with musculoskeletal back pain, including outcomes of
19 rehabilitation care. Secondhand smoke exposure contributes to an estimated 40,000 deaths
20 among non-smoking adults and 400 deaths of infants every year. Besides the health impact,
21 smoking also increases health care utilization, health care costs, and absenteeism from
22 work.

23
24 Electronic cigarettes (vapes) usually contain nicotine and are an emerging issue in tobacco
25 use and cessation. According to the Centers for Disease Control, in 2022, more than 2.55
26 million U.S. middle and high school students had used e-cigarettes in the past 30 days. This
27 includes 14.1% of high school students and 3.3% of middle school students. The Food and
28 Drug Administration reported that e-cigarette use, from 2017 to 2018, increased 78%
29 among high school students and 48% among middle school students. Additionally, 4.5%
30 of adults aged 18 or older are e-cigarette users, with highest use among those between 18-
31 24 years of age. Adults who are between the ages of 18-44 are more likely to smoke both
32 cigarettes and vape in comparison with adults 45 years of age or older (Kramarow &
33 Elgaddal, 2023).

34
35 E-cigarettes are not currently approved by the FDA as a quit-smoking aid. The U.S.
36 Preventive Services Task Force, has concluded that evidence is insufficient to “assess the
37 balance of benefits and harms of e-cigarettes for tobacco cessation in adults, including
38 pregnant persons.” The USPSTF (2021) recommends that clinicians direct patients who
39 use tobacco to other tobacco cessation interventions with proven effectiveness and
40 established safety. Based on evolving evidence, ASH does not currently support e-
41 cigarettes as a viable method of tobacco cessation or nicotine replacement.

1 **SCREENING RECOMMENDATIONS**

2 The 2008 clinical practice guideline by the U.S. Department of Health and Human Services
3 advises that healthcare providers should inquire about tobacco use among all patients and
4 consistently document this information in their medical records (Fiore et al., 2008).

5
6 The inclusion of tobacco use status has been recommended in patient intake forms and
7 clinic screening systems as a fifth vital sign.

8
9 The strength of evidence for this recommendation was designated as Level A, meaning that
10 “multiple well-designed randomized clinical trials, directly relevant to the
11 recommendation, yielded a consistent pattern of findings.” In a meta-analysis of nine
12 studies, it was found that including patient report of tobacco use status in patient records
13 through the use of screening systems significantly increased the rate of clinician
14 intervention. However, a meta-analysis showed that use of a clinic system to identify and
15 track patients’ tobacco use status, alone, did not significantly increase rates of cessation.

16
17 In addition, the USPSTF (2021) provided the following recommendations for adults:

18 *Grade A Recommendation:* Clinicians should ask all adults about tobacco use, advise them
19 to stop using tobacco and provide behavioral interventions and US Food and Drug
20 Administration (FDA) approved pharmacotherapy for cessation to nonpregnant adults who
21 use tobacco.

22
23 *Grade A Recommendation:* Clinicians should ask all pregnant persons about tobacco use,
24 advise them to stop using tobacco and provide behavioral interventions for cessation to
25 pregnant persons who use tobacco.

26
27 **Documentation Requirements to Substantiate Medical Necessity**

CPT® Code	CPT® Code Description
99406	Smoking and tobacco use cessation counseling visit; intermediate, greater than three (3) minutes up to 10 minutes
99407	Smoking and tobacco use cessation counseling visit; intensive, greater than 10 minutes

1 The 5As comprise a framework frequently used in clinical practice to guide behavioral
 2 interventions. The following s are examples of how each step can be applied to tobacco
 3 cessation.

4
 5 **1) Ask:**

- 6 • Ask every patient about tobacco use at each visit.
- 7 • Record the response in the patient’s chart.

8
 9 **2) Advise:**

- 10 • Provide the patient with a clear, non-judgmental statement about how important it
 11 is to stop smoking.
- 12 • Discuss the increased risk of tobacco use to the patient’s health.
- 13 • Discuss benefits of quitting for health, family, and economics (e.g., cost savings).

14
 15 **3) Assess:**

- 16 • Ask the patient about their willingness to quit.
- 17 • Provide interventions to a patient not yet willing to quit. Explore why they are not
 18 motivated to quit at this time. What are the advantages and disadvantages of
 19 smoking? Identify the patient’s core values and how they are related to tobacco use.
- 20 • If they are willing to quit, offer brief intervention, referral sources, schedule follow
 21 up plan.
- 22 • Assess for any medical and/or psychological condition(s) which may contraindicate
 23 or complicate tobacco cessation (e.g., COPD, schizophrenia). Consultation with the
 24 primary care physician in such circumstances should be obtained prior to cessation
 25 of tobacco use.

26
 27 **4) Assist:**

- 28 • Help the patient make a quit plan. Set a date, ideally within 2 weeks.
- 29 • Help the patient change their environment (e.g., cleaning ash trays out of the home).
- 30 • Assist patient with establishing a social support system for help with quitting.
- 31 • Identify and plan for dealing with tobacco triggers or other challenges before or
 32 after quitting tobacco (e.g., co-workers who smoke, stress).
- 33 • Discuss relapse prevention; Develop coping skills to maintain a desire to quit.
- 34 • Ask about the patient’s interest in medications and refer if medications are desired.
- 35 • Provide supplemental self-help materials and referrals information such as quit
 36 lines.
- 37 • Assess for environmental barriers (e.g., others smoking at home).
- 38 • Discuss previous quit attempts (successes and/or barriers).
- 39 • Given alcohol’s relation to relapse, consider limiting use while quitting.
- 40 • Discuss nicotine withdrawal symptoms.

- 1 • Discuss steps taken prior to quitting, such as removing all tobacco products from
- 2 the patient’s environment and avoiding using tobacco/nicotine products in places
- 3 where the patient spends the majority of their time.
- 4 • Discuss making home smoke-free.
- 5 • Provide support to address family and friends who use tobacco.
- 6 • Provide encouragement and support to quit.

7

8 **5) Arrange:**

- 9 • Follow up within the first week after quit date and again within the first month.
- 10 • Follow-up can be by phone, texting, in person or by e-mail.
- 11 • Congratulate successes.
- 12 • Provide out of office visit clinician support to maintain quitting (e.g., email, phone,
- 13 texting, walk-in).
- 14 • Encourage and support a prolonged quitting.
- 15 • Relapse prevention/intervention to support long term tobacco cessation.
- 16 • Reinforce the positive health benefits immediately following quitting and for
- 17 prolonged cessation.
- 18 • Records should indicate that patients participating in a tobacco cessation program
- 19 are asked about their tobacco use at every visit (prior, during, and after quitting).

20

21 According to the USPSTF, there is a dose-response relationship between quit rates and the

22 intensity of counseling (that is, more or longer sessions improve quit rates). Quit rates

23 appear to plateau after 90 minutes of total counseling contact time. Combination therapy

24 with counseling and medications is more effective at increasing cessation rates than either

25 component alone.

26

27 **EFFECTIVE INTERVENTIONS**

28 The practitioner should carry out an assessment to determine the most appropriate course

29 of tobacco cessation treatment for the patient. Figure 1 (below) provides a guideline


30 (adapted from Hughes, 2013) that the practitioner can utilize for the assessment and

31 management of tobacco cessation treatment. If the counseling and/or medication

32 interventions are outside of the expertise and scope of practice of the practitioner, then it

33 is helpful to educate the patient that counseling and medication can be effective and refer

34 the patient to an appropriate health care professional for further assistance.



1
2 Figure 1: Tobacco Cessation Intervention Assessment Algorithm (adapted from Hughes,
3 2013)

1 **Counseling**

2 Community-based tobacco-control programs have been effective, judging by the decline
3 in adult smoking prevalence in the U.S. from 20.9% in 2005 to 11.5% in 2021 (Centers for
4 Disease Control and Prevention, 2023). At the individual level, it has been documented that
5 personalized advice from their doctor influences patients to quit, when compared to
6 patients not advised to quit. Brief counseling of 3 minutes or less by a physician has been
7 shown to be effective in achieving prolonged abstinence, compared to no intervention.
8 Higher-intensity counseling sessions >10 minutes have achieved abstinence rates of 22.1%,
9 nearly twice those of brief counseling of <3 minutes, 13.4%. Use of state quit lines for
10 telephone counseling has been shown to be effective compared to no counseling or self-
11 help only.

12
13 **Evidence-Based Behavior Modification Techniques**

14 A recommendation published by the Department of Health and Human Services (Fiore et
15 al., 2008) reported that 2 types of counseling and behavioral therapies result in higher
16 abstinence rates: (a) providing smokers with practical counseling (problem-solving
17 skills/skills training), and (b) providing support and encouragement as part of treatment.
18 The panel recommended that these types of counseling elements should be included in
19 smoking cessation interventions. Examples of these include:

- 20 • **Problem solving/ skills training:** Recognize danger situations – Identify events,
21 internal states, or activities that increase the risk of smoking or relapse.
- 22 • **Develop coping skills:** Identify and practice coping or problem-solving skills.
23 Typically, these skills are intended to cope with danger situations.
- 24 • **Provide basic information:** Provide basic information about smoking and
25 successful quitting.

26
27 According to a meta-analysis by Hartmann-Boyce et al. (2021), smoking cessation rates
28 can be increased at 6 months or longer through behavioral support without evidence that
29 suggests that there is increased harm. This is true whether psychopharmacotherapy is also
30 provided, although this effect is slightly more pronounced when the latter is absent. In fact,
31 evidence of benefits is strongest when counseling of any kind is employed and guaranteed
32 financial incentives. There might also be benefit from interventions that are more
33 individually tailored; delivered by text message, email, or audio recording; delivered by a
34 lay health advisor; and content with motivational components, as well as a focus on how
35 to quit.

36
37 Counseling can be effective when used alone, however the combination of counseling and
38 medication is more effective than either strategy used on its own (Stead et al., 2012). The
39 use of medications is effective in combination with counseling, except for situations in
40 which it may be contraindicated, or with populations in which medication use has not been
41 found to be effective.

1 Medications

2 In addition to counseling, all smokers making a quit attempt may be offered medications,
3 or referrals for medication evaluations as appropriate.

4
5 Though evidence and guidelines suggest medication, practitioners must consult within
6 their scope of licensure. This guideline does not suggest communication about medication
7 if such activity is outside the practitioner’s scope of practice.

8
9 Over-the-counter products include nicotine gum, patches, and lozenges. It is important to
10 thoroughly review the directions prior to use. Prescription nicotine replacement products
11 include nasal and oral inhalers. Oral prescription medications for tobacco cessation that do
12 not contain nicotine include bupropion and varenicline.

13
14 Electronic cigarettes (e-cigarettes or electronic nicotine delivery systems) are a group of
15 products that generally provide aerosolized nicotine without the use of tobacco. They are
16 readily available to the public and are touted as an aid to tobacco cessation or as a
17 replacement for cigarettes where smoking is prohibited (Grana, 2014). The short- and long-
18 term efficacy and comparative efficacy with approved tobacco cessation products is not
19 yet fully known. A systematic review showed good results with smoking abstinence at one
20 month, but abstinence at 3 and 6 months was the same as placebo (McRobbie et al., 2014).
21 The toxicity of e-cigarettes is not yet clear and further research is needed to evaluate their
22 safety for the direct user and those with second-hand exposure (Hartmann-Boyce et al.,
23 2021). When used as a therapeutic intervention, the use of e-cigarettes may have a negative
24 effect on nicotine abstinence in comparison to nicotine replacement therapies. This is, most
25 smokers who quit smoking cigarettes with the help of e-cigarettes continued using e-
26 cigarettes until the end of random controlled trials (Hanewinkel et al., 2022).

27
28 Nicotine withdrawal symptoms include irritability, cravings, depression, anxiety, cognitive
29 and attention deficits, sleep disturbances, and increased appetite. These symptoms may
30 begin within a few hours after the last cigarette, quickly driving people back to tobacco
31 use. Symptoms peak within the first few days of smoking cessation and may subside within
32 a few weeks. For some people, however, symptoms may persist for months. The former
33 tobacco user should receive recognition of any success made during a quit attempt and
34 receive strong encouragement to remain abstinent. Relapse is most likely to occur soon
35 after quitting, but the risk for relapse can continue for months, or even years. All very
36 recent quitters should be given assistance; therefore, it is important to regularly ask those
37 who have quit if they are facing any challenges, such as temptations to smoke, close calls
38 for slips and relapses, or serious thoughts about starting again. Former tobacco users who
39 report such challenges should be given additional tobacco cessation assistance.

1 PRACTITIONER SCOPE AND TRAINING

2 Practitioners should practice only in the areas in which they are competent based on their
3 education training and experience. Levels of education, experience, and proficiency may
4 vary among individual practitioners. It is ethically and legally incumbent on a practitioner
5 to determine where they have the knowledge and skills necessary to perform such services
6 and whether the services are within their scope of practice.

7
8 It is best practice for the practitioner to appropriately render services to a patient only if
9 they are trained to competency, equally skilled, and adequately competent to deliver a
10 service compared to others trained to perform the same procedure. If the service would be
11 most competently delivered by another health care practitioner who has more skill and
12 training, it would be best practice to refer the patient to the more expert practitioner.

13
14 Best practice can be defined as a clinical, scientific, or professional technique, method, or
15 process that is typically evidence-based and consensus driven and is recognized by a
16 majority of professionals in a particular field as more effective at delivering a particular
17 outcome than any other practice (Joint Commission International Accreditation Standards
18 for Hospitals, 2020).

19
20 Depending on the practitioner’s scope of practice, training, and experience, a patient’s
21 condition and/or symptoms during examination or the course of treatment may indicate the
22 need for referral to another practitioner or even emergency care. In such cases it is essential
23 for the practitioner to refer the patient for appropriate co-management (e.g., to their primary
24 care physician) or if immediate emergency care is warranted, to contact 911 as appropriate.
25 See the *Managing Medical Emergencies (CPG 159 – S)* clinical practice guideline for
26 information.

27 PRACTITIONER RESOURCES

28 One way to assist patients with tobacco cessation is by using a tear sheet. The tear sheet
29 can allow clinicians to individualize an intervention and can be given to patients as a
30 takeaway.

- 31 • Tear Sheet for Use with Patients – English
32 ([http://www.ahrq.gov/sites/default/files/wysiwyg/professionals/clinicians-
35 providers/guidelines-
36 recommendations/tobacco/clinicians/tearsheets/tearsheet.pdf](http://www.ahrq.gov/sites/default/files/wysiwyg/professionals/clinicians-
33 providers/guidelines-
34 recommendations/tobacco/clinicians/tearsheets/tearsheet.pdf))
- 37 • Quick Reference Guide for Clinicians
38 ([https://www.ahrq.gov/sites/default/files/wysiwyg/professionals/clinicians-
41 providers/guidelines-
42 recommendations/tobacco/clinicians/references/quickref/tobaqrg.pdf](https://www.ahrq.gov/sites/default/files/wysiwyg/professionals/clinicians-
39 providers/guidelines-
40 recommendations/tobacco/clinicians/references/quickref/tobaqrg.pdf))
- 43 • Material Links for Clinical Websites and Blogs (<http://www.smokefree.gov/>)

1 Spanish Language

- 2 • Tear Sheet for Use with Patients - Spanish
- 3 ([http://www.ahrq.gov/professionals/clinicians-providers/guidelines-](http://www.ahrq.gov/professionals/clinicians-providers/guidelines-recommendations/tobacco/clinicians/tearsheets/tearsheetssp.html)
- 4 [recommen-dations/tobacco/clinicians/tearsheets/tearsheetssp.html](http://www.ahrq.gov/professionals/clinicians-providers/guidelines-recommendations/tobacco/clinicians/tearsheets/tearsheetssp.html))
- 5 • Material links for Clinical Websites and Blogs – Spanish
- 6 (<https://espanol.smokefree.gov/>)

8 MEMBER RESOURCES

9 Educating patients about tobacco cessation options and available resources can assist the
10 patient. Publicly available resources can be found at:

- 11 • How to Quit Smoking (<https://www.cdc.gov/tobacco/about/how-to-quit.html>)
- 12 • Tools to Help You Quit (<https://www.smokefree.gov/>)
- 13 • Understanding Tobacco Cravings
- 14 ([https://www.ashlink.com/ASH/public/Members/PreventiveHealthProgram.aspx?](https://www.ashlink.com/ASH/public/Members/PreventiveHealthProgram.aspx?nameid=UnderstandingTobaccoCravings&category=Tobacco)
- 15 [nameid=UnderstandingTobaccoCravings&category=Tobacco](https://www.ashlink.com/ASH/public/Members/PreventiveHealthProgram.aspx?nameid=UnderstandingTobaccoCravings&category=Tobacco))

16
17 Federal resources are available to patients to assist in quitting tobacco products:

- 18 • Visit <https://smokefree.gov/>
- 19 • Visit the CDC’s website on how to quit smoking (with links to Spanish content as
20 well): <https://www.cdc.gov/tobacco/campaign/tips/quit-smoking/index.html>
- 21 • Talk to a Smoking Cessation Counselor
 - 22 ○ Call 1-800-QUITNOW (1-800-784-8669), a national portal to a network
 - 23 of state quitlines
 - 24 ○ American Lung Association: Lung Helpline and Tobacco Quitline:
 - 25 ▪ 1-800-LUNG-USA (1-800-586-4872) & for the hearing impaired
 - 26 TTY 1-800-501-1068
- 27 • Get Instant Messaging Live Help
- 28 (https://livehelp.cancer.gov/app/chat/chat_launch)
- 29 • Approved Smoking Cessation Products
- 30 (<http://www.fda.gov/ForConsumers/ConsumerUpdates/ucm198176.htm>)

32 References

33 Ahluwalia JS, Gibson CA, Kenney RE, Wallace DD, Resnicow K. Smoking status as a
34 vital sign. *J Gen Intern Med.* 1999;14(7):402-408

35
36 American Medical Association. Current Procedural Terminology (CPT): Current Year.
37 Rev ed. Chicago, IL: American Medical Association

38
39 Centers for Disease Control and Prevention. About E-Cigarettes (Vapes). Retrieved on
40 February 23, 2026 from <https://www.cdc.gov/tobacco/e->

1 cigarettes/about.html?CDC_AAref_Val=https://www.cdc.gov/tobacco/basic_informat
2 ion/e-cigarettes/about-e-cigarettes.html

3
4 Centers for Disease Control and Prevention. Current cigarette smoking among adults in the
5 United States. Retrieved February 23, 2026, from
6 <https://www.cdc.gov/tobacco/php/data-statistics/adult-data-cigarettes/index.html>

7
8 Cherkin DC, Deyo RA, Sherman KJ, et al. Characteristics of visits to licensed
9 acupuncturists, chiropractors, massage therapists, and naturopathic physicians. *J Am*
10 *Board Fam Pract.* Nov-Dec 2002;15(6):463-472

11
12 Cho, ER et al. Smoking Cessation and Short- and Longer-Term Mortality. *NEJM*
13 *evidence* vol. 3,3 (2024): EVIDoa2300272. doi:10.1056/EVIDoa2300272

14
15 Doll R, Hill AB. The mortality of doctors in relation to their smoking habits: A preliminary
16 report. 1954. *BMJ.* 2004;328(7455):1529-1533; discussion 1533

17
18 Doll R, Peto R, Boreham J, Sutherland I. Mortality in relation to smoking: 50 years'
19 observations on male British doctors. *BMJ.* 2004;328(7455):1519

20
21 Farelley M, Pechacek, TF, Thomas, KY, Nelson, D. The impact of tobacco control
22 programs on adult smoking. *Am J Public Health.* Feb 2008 2008;98(2):304-309

23
24 Fiore MC, Jaén CR, Baker TB, et al. Treating Tobacco Use and Dependence: 2008
25 Update. Clinical Practice Guideline. Rockville, MD: U.S. Department of Health and
26 Human Services. Public Health Service. May 2008 Retrieved on February 23, 2026
27 from <https://www.ncbi.nlm.nih.gov/books/NBK63952/>

28
29 Goldberg MS, Scott SC, Mayo NE. A review of the association between cigarette smoking
30 and the development of nonspecific back pain and related outcomes. *Spine (Phila Pa*
31 *1976).* Apr 15 2000;25(8):995-1014

32
33 Grana R, Benowitz N, Glantz SA. E-cigarettes: a scientific review. *Circulation.*
34 2014;129(19):1972-1986. <https://doi.org/10.1161/CIRCULATIONAHA.114.007667>

35
36 Hanewinkel R, Niederberger K, Pedersen A, Unger JB, Galimov A. E-cigarettes and
37 nicotine abstinence: A meta-analysis of randomised controlled trials. *Eur Respir Rev.*
38 2022;31(163):210215. <https://doi.org/10.1183/16000617.0215-2021>

39
40 Hartmann-Boyce J, McRobbie H, Lindson N, Bullen C, Begh R, Theodoulou A, Notley C,
41 Rigotti NA, Turner T, Butler AR, Fanshawe TR, Hajek P. Electronic cigarettes for

- 1 smoking cessation. *Cochrane Database Syst Rev.* 2022;4(4):CD010216.
2 <https://doi.org/10.1002/14651858.CD010216.pub5>
3
- 4 Hartmann-Boyce J, Livingstone-Banks J, Ordóñez-Mena JM, Fanshawe TR, Lindson N,
5 Freeman SC, Sutton AJ, Theodoulou A, Aveyard P. Behavioural interventions for
6 smoking cessation: an overview and network meta-analysis. *Cochrane Database Syst*
7 *Rev.* 2021;1:CD013229
8
- 9 Hays JT, Ebbert JO, Sood A. Treating tobacco dependence in light of the 2008 US
10 Department of Health and Human Services clinical practice guideline. *Mayo Clin Proc.*
11 *Aug* 2009;84(8):730-735; quiz 735-736
12
- 13 Hill FJ. Complementary and alternative medicine: the next generation of health promotion?
14 *Health Promot Int.* Sep 2003;18(3):265-272
15
- 16 Hughes J. An updated algorithm for choosing among smoking cessation treatments. *J Subst*
17 *Abuse Treat.* 2013;45(2):215
18
- 19 Joint Commission International. *Joint Commission International Accreditation Standards*
20 *for Hospitals.* 7th ed. Oak Brook, IL: Joint Commission Resources; 2020
21
- 22 Kramarow EA, Elgaddal N. Current electronic cigarette use among adults aged 18 and
23 over: United States, 2021. Retrieved February 23, 2026 from
24 <https://www.cdc.gov/nchs/data/databriefs/db475.pdf>
25
- 26 Kreuter MW, Chheda SG, Bull FC. How does physician advice influence patient behavior?
27 Evidence for a priming effect. *Arch Fam Med.* 2000;9(5):426-433
28
- 29 McGeary DD, Mayer TG, Gatchel RJ, Anagnostis C. Smoking status and
30 psychosocioeconomic outcomes of functional restoration in patients with chronic
31 spinal disability. *Spine J.* Mar-Apr 2004;4(2):170-175
32
- 33 McRobbie H, Bullen C, Hartmann-Boyce J, Hajek P. Electronic cigarettes for smoking
34 cessation and reduction. *Cochrane Database Syst Rev.* 2014;(12):CD010216.
35 <https://doi.org/10.1002/14651858.CD010216.pub2>
36
- 37 MedlinePlus [Internet]. Smoking cessation medications. Retrieved on February 23, 2026
38 from <https://medlineplus.gov/ency/article/007439.htm>
39
- 40 Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual causes of death in the United
41 States, 2000. *JAMA.* 2004;291(10):1238-1245

- 1 National Institute on Drug Abuse. Tobacco, Nicotine, and E-Cigarettes Research Report.
 2 Retrieved on February 23, 2026 from
 3 [https://www.drugabuse.gov/publications/research-reports/tobacco-nicotine-e-](https://www.drugabuse.gov/publications/research-reports/tobacco-nicotine-e-cigarettes/introduction)
 4 [cigarettes/introduction](https://www.drugabuse.gov/publications/research-reports/tobacco-nicotine-e-cigarettes/introduction)
 5
- 6 Okuyemi KS, Nollen NL, Ahluwalia JS. Interventions to facilitate smoking cessation. *Am*
 7 *Fam Physician*. 2006;74(2):262-271
 8
- 9 Patel MS, Steinberg MB. In the Clinic. Smoking Cessation. *Ann Intern Med*.
 10 2016;164(5):ITC33-ITC48. <https://doi.org/10.7326/AITC201603010>
 11
- 12 Pincus T, Santos R, Breen A, Burton AK, Underwood M, Multinational Musculoskeletal
 13 Inception Cohort Study Collaboration. A review and proposal for a core set of factors
 14 for prospective cohorts in low back pain: a consensus statement. *Arthritis Rheum*.
 15 2008;59(1):14-24. <https://doi.org/10.1002/art.23251>
 16
- 17 Rehtine GR, 2nd, Frawley W, Castellvi A, Gowski A, Chrin AM. Effect of the spine
 18 practitioner on patient smoking status. *Spine (Phila Pa 1976)*. Sep 1 2000;25(17):2229-
 19 2233
 20
- 21 Stead LF, Lancaster T. Combined pharmacotherapy and behavioral interventions for
 22 smoking cessation. *Cochrane Database Syst Rev*. 2012;10:CD008286
 23
- 24 U.S. Food and Drug Administration. E-Cigarettes, Vapes, and other Electronic Nicotine
 25 Delivery Systems (ENDS). Retrieved on February 23, 2026 from
 26 [https://www.fda.gov/tobacco-products/products-ingredients-components/e-cigarettes-](https://www.fda.gov/tobacco-products/products-ingredients-components/e-cigarettes-vapes-and-other-electronic-nicotine-delivery-systems-ends)
 27 [vapes-and-other-electronic-nicotine-delivery-systems-ends](https://www.fda.gov/tobacco-products/products-ingredients-components/e-cigarettes-vapes-and-other-electronic-nicotine-delivery-systems-ends)
 28
- 29 U.S. Preventive Services Task Force (USPSTF). Tobacco Smoking Cessation in Adults,
 30 Including Pregnant Persons: Interventions. U.S. Public Health Service Office of the
 31 Surgeon General; National Center for Chronic Disease Prevention and Health
 32 Promotion (US) Office on Smoking and Health. Smoking Cessation: A Report of the
 33 Surgeon General [Internet]. Washington, DC: US Department of Health and Human
 34 Services; 2020. Chapter 6, Interventions for Smoking Cessation and Treatments for
 35 Nicotine Dependence. Available from:
 36 <https://www.ncbi.nlm.nih.gov/books/NBK555596/>
 37
- 38 Whitlock EP, Orleans CT, Pender N, Allan J. Evaluating primary care behavioral
 39 counseling interventions: An evidence-based approach. *Am J Prev Med*.
 40 2002;22(4):267-284