

1 **Clinical Policy:** **Preventive Care Services**

2  
3 **Date of Implementation:** **April 19, 2012**

4  
5 **Effective Date:** **February 19, 2026**

6  
7 **Product:** **Specialty**

8  
9  
10 **GUIDELINES**

11 American Specialty Health – Specialty (ASH) considers comprehensive preventive  
12 medicine evaluation and management (E/M) services provided by a qualified healthcare  
13 professional as medically necessary for detecting risk factors based on patient age, gender,  
14 and risk-status. Preventive medicine E/M services should follow established guidelines as  
15 determined by the U.S. Preventive Services Task Force (USPSTF).

16  
17 Preventive medicine counseling is established as clinically effective and professionally  
18 recognized for providing patient assessment and related provision of evidence-based  
19 information about self-care, risk factor reduction and professional interventions to more  
20 effectively manage high risk health behaviors and to pursue appropriate self-care and  
21 screening activities. Preventive medicine management focuses on health concerns such as  
22 injury prevention; alcohol or drug use; tobacco use; inadequate nutrition and diet; sedentary  
23 lifestyle; pursuing appropriate disease screening and immunizations; and the appropriate  
24 management of serum glucose, hypertension, and obesity. Preventive medicine counseling  
25 and risk factor reduction interventions should follow USPSTF established guidelines.

26  
27 Other CPGs related to specific screening and preventive care recommendations are  
28 available on [ashlink.com](http://ashlink.com).

29  
30 **INTRODUCTION**

31 The Agency for Healthcare Research and Quality (AHRQ) is under the U.S. Department  
32 of Health and Human Services (HHS) and sponsors the USPSTF, the leading independent  
33 panel of private-sector experts in prevention and primary care. The USPSTF conducts  
34 rigorous, impartial assessments of scientific evidence for the effectiveness of a broad range  
35 of clinical preventive services, including screening, counseling, and preventive  
36 medications. Its recommendations are considered the "gold standard" for clinical  
37 preventive services.

38  
39 By evaluating the benefits of individual services based on age, gender, and risk factors for  
40 disease, the USPSTF makes recommendations about which preventive services should be  
41 incorporated routinely into primary medical care and for which populations.

1 Preventive Care Guidelines are standards of care developed to encourage the appropriate  
 2 provision of preventive services to patients, according to their age, gender, and risk-status.  
 3 These services include screenings, immunizations, and physical examinations.

4  
 5 Preventive services are recommendations for when the participant does not have any other  
 6 health risks or a diagnosed health condition. Participants with any health risk or diagnosis  
 7 may have additional health care service recommendations or the frequency at which the  
 8 services are recommended by their practitioners may vary.

9  
 10 Preventive medicine evaluation and management services are comprehensive in nature,  
 11 reflect an age and gender appropriate history and examination, and include counseling,  
 12 anticipatory guidance, and risk factor reduction interventions, usually separate from  
 13 disease-related diagnoses. A spinal screening (e.g., evaluating posture, gait, range of  
 14 motion, and joints for subluxations) by itself does not meet the above-described criteria,  
 15 and would not be considered a preventive medicine evaluation and management service.

#### 16 **U.S. PREVENTIVE SERVICES TASK FORCE (USPSTF) GRADING SCALE**

17 The USPSTF assigns 1 of 5 letter grades to each of its recommendations (A, B, C, D, or I).  
 18 It also has defined levels of certainty regarding net benefit. See tables below.

Grade	Definition	Suggestions for Practice
<b>A</b>	The USPSTF recommends the service. There is high certainty that the net benefit is substantial.	Offer or provide this service.
<b>B</b>	The USPSTF recommends the service. There is high certainty that the net benefit is moderate or there is moderate certainty that the net benefit is moderate to substantial.	Offer or provide this service.
<b>C</b>	The USPSTF recommends selectively offering or providing this service to individual patients based on professional judgment and patient preferences. There is at least moderate certainty that the net benefit is small.	Offer or provide this service for <b>selected</b> patients depending on individual circumstances.
<b>D</b>	The USPSTF recommends against the service. There is moderate or high certainty that the service has no net benefit or that the harms outweigh the benefits.	Discourage the use of this service.

<b>Grade</b>	<b>Definition</b>	<b>Suggestions for Practice</b>
<b>I</b>	The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of the service. Evidence is lacking, of poor quality, or conflicting, and the balance of benefits and harms cannot be determined.	Read the clinical considerations section of USPSTF Recommendation Statement. If the service is offered, patients should understand the uncertainty about the balance of benefits and harms.

1  
2

**LEVELS OF CERTAINTY REGARDING NET BENEFIT**

<b>Level of Certainty*</b>	<b>Description</b>
<b>High</b>	The available evidence usually includes consistent results from well-designed, well-conducted studies in representative primary care populations. These studies assess the effects of the preventive service on health outcomes. This conclusion is therefore unlikely to be strongly affected by the results of future studies.
<b>Moderate</b>	The available evidence is sufficient to determine the effects of the preventive service on health outcomes, but confidence in the estimate is constrained by such factors as: <ul style="list-style-type: none"> <li>• The number, size, or quality of individual studies.</li> <li>• Inconsistency of findings across individual studies.</li> <li>• Limited generalizability of findings to routine primary care practice.</li> <li>• Lack of coherence in the chain of evidence.</li> </ul> As more information becomes available, the magnitude or direction of the observed effect could change, and this change may be large enough to alter the conclusion.
<b>Low</b>	The available evidence is insufficient to assess effects on health outcomes. Evidence is insufficient because of: <ul style="list-style-type: none"> <li>• The limited number or size of studies.</li> <li>• Important flaws in study design or methods.</li> <li>• Inconsistency of findings across individual studies.</li> <li>• Gaps in the chain of evidence.</li> <li>• Findings not generalizable to routine primary care practice.</li> <li>• Lack of information on important health outcomes.</li> </ul> More information may allow estimation of effects on health outcomes.

\* The USPSTF defines certainty as "likelihood that the USPSTF assessment of the net benefit of a preventive service is correct." The net benefit is defined as benefit minus harm of the preventive service as implemented in a general, primary care population. The USPSTF assigns a certainty level based on the nature of the overall evidence available to assess the net benefit of a preventive service.

**QUALITY OF EVIDENCE**

The USPSTF grades the quality of the overall evidence for a service on a 3-point scale (good, fair, and poor):

**Good:** Evidence includes consistent results from well-designed, well-conducted studies in representative populations that directly assess effects on health outcomes.

**Fair:** Evidence is sufficient to determine effects on health outcomes, but the strength of the evidence is limited by the number, quality, or consistency of the individual studies, generalizability to routine practice, or indirect nature of the evidence on health outcomes.

**Poor:** Evidence is insufficient to assess the effects on health outcomes because of limited number or power of studies, important flaws in their design or conduct, gaps in the chain of evidence, or lack of information on important health outcomes.

**CPT Code and Documentation Requirements to Substantiate Medical Necessity**

Preventive Medicine Assessments CPT Code Description	
<i>Initial</i> comprehensive preventive medicine evaluation and management of an individual including an age and gender appropriate history, examination, counseling/anticipatory guidance/risk factor reduction interventions, and the ordering of laboratory/diagnostic procedures, new patient; <u>by age</u> .	
CPT Code	Applicable Age Range
99381	infant (age younger than 1 year)
99382	early childhood (age 1 through 4 years)
99383	late childhood (age 5 through 11 years)
99384	adolescent (age 12 through 17 years)
99385	18-39 years
99386	40-64 years
99387	65 years and older

1 Preventive Medicine Counseling services are time-based and should follow established  
 2 guidelines as determined by the USPSTF and other sources described in this document.  
 3 Such annual visits should include the following:

- 4 • An assessment of common risk factors (age and at-risk behavior dependent);
- 5 • Discussion and identification of risks within each of the appropriate categories;
- 6 • Discussion of intervention options; and
- 7 • Programmatic options where available which are in line with USPSTF  
 8 recommendations.
- 9 • Evidence-based services that have in effect an A or B grade according to the current  
 10 recommendations of the United States Preventive Services Task Force (Task Force)  
 11 with respect to the individual involved.

12  
 13 An example of code 99383 would be: a 9-year-old male presenting for his first health  
 14 supervision and evaluation. The practitioner reviews his complete medical (including  
 15 immunizations), family, social, religious, and cultural history. A complete review of  
 16 systems and complete physical are performed. Speech and blood pressure are checked.  
 17 Growth, development and the patient's behavior are assessed. Anticipatory guidance is  
 18 provided to the patient regarding good health habits and self-care, and to the parents  
 19 regarding good parenting practices. Risk factors are identified, and interventions discussed.  
 20 Medically appropriate lab tests are ordered if indicated.

21  
 22 Medical record documentation should reflect the procedures performed and the clinical  
 23 rationale for them (e.g., age appropriate and/or at-risk behavior).  
 24

<b>Preventive Medicine Counseling CPT Code Description</b>	
Preventive medicine counseling and/or risk factor reduction intervention(s) provided to an individual [codes 99401 – 99404] or in a group setting [codes 99411 – 99412] (separate procedure) <u>by time</u>	
<b>CPT CODE</b>	<b>Applicable Time</b>
99401	approximately 15 minutes
99402	approximately 30 minutes
99403	approximately 45 minutes
99404	approximately 60 minutes
99411	approximately 30 minutes
99412	approximately 60 minutes

1 An example of code 99404 would be: a 31-year-old female patient who smokes, is  
 2 sedentary and obese, and has familial hypercholesterolemia, meets with her practitioner to  
 3 discuss dietary and lifestyle changes to reduce her risk of heart disease. The discussion  
 4 would focus on her risk of heart disease based on her current condition and family history,  
 5 as well as discussion of the potential benefits of losing weight, giving up smoking, and  
 6 lowering her cholesterol. The following options would also be reviewed, and assistance  
 7 provided with prioritizing her goals for smoking cessation, diet change, and weight loss.

8  
 9 Medical record documentation should reflect the procedures performed and the clinical  
 10 rationale for them (e.g., age appropriate and/or at-risk behavior).

### 11 **Practitioner Resources**

- 12 • Health Recommendations Tool (<http://epss.ahrq.gov>)
- 13 • Quick Guide to Healthy Living (<https://health.gov/myhealthfinder>)
- 14 • Home Page - Healthypeople (<https://health.gov/healthypeople>)

### 15 **Member Resources**

- 16 • Preventive Health  
 17 (<http://healthfinder.gov>)
- 18 • Preventive Health Topics  
 19 (<http://www.nlm.nih.gov/medlineplus/healthtopics.html>)

## 20 **PRACTITIONER SCOPE AND TRAINING**

21 Practitioners should practice only in the areas in which they are competent based on their  
 22 education, training and experience. Levels of education, experience, and proficiency may  
 23 vary among individual practitioners. It is ethically and legally incumbent on a practitioner  
 24 to determine where they have the knowledge and skills necessary to perform such services  
 25 and whether the services are within their scope of practice.

26 It is a best practice for the practitioner to appropriately render services to a patient only if  
 27 they are trained to competency, equally skilled, and adequately competent to deliver a  
 28 service compared to others trained to perform the same procedure. If the service would be  
 29 most competently delivered by another health care practitioner who has more skill and  
 30 training, it would be best practice to refer the patient to the more expert practitioner.

31 Best practice can be defined as a clinical, scientific, or professional technique, method, or  
 32 process that is typically evidence-based and consensus driven and is recognized by a  
 33 majority of professionals in a particular field as more effective at delivering a particular  
 34 outcome than any other practice (Joint Commission International Accreditation Standards  
 35 for Hospitals, 2020).

1 Depending on the practitioner’s scope of practice, training, and experience, a patient’s  
 2 condition and/or symptoms during examination or the course of treatment may indicate the  
 3 need for referral to another practitioner or even emergency care. In such cases it is essential  
 4 for the practitioner to refer the patient for appropriate co-management (e.g., to their primary  
 5 care physician) or if immediate emergency care is warranted, to contact 911 as appropriate.  
 6 See the *Managing Medical Emergencies (CPG 159 – S)* clinical practice guideline for  
 7 information.

## 9 **References**

10 American Academy of Family Physicians (AAFP). *AAFP Clinical Preventive Services*  
 11 *Recommendations* Retrieved on January 4, 2026 from [https://www.aafp.org/family-](https://www.aafp.org/family-physician/patient-care/clinical-recommendations/aafp-cps.html)  
 12 [physician/patient-care/clinical-recommendations/aafp-cps.html](https://www.aafp.org/family-physician/patient-care/clinical-recommendations/aafp-cps.html)

14 American Medical Association. (current year). *Current Procedural Terminology (CPT)*  
 15 *Current year* (rev. ed.). Chicago: AMA

17 Joint Commission International. (2020). *Joint Commission International Accreditation*  
 18 *Standards for Hospitals (7<sup>th</sup> ed.)*: Joint Commission Resources

20 U.S. Department of Health and Human Services. (2024). *Healthfinder*. Retrieved January  
 21 4, 2026 from <http://www.healthfinder.gov/>

23 U.S. Department of Health and Human Services, Health Resources and Services  
 24 Administration. (March 2024). *Women’s Preventive Services Guidelines*. Retrieved  
 25 January 4, 2026 from <https://www.hrsa.gov/womens-guidelines/index.html>

27 U.S. Preventive Services Task Force (USPSTF). *Grade definitions*. Retrieved January 4,  
 28 2026 from [https://www.uspreventiveservicestaskforce.org/uspstf/about-](https://www.uspreventiveservicestaskforce.org/uspstf/about-uspstf/methods-and-processes/grade-definitions#july2012.jsp)  
 29 [uspstf/methods-and-processes/grade-definitions#july2012.jsp](https://www.uspreventiveservicestaskforce.org/uspstf/about-uspstf/methods-and-processes/grade-definitions#july2012.jsp)

31 U.S. Preventive Services Task Force (USPSTF). *A and B Recommendations*. Retrieved  
 32 January 4, 2026 from  
 33 [https://www.uspreventiveservicestaskforce.org/uspstf/recommendation-topics/uspstf-](https://www.uspreventiveservicestaskforce.org/uspstf/recommendation-topics/uspstf-and-b-recommendations)  
 34 [and-b-recommendations](https://www.uspreventiveservicestaskforce.org/uspstf/recommendation-topics/uspstf-and-b-recommendations)

36 U.S. Preventive Services Task Force (USPSTF). *About the USPSTF: Task Force*  
 37 *Resources*. Retrieved January 4, 2026 from  
 38 [https://www.uspreventiveservicestaskforce.org/uspstf/about-uspstf/task-force-](https://www.uspreventiveservicestaskforce.org/uspstf/about-uspstf/task-force-resources)  
 39 [resources](https://www.uspreventiveservicestaskforce.org/uspstf/about-uspstf/task-force-resources)