

1 **Clinical Practice Guideline:** **Repair of Rupture of Foot and Ankle Flexor or**
 2 **Extensor Tendons**

4 **Date of Implementation:** **September 17, 2015**

6 **Product:** **Specialty**

9 **GUIDELINES**

10 A. American Specialty Health – Specialty (ASH) considers procedures consisting of
 11 **CPT® Code 27658, 27659, 27664, and 27665** to be medically necessary for repair of
 12 flexor or extensor tendons **for the following diagnoses:**

- 13 • Strain of Achilles tendon, sprain of other ligament of ankle, and strain of
 14 unspecified muscle and tendon at ankle and foot level (ICD-10 Codes S86.011A -
 15 S86.019S, S93.491A - S93.499S, S96.919A - S96.919S)
- 16 • Spontaneous rupture of tendons of foot and ankle (not amenable to casting or
 17 bracing) (ICD-10 Codes M66.271 - M66.279, M66.371 - M66.379, M66.871 -
 18 M66.879)
- 19 • Abscess of tendon sheath or bursa, other specified bursopathies, synovial
 20 hypertrophy, and other specified disorders of synovium and tendon of lower leg,
 21 ankle and foot, other site and multiple sites (ICD-10 Codes M65.061 - M65.079,
 22 M65.08, M67.261 - M67.279, M67.28 - M67.29, M67.871 - M67.879, M67.88 -
 23 M67.89, M71.071 - M71.079, M71.08 - M71.09, M71.871 - M71.879, M71.88 -
 24 M71.89)
- 25 • Unspecified disorder of synovium and tendon of lower leg, ankle and foot, other
 26 site, and multiple sites (not amenable to casting or bracing) (ICD-10 Codes
 27 M67.961 - M67.979, M67.98 - M67.99)

28 B. ASH considers procedures consisting of **CPT® Code 28200, 28202, 28208, and 28210**
 29 to be medically necessary for repair of the foot flexor or extensor tendons **for the**
 30 **following diagnoses:**

- 31 • Strain of Achilles tendon, sprain of other ligament of ankle, and strain of
 32 unspecified muscle and tendon at ankle and foot level (ICD-10 Codes S86.011A -
 33 S86.019S, S93.491A - S93.499S, S96.919A - S96.919S)
- 34 • Spontaneous rupture of tendons of foot and ankle (not amenable to casting or
 35 bracing) (ICD-10 Codes M66.271 - M66.279, M66.371 - M66.379, M66.871 -
 36 M66.879)

1 **CPT® Codes and Descriptions**

CPT® Code	CPT® Code Description
27658	Repair, flexor tendon, leg; primary, without graft, each tendon
27659	Repair, flexor tendon, leg; secondary, with or without graft, each tendon
27664	Repair, extensor tendon, leg; primary, without graft, each tendon
27665	Repair, extensor tendon, leg; secondary, with or without graft, each tendon
28200	Repair, tendon, flexor, foot; primary or secondary, without free graft, each tendon
28202	Repair, tendon, flexor, foot; secondary with free graft, each tendon (includes obtaining graft)
28208	Repair, tendon, extensor, foot; primary or secondary, each tendon
28210	Repair, tendon, extensor, foot; secondary with free graft, each tendon (includes obtaining graft)

2

3 **DESCRIPTION/BACKGROUND**

4 When a tendon ruptures it can be extremely painful and cause disability of the foot and
5 ankle. Ruptures of ankle and foot tendons can occur from a single traumatic event (sprains
6 and strains of ankle and foot, other tendon ICD-10 Codes S86.011A - S86.019S, S93.491A
7 - S93.499S, S96.919A - S96.919S). A nontraumatic rupture may occur in degenerative
8 tendons where tendinosis causes a gradual attenuation and tendon rupture, as a result of
9 certain systemic diseases, or steroid use among other possible factors. An untreated tendon
10 rupture can result in damage to the joints of the foot and ankle. Any of these structures may
11 rupture, resulting in a serious injury that may require surgical repair.

12

13 **Tibialis Anterior Tendon Rupture**

14 The tibialis anterior tendon is vulnerable to laceration because of its subcutaneous position
15 over the anterior aspect of the foot and ankle. Laceration of this tendon may weaken
16 dorsiflexion of the ankle and extension will be impacted; however, rupture of the tendon is
17 rare because the tendon will remain intact due to the secondary action of the extensor
18 hallucis longus (EHL) and extensor digitorum longus (EDL). The foot laceration should be
19 evaluated if there is suspicion of partial or complete tendon rupture. The structures heal
20 well and have minimal dysfunction when repaired acutely.

21

22 In rupture of the tibialis anterior tendon, the ruptured tendon end often becomes caught at
23 one of the extensor retinacular layers and may be easily palpated beneath the skin.
24 Nontraumatic tendon rupture usually occurs by partial avulsion from the insertion and
25 elongation of the degenerative tendon within the inferior extensor retinaculum.

1 Tibialis anterior deficiency commonly presents with spontaneous rupture of the tendon at
2 or near its insertion. It typically occurs in middle aged athletes and often accompanies other
3 comorbid conditions such as diabetes, inflammatory arthritis, gout, obesity, and steroid
4 use. The tendon in this location may demonstrate a zone of relative hypovascularity near
5 its insertion that may predispose it to tendinosis and rupture in this area. Repair may be
6 performed early or late, but the results of the surgery are better if repair is performed within
7 the first 3 to 6 weeks. Non-operative treatment is acceptable for elderly, inactive patients,
8 but primary or delayed repair is helpful for active individuals regardless of age.

9
10 Sammarco et al. (2009) performed a retrospective review of cases of patients with anterior
11 tibialis tendon rupture who had undergone operative procedures. Nineteen tibialis anterior
12 tendon ruptures were surgically repaired in eighteen patients ranging in age from 21 to 78
13 years. Early repair was performed for one traumatic and seven atraumatic ruptures three
14 days to six weeks after the injury. If the tendon ends could be approximated or if the tendon
15 could be brought to its insertion, the ends were debrided, and a direct tendon repair was
16 performed. In four cases, the tendon was repaired directly to bone with a suture anchor.
17 Direct repair was achieved in the cases of seven ruptures, including four that underwent
18 early repair and three that underwent delayed reconstruction. If the tendon ends could not
19 be approximated or if the tendon could not be placed onto its insertion site, an
20 interpositional tendon graft was used to bridge the gap and to reinforce the repair.

21 22 **Flexor Hallucis Longus Rupture**

23 Rupture of the flexor hallucis longus is uncommon but can occur in several areas. Tendon
24 rupture can occur following corticosteroid injection into the tendon at the
25 metatarsophalangeal joint. Symptoms include pain, sudden loss of active flexion at the
26 interphalangeal joint, and weakness in flexion at the metatarsophalangeal joint. Chronic,
27 complete, spontaneous rupture has also been reported in athletes. Treatment in athletes
28 consists of surgical repair. If the tendon has retracted proximally, and cannot be
29 reapproximated, it may be sutured using a tendon graft.

30
31 In some patients with low demands on the foot, complaints of a hyperextended distal
32 phalanx rubbing on top of the shoe persist after repair with tendon graft. Arthrodesis of the
33 hallux interphalangeal joint in 20 degrees of flexion may provide relief of symptoms
34 without altering performance.

35
36 In a review of treatments in the management of chronic Achilles tendon rupture, flexor
37 hallucis longus (FHL) tendon transfer was widely reported as treatment in twenty-two
38 studies. Other tendon transfer methods, such as semitendinosus tendon transfer (ST
39 transfer), peroneus brevis tendon transfer (PB transfer) and hamstring tendon transfer, were
40 reported in seven, six and two studies, respectively. A total of ten studies used
41 gastrocnemius flaps with no augmentation, whilst six studies describe additional flexor
42 hallucis longus augmentation. Direct repair techniques such as V–Y and Z plasty were

1 reported both as stand-alone techniques and combined with a synthetic acellular human
2 dermal tissue matrix graft jacket (Arshad et al., 2021).

3
4 With respect to acute Achilles tendon rupture, the American College of Foot and Ankle
5 Surgeons (2021) created a clinical consensus statement to address selected aspects of care
6 of the acute Achilles tendon injury. The panel reached consensus on the statement that
7 patients with increased risk factors for postoperative complications (diabetes, obesity,
8 cigarette smoking) have special considerations with regard to deciding operative versus
9 nonoperative management of the acute Achilles tendon rupture. The panel also agreed that
10 acute partial Achilles tendon ruptures should be treated nonoperatively and that early
11 weightbearing and progressive physical therapy should be used after repair or at initiation
12 of nonoperative management.

13 14 **PRACTITIONER SCOPE AND TRAINING**

15 Practitioners should practice only in the areas in which they are competent based on their
16 education, training, and experience. Levels of education, experience, and proficiency may
17 vary among individual practitioners. It is ethically and legally incumbent on a practitioner
18 to determine where they have the knowledge and skills necessary to perform such services
19 and whether the services are within their scope of practice.

20
21 It is best practice for the practitioner to appropriately render services to a member only if
22 they are trained, equally skilled, and adequately competent to deliver a service compared
23 to others trained to perform the same procedure. If the service would be most competently
24 delivered by another health care practitioner who has more skill and training, it would be
25 best practice to refer the member to the more expert practitioner.

26
27 Best practice can be defined as a clinical, scientific, or professional technique, method, or
28 process that is typically evidence-based and consensus driven and is recognized by a
29 majority of professionals in a particular field as more effective at delivering a particular
30 outcome than any other practice (Joint Commission International Accreditation Standards
31 for Hospitals, 2020).

32
33 Depending on the practitioner's scope of practice, training, and experience, a member's
34 condition and/or symptoms during examination or the course of treatment may indicate the
35 need for referral to another practitioner or even emergency care. In such cases it is prudent
36 for the practitioner to refer the member for appropriate co-management (e.g., to their
37 primary care physician) or if immediate emergency care is warranted, to contact 911 as
38 appropriate. See the *Managing Medical Emergencies (CPG 159 – S)* clinical practice
39 guideline for information.

1 **References**

- 2 American College of Ankle and Foot Surgeons (ACFAS) Position statement on cosmetic
3 surgery (2020). Retrieved on April 12, 2024 from: [https://www.acfas.org/policy-](https://www.acfas.org/policy-advocacy/policy-position-statements/acfas-position-statement-on-cosmetic-surgery)
4 [advocacy/policy-position-statements/acfas-position-statement-on-cosmetic-surgery](https://www.acfas.org/policy-advocacy/policy-position-statements/acfas-position-statement-on-cosmetic-surgery)
5
- 6 American Medical Association. (current year). Current Procedural Terminology (CPT)
7 Current year (rev. ed.). Chicago: AMA
8
- 9 American Medical Association. (current year). ICD-10-CM. American Medical
10 Association
11
- 12 Arshad, Z., Lau, E. J. S., Leow, S. H., & Bhatia, M. (2021). Management of chronic
13 Achilles ruptures: a scoping review. *International orthopaedics*, 45(10), 2543-2559
14
- 15 Baxter, D. E., Porter, D. A., & Schon, L. (2008). *Baxter's the Foot and Ankle in Sport:*
16 Mosby Elsevier
17
- 18 Joint Commission International. (2020). Joint Commission International Accreditation
19 Standards for Hospitals (7th ed.): Joint Commission Resources
20
- 21 Naldo, J., Agnew, P., Brucato, M., Dayton, P., & Shane, A. (2021). ACFAS Clinical
22 Consensus Statement: Acute Achilles Tendon Pathology. *The Journal of foot and ankle*
23 *surgery: official publication of the American College of Foot and Ankle Surgeons*,
24 60(1), 93–101. <https://doi.org/10.1053/j.jfas.2020.02.006>
25
- 26 Sammarco, V. J., Sammarco, G. J., Henning, C., & Chaim, S. (2009). Surgical Repair of
27 Acute and Chronic Tibialis Anterior Tendon Ruptures. *The Journal of Bone and Joint*
28 *Surgery Am*, 91(2), 325-332. doi: 10.2106/jbjs.g.01386