

1 **Clinical Practice Guideline: Unna Boot**
 2
 3 **Date of Implementation: February 18, 2016**
 4
 5 **Product: Specialty**
 6
 7

8 **GUIDELINES**

9 American Specialty Health – Specialty (ASH) considers services consisting of CPT Code
 10 29580 - Unna boot bandage - to be medically necessary when used for a primary diagnosis
 11 from the table below with the following indications:

- 12 • To treat venous vascular insufficiency;
- 13 • For the treatment of ulcers with and without inflammation of the lower extremities
 14 which are caused by increased venous pressure, venous insufficiency or capillary
 15 dysfunction; and
- 16 • For the management of sprains, strains, dislocations and minor fractures.

17
 18 Unna boot application is *not* indicated for use with ulcers resulting from arterial disease or
 19 diabetes.

20
 21 **Other considerations:**

- 22 A. This code is to be used for management of sprains, strains, dislocations and minor
 23 fractures requiring immobilization as part of the treatment plan. Although an Unna
 24 boot is frequently used to help with reduction of lower extremity edema, it is
 25 considered a part of the dressing and should not be billed separately in wound
 26 debridement.
- 27 B. An Unna boot, as well as other dressings such as Kling, Profore, etc., may be used
 28 as a dressing as an adjunct to wound debridement and will be covered as a supply;
 29 however, should not be billed with CPT 29580.
- 30 C. The method of application (primary or secondary dressing) will be left to the
 31 discretion of the provider.
- 32 D. An Unna Boot is considered to be a compression dressing, not a cast. Therefore,
 33 the supply for Unna boot is included in the payment of the procedure and not paid
 34 separately.

35
 36 **CPT Codes and Descriptions applicable when medically necessary per the criteria**
 37 **listed above**

CPT Code	Description
29580	Strapping; Unna boot

1 **ICD-10 Codes and Descriptions applicable when medically necessary per the criteria**
 2 **listed above**

ICD-10 Code	Description
Information listed in brackets below has been added for clarification purposes	
I80.00 - I80.299	Phlebitis and thrombophlebitis of superficial vessels of unspecified lower extremity - Phlebitis and thrombophlebitis of other deep vessels of unspecified lower extremity
I83.002 - I83.009, I83.012 - I83.019, I83.022 - I83.029, I83.10 - I83.12, I83.202 - I83.209, I83.212 - I83.219, I83.222 - I83.229, I83.811 - I83.899	Varicose veins of lower extremities
I87.011 - I87.099	Postthrombotic syndrome of lower extremity
I87.2	Venous insufficiency (chronic) (peripheral)
I87.311 - I87.319	Chronic venous hypertension (idiopathic) with ulcer of lower extremity
I87.331 - I87.339	Chronic venous hypertension (idiopathic) with ulcer and inflammation of lower extremity
L89.500 – L89.629, L89.90 - L89.95	Pressure ulcer of lower extremity
L97.201 - L97.929, L98.491 – L98.499	Non-pressure chronic ulcer of lower extremity
M25.471 - M25.476	Effusion, ankle and foot
M66.271 – M66.279, M66.28 – M66.29, M66.361 - M66.379, M66.38 – M66.39	Spontaneous rupture of tendons, ankle and foot
M84.361A - M84.379S	Stress fracture, lower extremity
R60.0 – R60.9	Localized, generalized, or unspecified edema

ICD-10 Code	Description
S81.801A - S81.859S, S86.021A - S86.929S, S91.001A - S91.359S, S96.021A - S96.899S	Open wound of lower extremity
S82.301A - S82.309S, S82.391A - S82.399S, S82.51XA - S82.66XS, S82.841A - S82.856S, S82.871A - S82.899S, S89.101A - S89.199S, S89.301A - S89.399S	Fracture of tibia and fibula
S86.011A - S86.019S, S93.401A - S93.699S, S96.011A - S96.0119S, S96.111A - S96.119S, S96.211A - S96.219S,	Strains and sprains of ankle and foot
S90.00XA - S90.32XS	Contusion of foot and ankle
S92.001A - S92.356S, S92.401A - S92.599S, S92.901A - S92.919S	Fracture of talus, calcaneus, metatarsal, foot, and toe(s)
S93.101A - S93.106S	Subluxation or dislocation of toe(s)
S93.111A - S93.139S, S93.311A - S93.326S	Subluxation or dislocation of interphalangeal, tarsal, and tarsometatarsal joint of foot
S93.301A - S93.306S, 393.331A - 393.336S, S93.01XA - S93.06XS	Subluxation or dislocation of foot and ankle
S96.811A - S96.819S, S96.911A - S96.919S	
T81.89XA - T81.89XS	Other complications of procedures, not elsewhere classified [i.e., non-healing surgical wound]

1
2
3
4
5
6
7
8

BACKGROUND

Unna Boots are named after a German dermatologist, Paul Gerson Unna. The Unna Boot is a commercially prepared, medically impregnated compression support dressing, usually made of cotton, which has a zinc oxide paste applied uniformly to the entire bandage. The zinc oxide paste in the Unna Boot helps ease skin irritation and keeps the area moist. The zinc promotes healing within wound sites, making it useful for burns and ulcers. Zinc oxide paste is superior to gelatins used in other dressings, because it does not harden or cake.

1 Calamine lotion or glycerin may also be used. The bandage is applied to the leg from the
2 toe to the knee by overlapping wraps of impregnated gauze. The Unna boot forms a semi-
3 rigid soft cast which should be left in place for 4 to 7 days. The Unna boot bandage restricts
4 the volume of the leg, controls edema, and encourages more normal prograde venous blood
5 flow with reduction in the subcutaneous blood pressure. The net effect is improved healing
6 of venous stasis ulcers of the lower extremities. Impregnated gauze wraps are commonly
7 referred to as a "soft cast" that are used to deliver sustained, graduated compression for the
8 management of lower extremity edema and ulcerations associated with venous
9 insufficiency. Unna Boots provide between 20-30 mmHg in pressure, making them useful
10 in a variety of wounds. It supports vascular problems, helps with healing leg ulcers,
11 swelling or lymphedema by giving compression to the areas that are wrapped. In general,
12 Unna Boots are used to treat wounds with light to moderate drainage and sometimes used
13 with hydrogel dressings. Unna Boots are more commonly used for patients who are active
14 and can move on their own, as opposed to patients who are confined to a wheelchair or
15 bed.

16 **PRACTITIONER SCOPE AND TRAINING**

17 Practitioners should practice only in the areas in which they are competent based on their
18 education, training and experience. Levels of education, experience, and proficiency may
19 vary among individual practitioners. It is ethically and legally incumbent on a practitioner
20 to determine where they have the knowledge and skills necessary to perform such services
21 and whether the services are within their scope of practice.
22

23
24 It is best practice for the practitioner to appropriately render services to a member only if
25 they are trained, equally skilled, and adequately competent to deliver a service compared
26 to others trained to perform the same procedure. If the service would be most competently
27 delivered by another health care practitioner who has more skill and training, it would be
28 best practice to refer the member to the more expert practitioner.
29

30 Best practice can be defined as a clinical, scientific, or professional technique, method, or
31 process that is typically evidence-based and consensus driven and is recognized by a
32 majority of professionals in a particular field as more effective at delivering a particular
33 outcome than any other practice (Joint Commission International Accreditation Standards
34 for Hospitals, 2020).
35

36 Depending on the practitioner's scope of practice, training, and experience, a member's
37 condition and/or symptoms during examination or the course of treatment may indicate the
38 need for referral to another practitioner or even emergency care. In such cases it is prudent
39 for the practitioner to refer the member for appropriate co-management (e.g., to their
40 primary care physician) or if immediate emergency care is warranted, to contact 911 as
41 appropriate. See the *Managing Medical Emergencies (CPG 159 – S)* clinical practice
42 guideline for information.

1 **References**

2 American Medical Association. (current year). Current Procedural Terminology (CPT)
3 Current year (rev. ed.). Chicago: AMA.

4
5 American Medical Association. (current year). ICD-10-CM. American Medical
6 Association.

7
8 Centers for Medicare and Medicaid Services. Local Coverage Article: Billing and Coding:
9 Wound Care (A53001). Retrieved on February 16, 2023 from
10 [https://www.cms.gov/medicare-coverage-](https://www.cms.gov/medicare-coverage-database/view/article.aspx?articleid=53001&ver=36&LCDId=35125&Date=12%2f01%2f2017&DocID=L35125)
11 [database/view/article.aspx?articleid=53001&ver=36&LCDId=35125&Date=12%2f01](https://www.cms.gov/medicare-coverage-database/view/article.aspx?articleid=53001&ver=36&LCDId=35125&Date=12%2f01%2f2017&DocID=L35125)
12 [%2f2017&DocID=L35125](https://www.cms.gov/medicare-coverage-database/view/article.aspx?articleid=53001&ver=36&LCDId=35125&Date=12%2f01%2f2017&DocID=L35125)

13
14 Centers for Medicare and Medicaid Services. Local Coverage Determination (LCD):
15 Wound Care (L35125). Retrieved on February 16, 2023 from
16 [https://www.cms.gov/medicare-coverage-database/details/lcd-](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=35125&ver=76)
17 [details.aspx?LCDId=35125&ver=76](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=35125&ver=76)

18
19 Joint Commission International. (2020). Joint Commission International Accreditation
20 Standards for Hospitals (7th ed.): Joint Commission Resources.