

Clinical Practice Guideline: **Cultural Competency in Pain Management and Musculoskeletal Care**

Date of Implementation: **July 17, 2025**

Effective Date: **July 23, 2026**

Product: **Specialty**

Related Policies:

QM 36: Corporate Commitment to Health Equity
 CPG 12: Medical Necessity Decision Assist Guideline for Rehabilitative Care
 CPG 110: Medical Record Maintenance and Documentation Practices
 CPG 111: Patient Assessments: Medical Necessity Decision Assist Guideline for Evaluations, Re-evaluations, and Consultations
 CPG 135: Physical Therapy Medical Policy/Guideline
 CPG 155: Occupational Therapy Medical Policy/Guideline
 CPG 166: Speech-Language Pathology/Speech Therapy Guidelines
 CPG 167: Therapeutic Massage Medical Policy/Guideline
 CPG 264: Acupuncture Services Medical Policy/Guideline
 CPG 278: Chiropractic Services Medical Policy/Guideline
 CPG 305: Virtual Physical and Occupational Therapy Services

INTRODUCTION

American Specialty Health (ASH) clinical programs are committed to improving the equitable and culturally conscious delivery of healthcare administrative and clinical services.

Health equity especially remains a critical concern in pain management. Pain is a complex experience influenced by various biological, psychological, and social factors. Health-related social needs, including stable housing/utilities, transportation, food security, employment status, and personal safety, can impact a patient's pain experience and treatment outcomes. Additionally, cultural perceptions significantly influence how individuals experience, articulate, and manage pain. Understanding and addressing these cultural differences are essential for equitable and effective pain management. By adopting a holistic and culturally sensitive approach, healthcare providers can help patients receive appropriate care and achieve positive health outcomes. It is incumbent on health care providers to conduct comprehensive pain assessments that consider these factors and develop personalized treatment plans that address both medical and social needs.

DESCRIPTION/BACKGROUND

Cultural competence in pain management and musculoskeletal care is essential for equitable and effective healthcare. Providers often encounter patients with diverse social

and cultural circumstances, including low health literacy levels, disabilities, varied ethnic backgrounds, trauma, and diverse gender identities with which the provider may not be familiar. Understanding and addressing these factors is crucial for achieving patient-centered care and optimal outcomes.

Large factors such as politics, history, economics, societal structure, geography, and culturally accepted biased policies govern whether people have access to education, housing, employment and health care. In turn, individual behaviors, opportunities, social status, health risk factors, the experience of pain, and health outcomes are influenced.

Key Definitions

Patient-Centered: Patient-centered care involves patients in their care decisions, including being mindful of their beliefs, values, and preferences.

Holistic Care: Holistic Care recognizes patients as complete individuals, encompassing various factors affecting their health and well-being. This can include cultural, familial, ethnic, health equity, environmental, spiritual, and other issues important to the person.

Social Determinants of Health (SDOH): Social Determinants of Health (SDOH) are the conditions in the environments where people are born, live, learn, work, play, worship, and age that affect a wide range of health, functioning, and quality-of-life outcomes and risks. There are five domains: 1) Economic stability, 2) Education access and quality, 3) Health care access and quality, 4) Neighborhood and built environment, and 5) Social and community context (U.S. Department of Health and Human Services, n.d.).

Health-Related Social Needs (HRSN): Health-related social needs (HRSN) are social and economic factors that impact individuals' health and well-being. These needs, such as financial instability, limited access to healthcare, and transportation issues, can lead to poorer health outcomes and increased healthcare utilization.

Health Equity: The CDC defines health equity as “the state in which everyone has a fair and just opportunity to attain their highest health.” Factors that can impede health equity include, but are not limited to, race, ethnicity, disability, sexual orientation, gender identity, socioeconomic status, geography, and preferred language. SDOH influence health equity and may impact health care access, patient presentation, clinical evaluations, treatment planning, and patient outcomes. These factors may, in turn, influence medical necessity considerations.

Health Equity Factors in Medical Necessity Review: Health Equity factors may be barriers to clinical progress documented during review of provider submissions for medical necessity. If the clinical quality evaluator notes a related health equity factor, they may communicate with the specialty provider regarding the patient’s situation and any possible

relationship to medical necessity. Standardized referral recommendations or resources for assisting with the patient's health-related needs may also be provided. The clinical quality evaluator will use the current Health and Safety Investigation Team (HSIT) guidelines if the clinical quality evaluator notes a related health and safety issue.

Cultural Competence: Cultural competence is understanding the importance of social and cultural influences on patients' health beliefs and behaviors. Providers should consider and adjust communication styles to accommodate the beliefs, values, and preferences of others. This includes cultural preferences like directness or indirectness and being mindful of body language, eye contact, and personal space, as these may have different meanings across cultures. Healthcare providers should acknowledge and address their biases about a patient's pain experience. Research indicates that some professionals hold misconceptions about pain tolerance in different racial groups, ethnicities, and sexes, leading to disparities in care. Ongoing education and strategies to combat these biases can promote equitable care for all patients.

Explicit Bias: Explicit bias refers to preferences, beliefs, and attitudes that individuals are consciously aware of and are intentional. They can often be clearly identified and communicated.

Implicit Bias: Implicit bias refers to unconscious mental processes that lead to automatic associations and reactions without intention. These biases, which include favorable or unfavorable evaluations of groups of people (stereotypes), significantly impact decision-making despite a lack of awareness of them.

PROVIDER STRATEGIES

Providers must recognize and integrate social and cultural factors that influence health beliefs and behaviors. Strategies providers can use include adapting communication styles, respecting diverse expressions of pain, and addressing any biases affecting assessment and treatment. Embracing these approaches can help build trust, improve treatment adherence, and lead to better health outcomes.

One framework for strategically managing pain is the "5 A's" noted in Walker et al. (2026). Five key actions for social care integration:

- Awareness: Identify individuals' social risks (e.g., limited transportation to pain clinic).
- Adjustment: Tailor treatment plans in response to barriers (e.g., offer telehealth when transport is an issue).

- Assistance: Link individuals to relevant resources (e.g. connections with community services and other healthcare professionals to coordinate care (e.g., coordinate with social workers to ensure transport options are known and accessible for all individuals).
- Advocacy: Use clinical insights to highlight when structures and systems create barriers to equitable care. Push for systems that make equitable care possible (e.g., work to promote policies that change the transportation infrastructure within the community).

Managing Explicit and Implicit Bias: Managing bias involves self-awareness, education, institutional assessments, and proactive strategies.

To address implicit bias, individuals should self-reflect using standardized assessments such as the Implicit Association Test (IAT) and ongoing self-analysis. Several IATs are available through Project Implicit at <https://implicit.harvard.edu/implicit/takeatest.html>. Consider the perspectives of those stereotyped by the provider's implicit bias through self-study, group educational activities, or direct interpersonal interaction. Implicit biases can also operate in institutions where common practices can be biased without the institution or its healthcare workers being aware. Implicit bias in institutions can be evaluated and altered through shared educational opportunities and changes in behaviors and policies. Proactively seeing patients as individuals rather than as a compilation of group affiliations can reduce the effects of implicit bias in clinical activities.

Management of explicit bias requires full evaluation and identification of biases as well as education on their impact, and implementation and enforcement of policies to effect change. Creating an environment where open discussions and awareness of biases and their impacts is crucial.

Examples of Explicit and Implicit Bias: Examples of both explicit and implicit biases can involve sex or gender, each affecting pain perception and expression differently. Studies have shown that women may report higher pain levels and different pain experiences compared to men. Women are more likely to suffer with chronic pain conditions such as fibromyalgia and temporomandibular joint disorders while men have a higher incidence of cluster headaches. In a review of the literature, Grinberg and Sela (2025) demonstrated that despite reporting higher levels of pain, women frequently received inadequate analgesia; Unconscious biases, a lack of specific protocols, and cultural stereotypes were contributing factors. Harkins et al. (2025) demonstrated that women with marginalized social identities have poorer pregnancy outcomes and receive inferior quality pain management during childbirth. Intersex people may also experience pain differently from other sexes especially due to hormonal and anatomic variations.

Preliminary research shows higher rates of pain in transgender and gender diverse individuals, especially if they are undergoing hormonal therapies. Gender-diverse youth are at increased risk for chronic pain conditions. Providers should be aware of these differences in people's pain experiences and avoid gender/sex biases in pain assessment and treatment. Explicit biases in institutions can lead to reduced access to pain management based on gender/sex and result in poorer health outcomes. Health care institutions that do not encompass the needs of sex and gender diversity can influence the development of diagnoses and treatment plans that are mismatched to the patients and therefore less effective.

Tailored Treatment Plans: Personalizing treatment strategies to the individual's specific goals and preferences is crucial for effective pain management and musculoskeletal care. For example, integrating the patient's traditional healing practices can enhance patient engagement, adherence to care plans, and health outcomes. Providers may consider engaging with the surrounding communities where their patients live to better understand local cultural contexts and build relationships with community and cultural leaders. Baslam et al. (2025) completed a scoping review of cultural tailoring of pain management in 38 studies with 27 unique pain management intervention approaches. Seventy percent of the approaches addressed racial and ethnic minorities; Educational interventions were the most common with content adaptation to culturally relevant language and gender considerations. The study authors suggested future broadening to include lower-income countries, more digital involvement and more monitoring of near and long-term outcomes. Another individual factor that may require more tailored attention is the person's age. Keep in mind that people of different ages and generations have varying perceptions of health and illness, and how they are expected to interact with health care providers.

Outcome Measures: Patient-reported outcome measures (PROMs) provide information on how a patient is functioning, their symptoms and related emotions, and quality of life. This information aids the provider in tailoring the treatment plan. Tools that have been validated in the specific cultural groups served should be used when available, especially if translating into another language or adapting for a patient's cultural background.

Clinicians can offer culturally appropriate information about pain management options and involve patients in decision-making. Regularly soliciting patient feedback regarding pain management experiences can allow for adjustments to care plans as needed. The use of patient-reported functional outcome measures (PROMs) helps monitor progress in functions that are important to the patient during pain management.

Communication and Patient Education: Practitioners should also be aware of and respect different cultural interpretations and expressions of pain. Recognizing that cultural backgrounds influence pain perception can enhance patient care. Some cultures may view discussing pain as necessary, while others may consider it a sign of vulnerability.

Healthcare providers can better communicate and understand patients' pain experiences by learning the unique terminologies and metaphors used across cultures.

Improving cultural competence in healthcare fosters better communication between providers and patients, creating a trusting environment. Enhancing communication includes assessing barriers to communication such as language, hearing or vision impairment, and health literacy. Examples of strategies might be providing interpreters and health materials in preferred languages.

Factors that may impact communication and patient education include:

- **Literacy and Numeracy:** Literacy is typically defined as the ability to read and write. It encompasses the skills necessary to understand, evaluate, and use written text in order to engage effectively in society (CDC, 2024). Numeracy refers to the ability to access, use, interpret, and communicate mathematical information and ideas (CDC, 2024). Health literacy and literacy are related, but not the same. It is estimated that about 22% of people in the United States speak a language other than English. Children and families who speak languages other than English have poorer pain outcomes compared with English-speaking children and families. Issues that influence outcomes include clinician bias, interpersonal miscommunications, misunderstandings of culture concepts of pain, and lack of access to interpretation services (Lim et al., 2025).

- **Health Literacy:** Health literacy is defined as an individual's ability to access, understand, and process health information in order to make appropriate health decisions. The World Health Organization states that "health literacy is a stronger predictor of an individual's health status than income, employment status, education level, and racial or ethnic group." Many patients may have limited health literacy, making it difficult for them to understand medical terminology and treatment plans. Even people with high literacy and numeracy may have limited health literacy. Providers should use plain language, visual aids, and teach-back methods to ensure patients comprehend their care instructions. This approach helps in building trust and improving adherence to treatment. Debonne et al. (2025), demonstrated that out of 23 studies and 41 patient education pieces, 95% were in English, less than 20% had any readability assessments, only one piece had all of the information required, and just 8 studies included patient representative involvement.

E-Health Literacy is defined by the World Health Organization (WHO) as "the ability to seek, find, understand, and appraise health information from electronic sources and apply the knowledge gained to address or solve a health problem." People's access to electronic sources of information and their ability to use them

effectively and safely can vary widely. This includes cell phones, computers, and patient portals. Providers can evaluate what the patient is able to access and use, have resources for assistance, and alternate methods for communication and information sharing.

- **Hearing Impairments:** Patients with hearing impairments may struggle to communicate their pain levels and symptoms effectively. They may also have difficulty hearing questions, directions, and participating in shared decision-making. Age is one of the most important risk factors for hearing loss, and as the population ages, the number of people with hearing loss will continue to increase (Haile et al., 2021). Hearing impairments and aging are common sources of stigma causing patients to try to hide their issues and thus reduce their access to pain management and other care. Providers should use clear, simple language and consider using sign language interpreters or written communication tools as appropriate to facilitate better understanding. Ensuring that patients can express their pain accurately is vital for appropriate pain management.
- **Vision Impairments:** Vision impairment may impact communication with the provider in several ways. It may make it difficult for the patient to read non-verbal cues from the provider. Vision impairment may also make it difficult for the patient to describe their pain, especially if the provider is using visual analog scales or written questionnaires. Appropriate educational material may be limited. Consider having consent forms and other handouts or patient education material available in large black and white print, electronic copies, braille, or as an audio version.

Screening for Social Determinants of Health:

Providers may screen patients for social needs that could be related to their health in general, as well as pain management options and outcomes. There are several screening tools from multiple agencies. Providers can choose the tools that most relate to their patient populations. Below are a few examples:

Health Leads Social Needs Screening Toolkit

https://healthleadsusa.org/wp-content/uploads/2023/05/Screening_Toolkit_2018.pdf

CMS Accountable Health Communities Health-Related Social Needs Screening Tool

<https://www.cms.gov/priorities/innovation/files/worksheets/ahcm-screeningtool.pdf>

PRAPARE Protocol for Responding to and Assessing Patients' Assets, Risks and Experiences

<https://prapare.org/prapare-toolkit/>

1 ***American Academy of Family Physicians Social Needs Screening Tool***

2 https://www.aafp.org/dam/AAFP/documents/patient_care/everyone_project/hops19-physician-form-sdoh.pdf

5 ***WellRX Questionnaire***

6 https://cdn.prod.website-files.com/63fd0d00f2ad370c6ec29450/68e4bac416d25e603913d06e_WellRX-Questionnaire-Fillable.pdf

10 **MEDICAL NECESSITY REVIEW**

11 During the review of materials submitted for medical necessity review, clinical quality
12 evaluators may become aware of barriers to care, clinical progress, and/or outcomes related
13 to social determinants of health (SDOH). These SDOH may include but are not limited to
14 economic instabilities, healthcare access, and social/community contexts. If the clinical
15 quality evaluator notes a related SDOH, they may communicate with the specialty provider
16 regarding the patient's situation and any possible relationship to medical necessity and
17 provide standardized referral recommendations or resources for assisting with the patient's
18 health-related social needs. If the clinical quality evaluator notes a related health and safety
19 issue, they will utilize the current HSIT guidelines

21 A comprehensive pain assessment by the provider submitting for medical necessity review
22 involves understanding patients' unique experiences. Considering factors like medical
23 history, socioeconomic status, health literacy, and cultural influences allows for more
24 accurate diagnoses and treatment plans. Medical necessity reviews involve evaluating the
25 appropriateness of pain management interventions based on clinical guidelines and patient-
26 specific factors. Providers are informed about the criteria used in these reviews so that
27 relevant patient information is submitted to demonstrate medical necessity.

29 **PRACTITIONER SCOPE AND TRAINING**

30 Practitioners should practice only in the areas in which they are competent based on their
31 education, training, and experience. Levels of education, experience, and proficiency may
32 vary among individual practitioners. It is ethically and legally incumbent on a practitioner
33 to determine where they have the knowledge and skills necessary to perform such services
34 and whether the services are within their scope of practice.

36 It is best practice for the practitioner to appropriately render services to a patient only if
37 they are trained to competency, equally skilled, and adequately competent to deliver a
38 service compared to others trained to perform the same procedure. If the service would be
39 most competently delivered by another health care practitioner who has more skill and
40 training, it would be best practice to refer the patient to the more expert practitioner.

Best practice can be defined as a clinical, scientific, or professional technique, method, or process that is typically evidence-based and consensus driven and is recognized by a majority of professionals in a particular field as more effective at delivering a particular outcome than any other practice (Joint Commission International Accreditation Standards for Hospitals, 2020).

Depending on the practitioner's scope of practice, training, and experience, a patient's condition and/or symptoms during examination or the course of treatment may indicate the need for referral to another practitioner or even emergency care. In such cases it is essential for the practitioner to refer the patient to appropriate co-management (e.g., to their primary care physician) or if immediate emergency care is warranted, to contact 911 as appropriate. See the *Managing Medical Emergencies (CPG 159 – S)* clinical practice guideline for information.

Helpful resources for patients:

FindHelp.org

- Free or reduced-cost resources like food, housing, financial assistance, healthcare and more.
- Just enter zip code to find resources.

211.org or Call 211

- Connect you to expert, caring help

988lifeline.org or Call/ text 988

- Provides 24/7, free support for people in distress, prevention and crisis resources for you or your loved ones, and best practices for professionals in the United States

Call/ text 911

- National police, fire, or emergency medical assistance from any phone in any location

USPainFoundation.org

- *References and support for people with pain*

References

Basalem N, Astek AA, Sroge RA, Ali SM, Gangannagaripalli J, Stanmore E, van der Veer SN. Cultural tailoring of pain management approaches: a scoping review. *Int J Equity Health*. Published 2025 Dec 20. doi:10.1186/s12939-025-02743-5.

Betancourt JR, Green AR, Carrillo JE, Ananeh-Firempong O 2nd. Defining cultural competence: a practical framework for addressing racial/ethnic disparities in health and health care. *Public Health Rep*. 2003;118(4):293-302.

Brady B, Veljanova I, Andary T, Southwell T, Chipchase L. Recognising ethnocultural diversity in chronic pain assessment: validation of the Pictorial Representation of

1 Illness and Self Measure (PRISM) for use with culturally diverse communities. *Health*
 2 *Qual Life Outcomes*. 2019;17(1):56. doi:10.1186/s12955-019-1126-9

4 Bull C, Teede H, Watson D, Callander EJ. Selecting and Implementing Patient-Reported
 5 Outcome and Experience Measures to Assess Health System Performance. *JAMA*
 6 *Health Forum*. 2022;3(4):e220326. Published 2022 Apr 1.
 7 doi:10.1001/jamahealthforum.2022.0326

9 Centers for Disease Control and Prevention. Understanding Literacy &
 10 Numeracy. <https://www.cdc.gov/healthliteracy/learn/understandingliteracy.html>.
 11 Published October 16, 2024. Accessed May 6, 2025.

13 Debonne C, Houdart A, Cachinho C, Ouvrier-Neyret A, Gérard T, Vaillant V, Tousignant-
 14 Laflamme Y, Gagnon MP, Sasseville M, Décary S, Naye F. Accessible Patient
 15 Education Materials for Low Back Pain Rarely Meet People's Information Needs: A
 16 Scoping Review. *Musculoskeletal Care*. 2025 Jun;23(2):e70130.
 17 doi:10.1002/msc.70130.

19 FitzGerald C, Hurst S. Implicit bias in healthcare professionals: a systematic review. *BMC*
 20 *Med Ethics*. 2017;18(1):19. Published 2017 Mar 1. doi:10.1186/s12910-017-0179-8

22 Grinberg K, Sela Y. A Literature Review on Pain Management in Women During Medical
 23 Procedures: Gaps, Challenges, and Recommendations. *Medicina (Kaunas)*. 2025 Jul
 24 26;61(8):1352. doi:10.3390/medicina61081352.

26 Haile LM, Kamenov K, Briant PS, et al. Hearing loss prevalence and years lived with
 27 disability, 1990–2019: findings from the global burden of disease study 2019. *Lancet*.
 28 2021;397(10278):996–1009. doi: 10.1016/S0140-6736(21)00516-X.

30 Halpin S, Tarrant N, de la Cruz Y. Social Drivers vs. Social Determinants: Using Clear
 31 Terms. National Association of Community Health
 32 Centers. [https://www.nachc.org/social-drivers-vs-social-determinants-using-clear-](https://www.nachc.org/social-drivers-vs-social-determinants-using-clear-terms/)
 33 [terms/](https://www.nachc.org/social-drivers-vs-social-determinants-using-clear-terms/). Published January 23, 2023. Accessed May 6, 2025.

35 Han S, Letra A, Alessandri-Bonetti A, Thomas D, Kapos FP, Sangalli L. Sex differences
 36 in pain in 2 large and diverse US databases. *Pain Rep*. 2025;11(1):e1371. Published
 37 2025 Dec 12. doi:10.1097/PR9.0000000000001371

39 Harkins SE, Hazi AK, Guglielminotti J, Landau R, Barcelona V. Discrimination, racism,
 40 and bias in childbirth pain management in the United States: a scoping review and
 41 directions for research and clinical care. *Int J Obstet Anesth*. 2025;63:104379.
 42 doi:10.1016/j.ijoa.2025.104379.

- 1 Joint Commission International. Joint Commission International Accreditation Standards
2 for Hospitals. 7th ed. Oak Brook, IL: Joint Commission Resources; 2020.
- 3
- 4 Keough, E, Nwasor EO, Boerner KE. Overview of sex and gender differences in human
5 pain. International Association for the Study of Pain. Published June 28, 2024.
6 Accessed May 7, 2026. <https://www.iasp-pain.org/resources/fact-sheets/156455/>
- 7
- 8 Lim PS, Fortier MA, Kain ZN. Pain disparities attributed to linguistic minoritization in
9 health care settings. *J Pain*. 2025;37S:104688. doi:10.1016/j.jpain.2024.104688.
- 10
- 11 Mansutti I, Tomé-Pires C, Chiappinotto S, Palese A. Facilitating pain assessment and
12 communication in people with deafness: a systematic review. *BMC Public Health*.
13 2023;23(1):1594.
- 14
- 15 Mohmand S, Chams S. Cultural considerations in the management of chronic pain. *CAND*
16 *J*. 2021;28(4):11-13. <https://doi.org/10.54434/candj.93>
- 17
- 18 Mulvey MR, Paley CA. The challenges and opportunities for cancer pain management in
19 primary and community care services: a scoping review. *Pain Manag*. 2026;16(3):245-
20 256. doi:10.1080/17581869.2025.2608575.
- 21
- 22 Reis FJJ, Nijs J, Parker R, Sharma S, Wideman TH. Culture and musculoskeletal pain:
23 strategies, challenges, and future directions to develop culturally sensitive physical
24 therapy care. *Braz J Phys Ther*. 2022;26(5):100442.
- 25
- 26 Rezvani Kakhki B, Sahebi S. Mapping global research on labor and postpartum pain
27 management: a bibliometric analysis (2000-2025). *Int J Equity Health*. 2026;25(1):87.
28 doi:10.1186/s12939-026-02805-2.
- 29
- 30 Schoenthaler A, Williams N. Looking beneath the surface: racial bias in the treatment and
31 management of pain. *JAMA Netw Open*. 2022;5(6):e2216281.
- 32
- 33 Stubbe DE. Practicing cultural competence and cultural humility in the care of diverse
34 patients. *Focus (Am Psychiatr Publ)*. 2020;18(1):49-51.
- 35
- 36 Truong M, Paradies Y, Priest N. Interventions to improve cultural competency in
37 healthcare: a systematic review of reviews. *BMC Health Serv Res*. 2014;14:99.
38 Published 2014 Mar 3. doi:10.1186/1472-6963-14-99
- 39
- 40 Vela MB, Erondy AI, Smith NA, Peek ME, Woodruff JN, Chin MH. Eliminating Explicit
41 and Implicit Biases in Health Care: Evidence and Research Needs. *Annu Rev Public*
42 *Health*. 2022;43:477-501. doi:10.1146/annurev-publhealth-052620-103528

- 1 Walker E, Karran EL, Jones MD, Gibbs MT, Sharma S. Social determinants of health
2 equity for people with pain: from understanding to action. *Pain Rep.* 2025;11(1):e1364.
3 Published 2025 Dec 23. doi:10.1097/PR9.0000000000001364
4
- 5 Wang ML, Jacobs O. From Awareness to Action: Pathways to Equity in Pain
6 Management. *Health Equity.* 2023;7(1):416-418. Published 2023 Aug 23.
7 doi:10.1089/heq.2023.0179
8
- 9 World Health Organization. Health literacy. [https://www.who.int/news-room/fact-](https://www.who.int/news-room/fact-sheets/detail/health-literacy)
10 [sheets/detail/health-literacy](https://www.who.int/news-room/fact-sheets/detail/health-literacy). Published August 5, 2024. Accessed May 6, 2025.