

1 **Clinical Practice Guideline: Axial/Spinal Decompression Therapy**

2

3 **Date of Implementation: July 13, 2006**

4

5 **Product: Specialty**

6

7

8 **GUIDELINES**

9 American Specialty Health – Specialty (ASH) considers nonsurgical axial/spinal
 10 decompression therapy to be unproven due to insufficient scientific evidence of efficacy in
 11 the treatment of neck, low back and related disorders. This includes any motorized
 12 mechanical traction device that is promoted as providing “decompression therapy” e.g.,
 13 VAX-D, IDD Therapy® [Intervertebral Differential Dynamics Therapy], DRS, DRX,
 14 DRX-2000, DRX-3000, DRX-5000, DRX-9000, Accu-SPINA™, Lordex Power Traction
 15 device, Mettler Traction Device [MTD 4000], Tru Trac 401, Integrity Spinal Care System
 16 Alpha-SPINA System, Dynatron DX2, Dynapro™ DX2, Spinex LDM, or any other
 17 device that claims to create spinal decompression.

18

19 The research evidence concerning nonsurgical axial/spinal decompression therapy is
 20 lacking and of low quality. Any estimate of treatment effect is uncertain, as is the clarity
 21 of risk, benefit, and burden to the patient.

22

23 There are significant burdens placed upon health plan members due to high out-of-pocket
 24 costs, time spent receiving the intervention, and the unsubstantiated/misleading marketing
 25 about the alleged proven effectiveness and safety of nonsurgical axial/spinal
 26 decompression therapy. These burdens have been recognized as significant by some
 27 professional licensing boards and state justice departments.

28

29 Similar conclusions have been reached by a broad range of health care organizations.
 30 Professionals and groups, who are proponents of nonsurgical axial/spinal decompression
 31 therapy, should pursue further investigation using experimental study designs and rigorous
 32 methodologies.

33

HCPCS/	Description
S9090	Vertebral Axial Decompression, per session; {most accurately describes services for the application of spinal decompression motorized traction devices }
Other CPT codes that have been associated with the use of nonsurgical spinal decompression therapy are:	
64722	Decompression; unspecified nerve(s) (specify) {a surgical code }
97012	Application of a modality to 1 or more areas; traction, mechanical

1 **DESCRIPTION/BACKGROUND**

2 Traction as a treatment option for low back pain and sciatica has existed for many years.
 3 Its use has progressed from continuous static traction to intermittent motorized traction.
 4 The most recent form of intermittent motorized traction is commonly referred to as
 5 axial/spinal decompression therapy. Developers and manufacturers of the equipment along
 6 with clinicians often consider it to be a unique form of traction. Proponents of nonsurgical
 7 axial/spinal decompression therapy claim it to be a safe and effective alternative to surgical
 8 interventions. Companies demonstrate intense marketing programs and claim high success
 9 rates. Axial/spinal decompression therapy is intended to create negative pressure within
 10 the spine so that as the spinal column is elongated, pressure is taken off the nerve root(s),
 11 and herniated disc material may be pulled back into place. Axial/spinal decompression
 12 therapy is generally performed using a specially designed computerized mechanical table
 13 that separates in the middle. Depending on the type of table being used, a patient is strapped
 14 in a prone or supine position to the lower part of the table using a pelvic harness and may
 15 hold handgrips at the top of the table. The table is then mechanically separated in the
 16 middle creating a distractive force to relieve pressure within the spine that may be causing
 17 pain. The amount of distractive force is tailored for each patient and usually lasts about 60
 18 seconds. Depending on the device utilized, static, intermittent, or cycled distractive force
 19 may be applied. Typical treatment protocols include 20 sessions, each lasting 30 to 40
 20 minutes. The process of distraction and relaxation is fully computerized using a
 21 programmable logic controller and is monitored by a licensed health care practitioner. The
 22 American Medical Association (AMA), Food and Drug Administration (FDA), and
 23 Centers for Medicare & Medicaid Services (CMS) all consider axial/spinal decompression
 24 therapy to be a form of traction. However, this therapy involves a special table and protocol
 25 that isn't the same as conventional or traditional traction with claims of spinal
 26 decompression.

27
 28 The tables utilized for axial/spinal decompression therapy are classified by the FDA as
 29 powered traction equipment. Examples of axial/spinal decompression therapy tables (and
 30 their manufacturers) include:

- 31 • VAX-D Table (VAX-D Manufacturing, Palm Harbor, FL)
- 32 • Decompression, Reduction, Stabilization (DRS) System (North American Medical
 33 Corporation, Atlanta, GA)
- 34 • DRX 2000 and DRX 9000 (Axiom Worldwide, Tampa, FL)
- 35 • Spina System (North American Medical Corporation, Atlanta, GA)

36
 37 Two popular units will be described here. Due to the number of available products, it would
 38 be impractical to provide information on all of them.

39 **VAX-D**

40 The manufacturer suggests that use of the VAX-D table applies distractive forces in a
 41 gradual, progressive fashion through extension of the lower end of the table. The level of
 42

1 tension is preset on a control panel and can be increased, allowing for various
 2 decompression phases and a rest phase. Various decompression phases allow alternating
 3 cycles of distraction and relaxation. Typically, a treatment cycle consists of 15 cycles of
 4 tension and relaxation. The patient lies prone on the VAX-D table. The table is split,
 5 allowing the table to slowly extend, thus decreasing load bearing in the intervertebral discs
 6 and/or intervertebral joint spaces. The VAX-D manufacturer claims specific parameters of
 7 their system make the device inherently safe. These safety features include the use of air
 8 pressure as the energy source; the ramp characteristics employed in applying the distraction
 9 tensions; the release rate of the distraction and relaxation cycles; the cycle periodicity; the
 10 upper limits on the distraction tensions; the positioning of the patient and the means of
 11 fixing the upper body; and the ability of the patient to release the handgrips if the distraction
 12 tension causes pain or discomfort. Information regarding the range and incidence of
 13 adverse effects that occur during VAX-D therapy is limited. Complications reported with
 14 VAX-D include:

- 15 • The development of a sharp burning, radiating pain during therapy
- 16 • Stress to the shoulder girdle and rotator cuff muscles
- 17 • Overstretching of the soft tissue of the back

18 **Decompression, Reduction, Stabilization (DRS) System**

19 Manufacturers recommend the DRS System for treatment of low back pain. This device
 20 uses a bed that is split into two cushions. The patient can step onto a foot pad, have a pelvic
 21 and chest harness attached, after which the patient and bed are lowered to a horizontal
 22 position. Distraction tension is applied by the pelvic harness while the patient's upper body
 23 is secured to the locked upper cushion via the chest harness. The DRS System is marketed
 24 for the treatment of low back pain associated with herniated and degenerated discs.
 25 According to the manufacturer, the DRS System applies pressures on the disc in a
 26 graduated manner, which bypasses the inherent neurological mechanisms that lead to firing
 27 of stretch receptors in the paravertebral structures. This decreased resistance to the
 28 distractive forces allows a reduction in intradiscal pressures, which promotes retraction of
 29 herniated disc material and facilitates influx of oxygen, proline, and other substrates.
 30

31 **EVIDENCE REVIEW**

32 Currently, there is not adequate scientific evidence which proves that axial/spinal
 33 decompression is an effective single intervention or adjunct to conservative therapy for
 34 back pain. In addition, axial/spinal decompression devices have not been adequately
 35 studied as alternatives to back surgery.
 36

37 Proponents of nonsurgical axial/spinal decompression therapy assert this form of traction
 38 is, however, unique for being proven able to reduce the relative pressure measured within
 39 intervertebral discs (decompression). The evidence typically cited to support this claim is
 40 from a study by Ramos, 1994. An evaluation of this study shows the conclusions are based
 41 upon data from only three subjects. This study demonstrated a number of methodological
 42

1 flaws likely to invalidate the results. These included not using a closed transducer system,
2 not taking into account temperature effects, absent hydrostatic conditions (in degenerative
3 discs), and no attempt reported to calibrate negative readings.

4
5 Regardless of the flaws, this study is not sufficient to arrive at conclusions about the
6 translation of basic science research into clinical care settings. The author (Ramos)
7 concluded additional study is needed to establish the relationship of negative intradiscal
8 pressures with clinical outcomes. The results from an early uncontrolled, retrospective
9 study (Gose et al., 1998) regarding the benefits of the VAX-D table appeared to be
10 encouraging. However, the findings need to be validated in prospective, randomized,
11 controlled clinical trials because the study was poorly designed. A subsequent randomized
12 study (Sherry et al., 2001) compared VAX-D to transcutaneous electrical nerve stimulation
13 (TENS) in the treatment of patients with chronic (> 3 months in duration) low back pain.
14 Successful outcome was defined as a 50% decrease in pain using the Visual Analog Pain
15 Scale and an improvement in the level of functioning as measured by patient-nominated
16 disability ratings. The TENS-treated group ($n=21$) reported a success rate of 0%, while the
17 group treated with VAX-D ($n=19$) showed a success rate of 68.4%. No confirmatory
18 conclusions can be drawn from this study given detailed statistics regarding the outcomes
19 for each group was not included in the analysis. Furthermore, patients were not blinded to
20 the treatment received. The Australian Medical Services Advisory Committee (MSAC,
21 2001) performed an assessment of the literature on VAX-D therapy. The Committee
22 concluded that "there is currently insufficient evidence pertaining to the effectiveness of
23 vertebral axial decompression (VAX-D) therapy..." In 2007, they requested that the
24 Agency for Healthcare Research and Quality (AHRQ) commission an evidence-based
25 technology assessment. The AHRQ report "Decompression Therapy for the Treatment of
26 Lumbosacral Pain" concluded the current evidence regarding the efficacy of axial/spinal
27 decompression therapy is too limited in quality and quantity to allow for evidence-based
28 conclusions. Adverse event reporting for axial/spinal decompression therapy was viewed
29 as infrequent. The Centers for Medicare & Medicaid Services (CMS) Technology
30 Advisory Committee did not recommend coverage of the VAX-D system because of the
31 absence of scientific data on its effectiveness.

32
33 In review of a single study of DRS therapy (Shealy and Borgmeyer, 1997), the authors
34 reported on a comparison of DRS therapy to conventional traction for both ruptured lumbar
35 discs and chronic facet arthrosis. This study suffered from three major flaws: one of the
36 authors was affiliated with the treatment center that conducted the trial; the scale used to
37 quantify the results was not clearly defined; and the study consisted of a small sample size
38 lacking clearly defined methods of randomization.

39
40 Macario and Pergolizzi (2006) conducted a systematic review of the literature to assess the
41 efficacy of nonsurgical axial/spinal decompression that is achieved with motorized traction
42 for chronic discogenic low back pain. The authors reviewed data from 10 studies between

1 1975 and 2003. Seven were randomized controlled trials of motorized traction using
2 various apparatus types, including split-tabletop, plain tabletop, and friction-free couch
3 with weights. A total of 408 individuals received placebo, and 438 individuals received
4 motorized spinal decompression. Follow-up averaged 28 weeks. None of the studies were
5 blinded, and only three had description of the randomization method. Six of the seven
6 randomized trials reported no difference with motorized spinal decompression, and one
7 study reported reduced pain but not disability. In the author's opinion, the efficacy of spinal
8 decompression achieved with motorized traction for discogenic low back remains
9 unproven. Daniel (2007) reported that there is very limited evidence in the scientific
10 literature to support the effectiveness of non-surgical axial/spinal decompression therapy.
11 One randomized controlled trial, one clinical trial, one case series and seven other papers
12 were available in the published literature for review by the author as part of an intended
13 systematic review. Due to the limited evidence a systematic review was not done, and each
14 study was reviewed individually. The author noted many of the reviewed studies utilized
15 the VAX-D unit. Furthermore, the intervention has not been compared to exercise, spinal
16 manipulation, standard medical care or other less expensive conservative treatments.

17
18 In a prospective case series study, Beattie et al. (2008) examined outcomes after an
19 intervention of a prone lumbar traction protocol using the VAX-D system. A total of 296
20 subjects with low back pain and evidence of a degenerative and/or herniated intervertebral
21 disc at one or more levels of the lumbar spine were included in this study. Patients
22 underwent an 8-week course of prone lumbar traction, using the VAX-D system, consisting
23 of five 30-minute sessions a week for four weeks, followed by one 30-min session a week
24 for four additional weeks. The numeric pain rating scale and the Roland-Morris Disability
25 Questionnaire were completed at pre-intervention, discharge (within two weeks of the last
26 visit), and at 30 days and 180 days after discharge. A total of 250 (84.4 %) subjects
27 completed the treatment protocol. On the 30-day follow-up, 247 (83.4 %) subjects were
28 available; on the 180-day follow-up, data were available for 241 (81.4 %) subjects. These
29 researchers noted significant improvements for all post-intervention outcome scores when
30 compared with pre-intervention scores ($p < 0.01$). The authors noted that causal
31 relationships between the outcomes and the intervention cannot be made. This study lacked
32 a comparison group.

33
34 Macario et al. (2008) discussed the retrospective chart audit of 100 patients with discogenic
35 low back pain (LBP) lasting more than 12 weeks treated with a 2-month course of
36 motorized spinal decompression via the DRX9000. Patients at a convenience sample of 4
37 clinics received 30-min DRX9000 sessions daily for the first 2 weeks tapering to 1
38 session/week. Treatment protocol included lumbar stretching, myofascial release, or heat
39 prior to treatment, with ice and/or muscle stimulation afterwards. Primary outcome was
40 verbal NRS 0 to 10 before and after the 8-week treatment. Of the 100 subjects, three
41 withdrew their protected health information, and three were excluded because their LBP
42 duration was less than 12 weeks. The remaining 94 subjects had diagnoses of herniated

1 disc (73% of patients), degenerative disc disease (68 %), or both (27%). Mean NRS
 2 equaled 6.05 (SD 2.3) at presentation and decreased significantly to 0.89 (SD 1.15) at end
 3 of 8-week treatment ($p < 0.0001$). Analgesic use also appeared to decrease (charts with
 4 data = 20) and activities of daily living improved (charts with data = 38). Follow-up (mean
 5 of 31 weeks) on 29/94 patients reported mean 83% LBP improvement, NRS of 1.7 (SD
 6 1.15), and satisfaction of 8.55/10 (median of 9). The authors concluded that this
 7 retrospective chart audit provides preliminary data that chronic LBP may improve with
 8 DRX9000 spinal decompression, however caution should be taken with this interpretation
 9 given it was not provided as a singular treatment. They stated that randomized double-
 10 blind trials are needed to measure the effectiveness of such systems. Schimmel et al. (2009)
 11 conducted a randomized sham-controlled trial of intervertebral axial decompression. Sixty
 12 subjects with chronic symptomatic lumbar disc degeneration or bulging disc with no
 13 radicular pain and no prior surgical treatment (dynamic stabilization, fusion, or disc
 14 replacement) were randomly assigned to a graded activity program with an Accu-SPINA
 15 device (20 traction sessions during six weeks, reaching >50% body weight), or to a graded
 16 activity program with a non-therapeutic level of traction (<10% body weight). In addition
 17 to traction, the device provided massage, heat, blue relaxing light, and music during the
 18 treatment sessions in both groups. Neither patients nor evaluators were informed about the
 19 intervention received until after the 14-week follow-up assessment, and intention-to-treat
 20 analysis was performed (93% of subjects completed follow-up). Both groups showed
 21 improvements in validated outcome measures (visual analog scores for back and leg pain,
 22 Oswestry Disability Index, and Short-Form 36), with no differences between the treatment
 23 groups. The authors reported that the added axial, intermittent, mechanical traction of IDD
 24 Therapy to a standard graded activity program has been shown not to be effective.

25
 26 Apfel et al. (2010) conducted a retrospective cohort study of adults with chronic LBP
 27 attributed to disc herniation and/or discogenic LBP who underwent a six-week treatment
 28 protocol of motorized non-surgical spinal decompression via the DRX9000. The main
 29 outcomes were changes in pain as measured on a verbal rating scale during a flexion-
 30 extension range of motion evaluation and changes in disc height as measured on CT scans.
 31 The authors identified 30 patients with lumbar disc herniation and an average duration of
 32 LBP of 12.5 weeks. During treatment, low back pain decreased from 6.2 (SD 2.2) to 1.6
 33 (2.3, $p < 0.001$) and disc height increased from 7.5 (1.7) mm to 8.8 (1.7) mm ($p < 0.001$).
 34 Increase in disc height and reduction in pain were significantly correlated ($r = 0.36$,
 35 $p = 0.044$). Reported limitations of this study are no control group and small sample size.
 36 The authors reported that a randomized controlled trial is needed to confirm the efficacy
 37 and elucidate the mechanism of this treatment modality. Choi et al. (2015) sought to
 38 identify how spinal decompression therapy and general traction therapy influence the pain,
 39 disability, and straight leg raise (SLR) ability of patients with intervertebral disc herniation.
 40 The subjects were 30 patients with chronic lumbar pain who were divided into a spinal
 41 decompression therapy group using a spinal decompression device (SDTG, $n = 15$), and a
 42 general traction therapy group (GTTG, $n = 15$). Both groups received conservative physical

1 therapy three times a week for four weeks. A comparison of the two groups found no
2 statistically significant differences. Authors concluded that spinal decompression therapy
3 and general traction therapy are effective at improving the pain, disability, and SLR of
4 patients with intervertebral disc herniation. Limitations of the study from a methodology
5 standpoint do not allow conclusions to be confirmed. Kang et al. (2016) conducted a study
6 to clarify the difference in therapeutic effects between traction and decompression
7 therapies, and their clinical therapeutic significance. For the experimental group, 15
8 subjects were randomly selected to receive decompression therapy and trunk stabilization
9 exercise. For the control group, 16 subjects were randomly selected to receive traction
10 therapy and trunk stabilization exercise. Authors concluded that decompression therapy
11 was demonstrated to be more effective clinically than conventional traction therapy as an
12 intervention method for disk disease.

13
14 Demirel et al., (2017) sought to determine whether or not non-invasive spinal
15 decompression therapy (NSDT) was effective in resorption of herniation, increasing disc
16 height in patients with lumbar disc herniation (LHNP). A total of twenty patients diagnosed
17 as LHNP and suffering from pain at least 8 weeks were enrolled to the study. Patients were
18 randomly allocated in study (SG) and control groups (CG). Both groups received
19 combination of electrotherapy, deep friction massage and stabilization exercise for fifteen
20 sessions. SG received additionally NSDT different from CG. Numeric Analog Scale,
21 Straight leg raise test, Oswestry Disability Index (ODI) were applied at baseline and after
22 treatment. Disc height and herniation thickness were measured on MRI which performed
23 at baseline and three months after therapy. Both treatments had positive effect for
24 improving pain, functional restoration and reduction in thickness of herniation. Although
25 reduction of herniation size was higher in SG than CG, no significant differences were
26 found between groups and any superiority to each other ($p > 0.05$). Given the study design,
27 the study showed that physiotherapy was helpful but that adding NSDT did not confer
28 additional benefits. Amjad et al. (2022) sought to determine the effects of non-surgical
29 spinal decompression (NSD) therapy in addition to routine physical therapy on pain,
30 lumbar range of motion (ROM), functional disability, back muscle endurance (BME), and
31 quality of life (QOL) in patients with lumbar radiculopathy. A total of sixty patients with
32 lumbar radiculopathy were randomly allocated into two groups, an experimental ($n = 30$)
33 and a control ($n = 30$) group, through a computer-generated random number table. Baseline
34 values were recorded before providing any treatment by using a visual analogue scale
35 (VAS), Urdu version of Oswestry disability index (ODI-U), modified-modified Schober's
36 test (MMST), prone isometric chest raise test, and Short Form 36-Item Survey (SF-36) for
37 measuring the pain at rest, functional disability, lumbar ROM, BME, and QOL,
38 respectively. All patients received twelve treatment sessions over 4 weeks, and then all
39 outcome measures were again recorded. By using the ANCOVA test, a statistically
40 significant ($p < 0.05$) between-group improvement was observed in VAS, ODI-U, BME,
41 lumbar ROM, role physical (RP), and bodily pain (BP) domains of SF-36, which was in
42 favour of NSD therapy group. For these outcomes, a medium to large effect size ($d = 0.61$ -

1 2.47, 95% CI: 0.09-3.14) was observed. It was concluded that a combination of non-
 2 surgical spinal decompression therapy with routine physical therapy is more effective,
 3 statistically and clinically, than routine physical therapy alone in terms of improving pain,
 4 lumbar range of motion, back muscle endurance, functional disability, and physical role
 5 domain of quality of life, in patients with lumbar radiculopathy, following 4 weeks of
 6 treatment.

7 **References**

9 Agency for Health Care Research and Quality (2007). Decompression Therapy for the
 10 Treatment of Lumbosacral Pain. *Centers for Medicare and Medicaid Services*, 100-
 11 3(1), 1-84.

13 Amjad F, Mohseni-Bandpei MA, Gilani SA, Ahmad A, Hanif A. Effects of non-surgical
 14 decompression therapy in addition to routine physical therapy on pain, range of motion,
 15 endurance, functional disability and quality of life versus routine physical therapy
 16 alone in patients with lumbar radiculopathy; a randomized controlled trial. *BMC*
 17 *Musculoskelet Disord*. 2022;23(1):255. Published 2022 Mar 16. doi:10.1186/s12891-
 18 022-05196-x

20 Apfel CC, Cakmakkaya OS, Martin W, Richmond C, Macario A, George E, et al.
 21 Restoration of disk height through non-surgical spinal decompression is associated
 22 with decreased discogenic low back pain: a retrospective cohort study. *BMC*
 23 *Musculoskelet Disord*. 2010 Jul 8;11:155.

25 Beattie PF, Nelson RM, Michener LA, Cammarata J, Donley J. Outcomes after a prone
 26 lumbar traction protocol for patients with activity-limiting low back pain: a prospective
 27 case series study. *Arch Phys Med Rehabil*. 2008 Feb;89(2):269-74.

29 Beurskens, A. J., H. C. de Vet, et al. (1997). Efficacy of traction for nonspecific low back
 30 pain. 12-week and 6-month results of a randomized clinical trial. *Spine*, 22(23): 2756-
 31 62.

33 Borman P., Keskin D., et al. (2003). The efficacy of lumbar traction in the management of
 34 patients with low back pain. *Rheumatology International*, 23(2): 82-6.

36 Centers for Medicare & Medicaid Services (CMS). National Coverage Determination
 37 160.16. Vertebral axial decompression (VAX-D). Effective date April 15, 1997.
 38 Accessed October 25, 2023. Available at URL address: [http://www.cms.gov/medicare-
 39 coverage-database/details/ncd-
 40 details.aspx?NCDId=124&ncdver=1&bc=BAABAAAAAAAA&](http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=124&ncdver=1&bc=BAABAAAAAAAA&)

- 1 Choi, J., Lee, S., & Hwangbo, G. (2015). Influences of spinal decompression therapy and
 2 general traction therapy on the pain, disability, and straight leg raising of patients with
 3 intervertebral disc herniation. *Journal of physical therapy science*, 27(2), 481–483.
 4 <https://doi.org/10.1589/jpts.27.481>
 5
- 6 Daniel DM. Non-surgical spinal decompression therapy: does the scientific literature
 7 support efficacy claims made in the advertising media? *Chiropr Osteopat.* 2007 May
 8 18;15:7.
 9
- 10 Deen, H. G., Jr., T. D. Rizzo, et al. (2003). Sudden progression of lumbar disc protrusion
 11 during vertebral axial decompression traction therapy. *Mayo Clinic Proceedings*,
 12 78(12): 1554-6.
 13
- 14 Demirel A, Yorubulut M, Ergun N. Regression of lumbar disc herniation by physiotherapy.
 15 Does non-surgical spinal decompression therapy make a difference? Double-blind
 16 randomized controlled trial. *J Back Musculoskelet Rehabil.* 2017 Sep 22;30(5):1015-
 17 1022.
 18
- 19 Gose, E. E., W. K. Naguszewski, et al. (1998). Vertebral axial decompression therapy for
 20 pain associated with herniated or degenerated discs or facet syndrome: an outcome
 21 study. *Neurology Research*, 20(3): 186-90.
 22
- 23 Harrison, D. E., R. Cailliet, et al. (2002). Changes in sagittal lumbar configuration with a
 24 new method of extension traction: nonrandomized clinical controlled trial. *Archives of*
 25 *Physical Medicine and Rehabilitation*, 83(11): 1585-91.
 26
- 27 Harte, A. A., G. D. Baxter, et al. (2003). The efficacy of traction for back pain: a systematic
 28 review of randomized controlled trials. *Archives of Physical Medicine and*
 29 *Rehabilitation*, 84(10): 1542-53.
 30
- 31 Kang, J. I., Jeong, D. K., & Choi, H. (2016). Effect of spinal decompression on the lumbar
 32 muscle activity and disk height in patients with herniated intervertebral disk. *Journal*
 33 *of physical therapy science*, 28(11), 3125–3130. <https://doi.org/10.1589/jpts.28.3125>
 34
- 35 Krause, M., K. M. Refshauge, et al. (2000). Lumbar spine traction: evaluation of effects
 36 and recommended application for treatment. *Manual Therapy*, 5(2): 72-81.
 37
- 38 Lee, R. Y. and J. H. Evans (2001). Loads in the lumbar spine during traction therapy.
 39 *Australian Journal of Physiotherapy*, 47(2): 102-8.
 40
- 41 Macario, A., & Pergolizzi, J. V. (2006). Systematic literature review of spinal
 42 decompression via motorized traction for chronic discogenic low back pain. *Pain*

- 1 *practice : the official journal of World Institute of Pain*, 6(3), 171–178.
2 <https://doi.org/10.1111/j.1533-2500.2006.00082.x>
3
- 4 Macario, A., Richmond, C., Auster, M., & Pergolizzi, J. V. (2008). Treatment of 94
5 outpatients with chronic discogenic low back pain with the DRX9000: a retrospective
6 chart review. *Pain practice : the official journal of World Institute of Pain*, 8(1), 11–
7 17. <https://doi.org/10.1111/j.1533-2500.2007.00167.x>
8
- 9 Maher, C. G. (2004). Effective physical treatment for chronic low back pain. *The*
10 *Orthopedic Clinics of North America*. 35(1): 57-64.
11
- 12 Medical Services Advisory Committee. (2001, June). [MSAC Application 1012].
13 Canberra, Australia: Medical Services Advisory Committee. Accessed October 25,
14 2023 from <http://www.msac.gov.au/internet/msac/publishing.nsf/Content/1012-public>
15
- 16 Meszaros, T. F., R. Olson, et al. (2000). Effect of 10%, 30%, and 60% body weight traction
17 on the straight leg raise test of symptomatic patients with low back pain. *The Journal*
18 *of Orthopaedic and Sports Physical Therapy*. 30(10): 595-601.
19
- 20 Naguszewski, W. K., R. K. Naguszewski, et al. (2001). Dermatomal somatosensory
21 evoked potential demonstration of nerve root decompression after VAX-D therapy.
22 *Neurology Research*, 23(7): 706-14.
23
- 24 Pellecchia, G. L. (1994). Lumbar traction: a review of the literature. *The Journal of*
25 *Orthopaedic and Sports Physical Therapy* 20(5): 262-7.
26
- 27 Philadelphia Panel. (2001). Philadelphia Panel evidence-based clinical practice guidelines
28 on selected rehabilitation interventions for low back pain. *Physical Therapy*.
29 81(10):1641-74.
30
- 31 Ramos, G. (2004). Efficacy of vertebral axial decompression on chronic low back pain:
32 study of dosage regimen. *Neurological Research* 26(3): 320-4.
33
- 34 Ramos, G. and W. Martin (1994). Effects of vertebral axial decompression on intradiscal
35 pressure. *Journal of Neurosurgery*. 81(3): 350-3.
36
- 37 Revel, M. (2000). Does traction still have a role in nonspecific low back disorders? *Joint*
38 *Bone Spine* 67(3): 146-9.
39
- 40 Schimmel, J. J., de Kleuver, M., Horsting, P. P., Spruit, M., Jacobs, W. C., & van Limbeek,
41 J. (2009). No effect of traction in patients with low back pain: a single centre, single
42 blind, randomized controlled trial of Intervertebral Differential Dynamics

1 Therapy. *European spine journal* : official publication of the European Spine Society,
2 the European Spinal Deformity Society, and the European Section of the Cervical Spine
3 Research Society, 18(12), 1843–1850. <https://doi.org/10.1007/s00586-009-1044-3>
4

5 Shealy, CN. Borgmeyer, V. (1997). Decompression, reduction, and stabilization of the
6 lumbar spine: a cost effective treatment for lumbosacral pain. *American Journal of*
7 *Pain Management*. 7(2):63-65.
8

9 Sherry, E., P. Kitchener, et al. (2001). A prospective randomized controlled study of VAX-
10 D and TENS for the treatment of chronic low back pain. *Neurological Research* 23(7):
11 780-4.
12

13 Werners, R., P. B. Pynsent, et al. (1999). Randomized trial comparing interferential therapy
14 with motorized lumbar traction and massage in the management of low back pain in a
15 primary care setting. *Spine* 24(15): 1579-84.