

1 **Clinical Practice Guideline:** **Axial/Spinal Decompression Therapy**
 2
 3 **Date of Implementation:** **July 13, 2006**
 4
 5 **Effective Date:** **January 29, 2026**
 6
 7 **Product:** **Specialty**
 8
 9

10 **GUIDELINES**

11 American Specialty Health – Specialty (ASH) considers nonsurgical axial/spinal
 12 decompression therapy to be unproven due to insufficient scientific evidence of efficacy in
 13 the treatment of neck, low back, and related disorders. This includes any motorized
 14 mechanical traction device that is promoted as providing ‘decompression therapy’ (e.g.,
 15 VAX-D, IDD Therapy® [Intervertebral Differential Dynamics Therapy], DRS, DRX,
 16 DRX-2000, DRX-3000, DRX-5000, DRX-9000, Accu-SPINA™, Lordex Power Traction
 17 device, Mettler Traction Device [MTD 4000], Tru Trac 401, Integrity Spinal Care System
 18 Alpha-SPINA System, Dynatron DX2, Dynapro™ DX2, Spinerx LDM, or any other
 19 device that claims to create spinal decompression).
 20

21 The research evidence concerning nonsurgical axial/spinal decompression therapy is
 22 lacking and of low quality. Any estimate of treatment effect is uncertain, as is the clarity
 23 of risk, benefit, and burden to the patient.
 24

25 There are significant burdens placed upon health plan members due to high out-of-pocket
 26 costs, time spent receiving the intervention, and the unsubstantiated/misleading marketing
 27 about the alleged proven effectiveness and safety of nonsurgical axial/spinal
 28 decompression therapy. These burdens have been recognized as significant by some
 29 professional licensing boards and state justice departments.
 30

31 Similar conclusions have been reached by a broad range of health care organizations.
 32 Professionals and groups, who are proponents of nonsurgical axial/spinal decompression
 33 therapy, should pursue further investigation using experimental study designs and rigorous
 34 methodologies.
 35

HCPCS/CPT Code	HCPCS/CPT Code Description
S9090	Vertebral Axial Decompression, per session; {most accurately describes services for the application of spinal decompression motorized traction devices}
Other CPT codes that have been associated with the use of nonsurgical spinal decompression therapy are:	

97012	Application of a modality to 1 or more areas; traction, mechanical
-------	--

1
2 **DESCRIPTION/BACKGROUND**

3 Traction as a treatment option for low back pain and sciatica has existed for many years.
4 Its use has progressed from continuous static traction to intermittent motorized traction.
5 The most recent form of intermittent motorized traction is commonly referred to as
6 axial/spinal decompression therapy. Developers and manufacturers of the equipment along
7 with clinicians often consider it to be a unique form of traction. Proponents of nonsurgical
8 axial/spinal decompression therapy claim it to be a safe and effective alternative to surgical
9 interventions. Companies demonstrate intense marketing programs and claim high success
10 rates. Axial/spinal decompression therapy is intended to create negative pressure within
11 the spine so that as the spinal column is elongated, pressure is taken off the nerve root(s),
12 and herniated disc material may be pulled back into place. Axial/spinal decompression
13 therapy is generally performed using a specially designed computerized mechanical table
14 that separates in the middle. Depending on the type of table being used, a patient is strapped
15 in a prone or supine position to the lower part of the table using a pelvic harness and may
16 hold handgrips at the top of the table. The table is then mechanically separated in the
17 middle creating a distractive force to relieve pressure within the spine that may be causing
18 pain. The amount of distractive force is tailored for each patient and usually lasts about 60
19 seconds. Depending on the device utilized, static, intermittent, or cycled distractive force
20 may be applied. Typical treatment protocols include 20 sessions, each lasting 30 to 40
21 minutes. The process of distraction and relaxation is fully computerized using a
22 programmable logic controller and is monitored by a licensed health care practitioner. The
23 American Medical Association (AMA), Food and Drug Administration (FDA), and
24 Centers for Medicare & Medicaid Services (CMS) all consider axial/spinal decompression
25 therapy to be a form of traction. However, this therapy involves a special table and protocol
26 that isn't the same as conventional or traditional traction with claims of spinal
27 decompression.

28
29 The tables utilized for axial/spinal decompression therapy are classified by the FDA as
30 powered traction equipment. Examples of axial/spinal decompression therapy tables (and
31 their manufacturers) include:

- 32 • VAX-D Table (VAX-D Manufacturing, Palm Harbor, FL)
- 33 • Decompression, Reduction, Stabilization (DRS) System (North American Medical
34 Corporation, Atlanta, GA)
- 35 • DRX 2000 and DRX 9000 (Axiom Worldwide, Tampa, FL)
- 36 • Spina System (North American Medical Corporation, Atlanta, GA)

37
38 Two popular units will be described here. Due to the number of available products, it would
39 be impractical to provide information on all of them.

VAX-D

The manufacturer suggests that use of the VAX-D table applies distractive forces in a gradual, progressive fashion through extension of the lower end of the table. The level of tension is preset on a control panel and can be increased, allowing for various decompression phases and a rest phase. Various decompression phases allow alternating cycles of distraction and relaxation. Typically, a treatment cycle consists of 15 cycles of tension and relaxation. The patient lies prone on the VAX-D table. The table is split, allowing the table to slowly extend, thus decreasing load bearing in the intervertebral discs and/or intervertebral joint spaces. The VAX-D manufacturer claims specific parameters of their system make the device inherently safe. These safety features include the use of air pressure as the energy source; the ramp characteristics employed in applying the distraction tensions; the release rate of the distraction and relaxation cycles; the cycle periodicity; the upper limits on the distraction tensions; the positioning of the patient and the means of fixing the upper body; and the ability of the patient to release the handgrips if the distraction tension causes pain or discomfort. Information regarding the range and incidence of adverse effects that occur during VAX-D therapy is limited. Complications reported with VAX-D include:

- The development of a sharp burning, radiating pain during therapy
- Stress to the shoulder girdle and rotator cuff muscles
- Overstretching of the soft tissue of the back

Decompression, Reduction, Stabilization (DRS) System

Manufacturers recommend the Decompression, Reduction, Stabilization (DRS) System for treatment of low back pain. This device uses a bed that is split into two cushions. The patient can step onto a foot pad, have a pelvic and chest harness attached, after which the patient and bed are lowered to a horizontal position. Distraction tension is applied by the pelvic harness while the patient's upper body is secured to the locked upper cushion via the chest harness. The DRS System is marketed for the treatment of low back pain associated with herniated and degenerated discs. According to the manufacturer, the DRS System applies pressures on the disc in a graduated manner, which bypasses the inherent neurological mechanisms that lead to firing of stretch receptors in the paravertebral structures. This decreased resistance to the distractive forces allows a reduction in intradiscal pressures, which promotes retraction of herniated disc material and facilitates influx of oxygen, proline, and other substrates.

EVIDENCE REVIEW

Currently, there is not adequate scientific evidence which proves that axial/spinal decompression is an effective single intervention or adjunct to conservative therapy for back pain. In addition, axial/spinal decompression devices have not been adequately studied as alternatives to back surgery.

1 Proponents of nonsurgical axial/spinal decompression therapy assert this form of traction
2 is, however, unique for being proven able to reduce the relative pressure measured within
3 intervertebral discs (decompression). The evidence typically cited to support this claim is
4 from a study by Ramos, 1994. An evaluation of this study shows the conclusions are based
5 upon data from only three subjects. This study demonstrated a number of methodological
6 flaws likely to invalidate the results. These included not using a closed transducer system,
7 not taking into account temperature effects, absent hydrostatic conditions (in degenerative
8 discs), and no attempt reported to calibrate negative readings.

9
10 Regardless of the flaws, this study is not sufficient to arrive at conclusions about the
11 translation of basic science research into clinical care settings. The author (Ramos)
12 concluded additional study is needed to establish the relationship of negative intradiscal
13 pressures with clinical outcomes. The results from an early uncontrolled, retrospective
14 study (Gose et al., 1998) regarding the benefits of the VAX-D table appeared to be
15 encouraging. However, the findings need to be validated in prospective, randomized,
16 controlled clinical trials because the study was poorly designed. A subsequent randomized
17 study (Sherry et al., 2001) compared VAX-D to transcutaneous electrical nerve stimulation
18 (TENS) in the treatment of patients with chronic (> 3 months in duration) low back pain.
19 Successful outcome was defined as a 50% decrease in pain using the Visual Analog Pain
20 Scale and an improvement in the level of functioning as measured by patient-nominated
21 disability ratings. The TENS-treated group ($n=21$) reported a success rate of 0%, while the
22 group treated with VAX-D ($n=19$) showed a success rate of 68.4%. No confirmatory
23 conclusions can be drawn from this study given detailed statistics regarding the outcomes
24 for each group was not included in the analysis. Furthermore, patients were not blinded to
25 the treatment received. The Australian Medical Services Advisory Committee (MSAC,
26 2001) performed an assessment of the literature on VAX-D therapy. The Committee
27 concluded that "there is currently insufficient evidence pertaining to the effectiveness of
28 vertebral axial decompression (VAX-D) therapy..." In 2007, they requested that the
29 Agency for Healthcare Research and Quality (AHRQ) commission an evidence-based
30 technology assessment. The AHRQ report "Decompression Therapy for the Treatment of
31 Lumbosacral Pain" concluded the current evidence regarding the efficacy of axial/spinal
32 decompression therapy is too limited in quality and quantity to allow for evidence-based
33 conclusions. Adverse event reporting for axial/spinal decompression therapy was viewed
34 as infrequent. The Centers for Medicare & Medicaid Services (CMS) Technology
35 Advisory Committee did not recommend coverage of the VAX-D system because of the
36 absence of scientific data on its effectiveness.

37
38 In review of a single study of DRS therapy (Shealy and Borgmeyer, 1997), the authors
39 reported on a comparison of DRS therapy to conventional traction for both ruptured lumbar
40 discs and chronic facet arthrosis. This study suffered from three major flaws: one of the
41 authors was affiliated with the treatment center that conducted the trial; the scale used to

1 quantify the results was not clearly defined; and the study consisted of a small sample size
2 lacking clearly defined methods of randomization.

3
4 Macario and Pergolizzi (2006) conducted a systematic review of the literature to assess the
5 efficacy of nonsurgical axial/spinal decompression that is achieved with motorized traction
6 for chronic discogenic low back pain. The authors reviewed data from 10 studies between
7 1975 and 2003. Seven studies were randomized controlled trials of motorized traction
8 using various apparatus types, including split-tabletop, plain tabletop, and friction-free
9 couch with weights. A total of 408 individuals received placebo, and 438 individuals
10 received motorized spinal decompression. Follow-up averaged 28 weeks. None of the
11 studies were blinded, and only three had descriptions of the randomization method. Six of
12 the seven randomized trials reported no difference with motorized spinal decompression,
13 and one study reported reduced pain but not disability. In the author's opinion, the efficacy
14 of spinal decompression achieved with motorized traction for discogenic low back remains
15 unproven. Daniel (2007) reported that there is very limited evidence in the scientific
16 literature to support the effectiveness of non-surgical axial/spinal decompression therapy.
17 One randomized controlled trial, one clinical trial, one case series and seven other papers
18 were available in the published literature for review by the author as part of an intended
19 systematic review. Due to the limited evidence a systematic review was not done, and each
20 study was reviewed individually. The author noted many of the reviewed studies utilized
21 the VAX-D unit. Furthermore, the intervention has not been compared to exercise, spinal
22 manipulation, standard medical care, or other less expensive conservative treatments.

23
24 In a prospective case series study, Beattie et al. (2008) examined outcomes after an
25 intervention of a prone lumbar traction protocol using the VAX-D system. A total of 296
26 subjects with low back pain and evidence of a degenerative and/or herniated intervertebral
27 disc at one or more levels of the lumbar spine were included in this study. Patients
28 underwent an 8-week course of prone lumbar traction, using the VAX-D system, consisting
29 of five 30-minute sessions a week for four weeks, followed by one 30-min session a week
30 for four additional weeks. The numeric pain rating scale and the Roland-Morris Disability
31 Questionnaire were completed at pre-intervention, discharge (within two weeks of the last
32 visit), and at 30 days and 180 days after discharge. A total of 250 (84.4 %) subjects
33 completed the treatment protocol. On the 30-day follow-up, 247 (83.4 %) subjects were
34 available; on the 180-day follow-up, data were available for 241 (81.4 %) subjects. These
35 researchers noted significant improvements for all post-intervention outcome scores when
36 compared with pre-intervention scores ($p < 0.01$). The authors noted that causal
37 relationships between the outcomes and the intervention cannot be made. This study lacked
38 a comparison group.

39
40 Macario et al. (2008) discussed the retrospective chart audit of 100 patients with discogenic
41 low back pain (LBP) lasting more than 12 weeks, treated with a 2-month course of
42 motorized spinal decompression via the DRX9000. Patients at a convenience sample of 4

1 clinics received 30-min DRX9000 sessions daily for the first 2 weeks tapering to 1
2 session/week. Treatment protocol included lumbar stretching, myofascial release, or heat
3 prior to treatment, with ice and/or muscle stimulation afterwards. Primary outcome was
4 verbal NRS 0 to 10 before and after the 8-week treatment. Of the 100 subjects, three
5 withdrew their protected health information, and three were excluded because their LBP
6 duration was less than 12 weeks. The remaining 94 subjects had diagnoses of herniated
7 disc (73% of patients), degenerative disc disease (68 %), or both (27%). Mean NRS equaled
8 6.05 (SD 2.3) at presentation and decreased significantly to 0.89 (SD 1.15) at end of 8-
9 week treatment ($p < 0.0001$). Analgesic use also appeared to decrease (charts with data =
10 20) and activities of daily living improved (charts with data = 38). Follow-up (mean of 31
11 weeks) on 29/94 patients reported mean 83% LBP improvement, NRS of 1.7 (SD 1.15),
12 and satisfaction of 8.55/10 (median of 9). The authors concluded that this retrospective
13 chart audit provides preliminary data that chronic LBP may improve with DRX9000 spinal
14 decompression, however caution should be taken with this interpretation given it was not
15 provided as a singular treatment. They stated that randomized double-blind trials are
16 needed to measure the effectiveness of such systems. Schimmel et al. (2009) conducted a
17 randomized sham-controlled trial of intervertebral axial decompression. Sixty subjects
18 with chronic symptomatic lumbar disc degeneration or bulging disc with no radicular pain
19 and no prior surgical treatment (dynamic stabilization, fusion, or disc replacement) were
20 randomly assigned to a graded activity program with an Accu-SPINA device (20 traction
21 sessions during six weeks, reaching $>50\%$ body weight), or to a graded activity program
22 with a non-therapeutic level of traction ($<10\%$ body weight). In addition to traction, the
23 device provided massage, heat, blue relaxing light, and music during the treatment sessions
24 in both groups. Neither patients nor evaluators were informed about the intervention
25 received until after the 14-week follow-up assessment, and intention-to-treat analysis was
26 performed (93% of subjects completed follow-up). Both groups showed improvements in
27 validated outcome measures (visual analog scores for back and leg pain, Oswestry
28 Disability Index, and Short-Form 36), with no differences between the treatment groups.
29 The authors reported that the added axial, intermittent, mechanical traction of IDD Therapy
30 to a standard graded activity program has been shown not to be effective.

31
32 Apfel et al. (2010) conducted a retrospective cohort study of adults with chronic LBP
33 attributed to disc herniation and/or discogenic LBP who underwent a six-week treatment
34 protocol of motorized non-surgical spinal decompression via the DRX9000. The main
35 outcomes were changes in pain as measured on a verbal rating scale during a flexion-
36 extension range of motion evaluation and changes in disc height as measured on CT scans.
37 The authors identified 30 patients with lumbar disc herniation and an average duration of
38 LBP of 12.5 weeks. During treatment, low back pain decreased from 6.2 (SD 2.2) to 1.6
39 (2.3, $p < 0.001$) and disc height increased from 7.5 (1.7) mm to 8.8 (1.7) mm ($p < 0.001$).
40 Increase in disc height and reduction in pain were significantly correlated ($r = 0.36$,
41 $p = 0.044$). Reported limitations of this study are no control group and small sample size.
42 The authors reported that a randomized controlled trial is needed to confirm the efficacy

1 and elucidate the mechanism of this treatment modality. Choi et al. (2015) sought to
2 identify how spinal decompression therapy and general traction therapy influence the pain,
3 disability, and straight leg raise (SLR) ability of patients with intervertebral disc herniation.
4 The subjects were 30 patients with chronic lumbar pain who were divided into a spinal
5 decompression therapy group using a spinal decompression device (SDTG, $n=15$), and a
6 general traction therapy group (GTTG, $n=15$). Both groups received conservative physical
7 therapy three times a week for four weeks. A comparison of the two groups found no
8 statistically significant differences. Authors concluded that spinal decompression therapy
9 and general traction therapy are effective at improving the pain, disability, and SLR of
10 patients with intervertebral disc herniation. Limitations of the study from a methodology
11 standpoint do not allow conclusions to be confirmed. Kang et al. (2016) conducted a study
12 to clarify the difference in therapeutic effects between traction and decompression
13 therapies, and their clinical therapeutic significance. For the experimental group, 15
14 subjects were randomly selected to receive decompression therapy and trunk stabilization
15 exercise. For the control group, 16 subjects were randomly selected to receive traction
16 therapy and trunk stabilization exercise. Authors concluded that decompression therapy
17 was demonstrated to be more effective clinically than conventional traction therapy as an
18 intervention method for disk disease.

19
20 Demirel et al., (2017) sought to determine whether non-invasive spinal decompression
21 therapy (NSDT) was effective in resorption of herniation, increasing disc height in patients
22 with lumbar disc herniation (LHNP). A total of twenty patients diagnosed as LHNP and
23 suffering from pain at least 8 weeks were enrolled to the study. Patients were randomly
24 allocated in study (SG) and control groups (CG). Both groups received combination of
25 electrotherapy, deep friction massage and stabilization exercise for fifteen sessions. SG
26 received additionally NSDT different from CG. Numeric Analog Scale, Straight leg raise
27 test, Oswestry Disability Index (ODI) were applied at baseline and after treatment. Disc
28 height and herniation thickness were measured on MRI which performed at baseline and
29 three months after therapy. Both treatments had positive effect for improving pain,
30 functional restoration, and reduction in thickness of herniation. Although reduction of
31 herniation size was higher in SG than CG, no significant differences were found between
32 groups and any superiority to each other ($p > 0.05$). Given the study design, the study
33 showed that physiotherapy was helpful but that adding NSDT did not confer additional
34 benefits. Amjad et al. (2022) sought to determine the effects of non-surgical spinal
35 decompression (NSD) therapy in addition to routine physical therapy on pain, lumbar range
36 of motion (ROM), functional disability, back muscle endurance (BME), and quality of life
37 (QOL) in patients with lumbar radiculopathy. A total of 60 patients with lumbar
38 radiculopathy were randomly allocated into two groups, an experimental ($n = 30$) and a
39 control ($n = 30$) group, through a computer-generated random number table. Baseline
40 values were recorded before providing any treatment by using a visual analogue scale
41 (VAS), Urdu version of Oswestry disability index (ODI-U), modified-modified Schober's
42 test (MMST), prone isometric chest raise test, and Short Form 36-Item Survey (SF-36) for

1 measuring the pain at rest, functional disability, lumbar ROM, BME, and QOL,
 2 respectively. All patients received twelve treatment sessions over 4 weeks, and then all
 3 outcome measures were again recorded. By using the ANCOVA test, a statistically
 4 significant ($p < 0.05$) between-group improvement was observed in VAS, ODI-U, BME,
 5 lumbar ROM, role physical (RP), and bodily pain (BP) domains of SF-36, which was in
 6 favor of NSD therapy group. For these outcomes, a medium to large effect size ($d = 0.61$ -
 7 2.47 , 95% CI: 0.09-3.14) was observed. It was concluded that a combination of non-
 8 surgical spinal decompression therapy with routine physical therapy is more effective,
 9 statistically and clinically, than routine physical therapy alone in terms of improving pain,
 10 lumbar range of motion, back muscle endurance, functional disability, and physical role
 11 domain of quality of life, in patients with lumbar radiculopathy, following 4 weeks of
 12 treatment.

13 14 **References**

- 15 Agency for Health Care Research and Quality (2007). Decompression Therapy for the
 16 Treatment of Lumbosacral Pain. *Centers for Medicare and Medicaid Services*, 100-
 17 3(1), 1-84
- 18
- 19 Amjad F, Mohseni-Bandpei MA, Gilani SA, Ahmad A, Hanif A. Effects of non-surgical
 20 decompression therapy in addition to routine physical therapy on pain, range of motion,
 21 endurance, functional disability and quality of life versus routine physical therapy
 22 alone in patients with lumbar radiculopathy; a randomized controlled trial. *BMC*
 23 *Musculoskelet Disord*. 2022;23(1):255. Published 2022 Mar 16. doi:10.1186/s12891-
 24 022-05196-x
- 25
- 26 Apfel CC, Cakmakkaya OS, Martin W, Richmond C, Macario A, George E, et al.
 27 Restoration of disk height through non-surgical spinal decompression is associated
 28 with decreased discogenic low back pain: a retrospective cohort study. *BMC*
 29 *Musculoskelet Disord*. 2010 Jul 8;11:155
- 30
- 31 Beattie PF, Nelson RM, Michener LA, Cammarata J, Donley J. Outcomes after a prone
 32 lumbar traction protocol for patients with activity-limiting low back pain: a prospective
 33 case series study. *Arch Phys Med Rehabil*. 2008 Feb;89(2):269-74
- 34
- 35 Beurskens, A. J., H. C. de Vet, et al. (1997). Efficacy of traction for nonspecific low back
 36 pain. 12-week and 6-month results of a randomized clinical trial. *Spine*, 22(23): 2756-
 37 62
- 38
- 39 Borman P., Keskin D., et al. (2003). The efficacy of lumbar traction in the management of
 40 patients with low back pain. *Rheumatology International*, 23(2): 82-6

- 1 Centers for Medicare & Medicaid Services (CMS). National Coverage Determination
 2 160.16. Vertebral axial decompression (VAX-D). Effective date April 15, 1997.
 3 Accessed November 24, 2025. Available at URL address:
 4 <http://www.cms.gov/medicare-coverage-database/details/ncd->
 5 [details.aspx?NCDId=124&ncdver=1&bc=BAABAAAAAAAA&](http://www.cms.gov/medicare-coverage-database/details/ncd-)
 6
- 7 Choi, J., Lee, S., & Hwangbo, G. (2015). Influences of spinal decompression therapy and
 8 general traction therapy on the pain, disability, and straight leg raising of patients with
 9 intervertebral disc herniation. *Journal of Physical Therapy Science*, 27(2), 481–483.
 10 <https://doi.org/10.1589/jpts.27.481>
 11
- 12 Daniel DM. Non-surgical spinal decompression therapy: does the scientific literature
 13 support efficacy claims made in the advertising media? *Chiropr Osteopat*. 2007 May
 14 18;15:7
 15
- 16 Deen, H. G., Jr., T. D. Rizzo, et al. (2003). Sudden progression of lumbar disc protrusion
 17 during vertebral axial decompression traction therapy. *Mayo Clinic Proceedings*,
 18 78(12): 1554-6
 19
- 20 Demirel A, Yorubulut M, Ergun N. Regression of lumbar disc herniation by physiotherapy.
 21 Does non-surgical spinal decompression therapy make a difference? Double-blind
 22 randomized controlled trial. *J Back Musculoskelet Rehabil*. 2017 Sep 22;30(5):1015-
 23 1022
 24
- 25 Gose, E. E., W. K. Naguszewski, et al. (1998). Vertebral axial decompression therapy for
 26 pain associated with herniated or degenerated discs or facet syndrome: an outcome
 27 study. *Neurology Research*, 20(3): 186-90
 28
- 29 Harrison, D. E., R. Cailliet, et al. (2002). Changes in sagittal lumbar configuration with a
 30 new method of extension traction: nonrandomized clinical controlled trial. *Archives of*
 31 *Physical Medicine and Rehabilitation*, 83(11): 1585-91
 32
- 33 Harte, A. A., G. D. Baxter, et al. (2003). The efficacy of traction for back pain: a systematic
 34 review of randomized controlled trials. *Archives of Physical Medicine and*
 35 *Rehabilitation*, 84(10): 1542-53
 36
- 37 Kang, J. I., Jeong, D. K., & Choi, H. (2016). Effect of spinal decompression on the lumbar
 38 muscle activity and disk height in patients with herniated intervertebral disk. *Journal*
 39 *of physical therapy science*, 28(11), 3125–3130. <https://doi.org/10.1589/jpts.28.3125>
 40
- 41 Krause, M., K. M. Refshauge, et al. (2000). Lumbar spine traction: evaluation of effects
 42 and recommended application for treatment. *Manual Therapy*, 5(2): 72-81

- 1 Lee, R. Y. and J. H. Evans (2001). Loads in the lumbar spine during traction therapy.
2 *Australian Journal of Physiotherapy*, 47(2): 102-8
3
- 4 Macario, A., & Pergolizzi, J. V. (2006). Systematic literature review of spinal
5 decompression via motorized traction for chronic discogenic low back pain. *Pain
6 practice: the official journal of World Institute of Pain*, 6(3), 171–178.
7 <https://doi.org/10.1111/j.1533-2500.2006.00082.x>
8
- 9 Macario, A., Richmond, C., Auster, M., & Pergolizzi, J. V. (2008). Treatment of 94
10 outpatients with chronic discogenic low back pain with the DRX9000: a retrospective
11 chart review. *Pain practice: the official journal of World Institute of Pain*, 8(1), 11–
12 17. <https://doi.org/10.1111/j.1533-2500.2007.00167.x>
13
- 14 Maher, C. G. (2004). Effective physical treatment for chronic low back pain. *The
15 Orthopedic Clinics of North America*. 35(1): 57-64.
16
- 17 Medical Services Advisory Committee. (2001, June). [MSAC Application 1012].
18 Canberra, Australia: Medical Services Advisory Committee. Accessed November 20,
19 2024 from <http://www.msac.gov.au/internet/msac/publishing.nsf/Content/1012-public>
20
- 21 Meszaros, T. F., R. Olson, et al. (2000). Effect of 10%, 30%, and 60% body weight traction
22 on the straight leg raise test of symptomatic patients with low back pain. *The Journal
23 of Orthopaedic and Sports Physical Therapy*. 30(10): 595-601
24
- 25 Naguszewski, W. K., R. K. Naguszewski, et al. (2001). Dermatomal somatosensory
26 evoked potential demonstration of nerve root decompression after VAX-D therapy.
27 *Neurology Research*, 23(7): 706-14
28
- 29 Pelledchia, G. L. (1994). Lumbar traction: a review of the literature. *The Journal of
30 Orthopaedic and Sports Physical Therapy* 20(5): 262-7
31
- 32 Philadelphia Panel. (2001). Philadelphia Panel evidence-based clinical practice guidelines
33 on selected rehabilitation interventions for low back pain. *Physical Therapy*.
34 81(10):1641-74
35
- 36 Ramos, G. (2004). Efficacy of vertebral axial decompression on chronic low back pain:
37 study of dosage regimen. *Neurological Research* 26(3): 320-4
38
- 39 Ramos, G. and W. Martin (1994). Effects of vertebral axial decompression on intradiscal
40 pressure. *Journal of Neurosurgery*. 81(3): 350-3

- 1 Revel, M. (2000). Does traction still have a role in nonspecific low back disorders? *Joint*
2 *Bone Spine* 67(3): 146-9
3
- 4 Schimmel, J. J., de Kleuver, M., Horsting, P. P., Spruit, M., Jacobs, W. C., & van Limbeek,
5 J. (2009). No effect of traction in patients with low back pain: a single centre, single
6 blind, randomized controlled trial of Intervertebral Differential Dynamics
7 Therapy. *European Spine Journal : Official Publication of the European Spine Society,*
8 *the European Spinal Deformity Society, and the European Section of the Cervical Spine*
9 *Research Society, 18(12), 1843–1850.* <https://doi.org/10.1007/s00586-009-1044-3>
10
- 11 Shealy, CN. Borgmeyer, V. (1997). Decompression, reduction, and stabilization of the
12 lumbar spine: a cost effective treatment for lumbosacral pain. *American Journal of*
13 *Pain Management.* 7(2):63-65
14
- 15 Sherry, E., P. Kitchener, et al. (2001). A prospective randomized controlled study of VAX-
16 D and TENS for the treatment of chronic low back pain. *Neurological Research* 23(7):
17 780-4
18
- 19 Werners, R., P. B. Pynsent, et al. (1999). Randomized trial comparing interferential therapy
20 with motorized lumbar traction and massage in the management of low back pain in a
21 primary care setting. *Spine* 24(15): 1579-84