

1	Clinical Practice Guideline:	Manipulation Under Anesthesia (MUA)
2		
3	Date of Implementation:	July 13, 2006
4		
5	Effective Date:	February 19, 2026
6		
7	Product:	Specialty
8		

9 **Table of Contents**

10 GUIDELINES 1

11 DESCRIPTION/BACKGROUND 2

12 EVIDENCE REVIEW..... 3

13 Spine 3

14 Shoulder..... 4

15 Knee10

16 Fracture and/or Dislocation16

17 Chronic Joint Contracture.....16

18 Elbow16

19 TMJ17

20 Other Joints and Conditions17

21 PRACTITIONER SCOPE AND TRAINING17

22 *References*18

23

24 **GUIDELINES**

25 American Specialty Health – Specialty (ASH) considers one session of manipulation under
 26 anesthesia (MUA) medically necessary for the following indications:

- 27 • Adhesive capsulitis (i.e., frozen shoulder) when there is failure of conservative
 28 management, including medications with or without articular injections, home
 29 exercise programs and physical therapy for at least 6 to 8 weeks at a minimum (CPT
 30 code 23700).
- 31 • Post-traumatic or postoperative arthrofibrosis of the knee (e.g., total knee
 32 replacement, anterior cruciate ligament reconstruction) when there is failure of
 33 conservative management, including exercise and physical therapy per surgeon’s
 34 recommendations (CPT code 27570).
- 35 • Reduction of a displaced fracture (e.g., vertebral, long bones) (CPT codes 22505
 36 and 25675).
- 37 • Reduction of acute/traumatic dislocation (e.g., vertebral, perched cervical facet)
 38 (e.g., CPT code 22505).

- Chronic contracture of upper or lower extremity joint (e.g., fixed contracture from a neuromuscular condition) when there is failure of conservative management including range of motion exercise programs and physical therapy for at least 6 to 8 weeks at a minimum.

Manipulation under anesthesia (MUA) is considered safe and effective and is a well-established method of treatment of the above conditions. When performed for these specific conditions, MUA generally requires a single session of treatment, most often performed unilaterally, involving a single joint. Data supporting the need for, and clinical efficacy of multiple, repeat MUA treatment sessions for these specific conditions, is lacking in the peer-reviewed published medical literature.

ASH considers MUA for acute or chronic pain conditions of any of the following joints (other than those listed above as medically necessary) as unproven and thus, not medically necessary:

- Ankle (CPT code 27860)
- Cervical, thoracic or lumbar spine (e.g., CPT code 22505)
- Elbow (CPT code 24300)
- Finger (e.g., CPT code 26340, 26675)
- Hip (CPT code 27275)
- Pelvis, Sacroiliac (CPT code 27198)
- Temporomandibular (CPT code 21073)
- Thumb (CPT code 26340)
- Toe (CPT code 28635, 28665)
- Wrist (CPT code 25259)

The available evidence does not enable ASH to determine if MUA is safe or effective relative to more conservative care. Well-designed studies are needed to evaluate and confirm its place in treatment of neck and low back pain and for other pain conditions related to the above joints.

DESCRIPTION/BACKGROUND

Manipulation under anesthesia (MUA) is the use of manual manipulation of the spine or other joints while the patient is anesthetized. The addition of an anesthetic allows for manipulation under circumstances where conscious manipulation would not be effective because of pain response, spasm, muscle contracture, and/or guarding. The manipulative procedure that the physician performs depends upon the goals of the procedure, the tissues involved, and the presence of potential complications and/or contraindication(s). Treatment may include passive soft tissue stretching, oscillation of joints, and articular adjustments. In general, patients selected for MUA have generally undergone more conservative treatment and failed to improve, unless it is an urgent situation with a

1 displaced vertebral fracture or long bone fracture. As such, in most cases, MUA is not a
2 first line therapy for musculoskeletal conditions.

3
4 The treatment is typically performed in a hospital or surgery center with the assistance of
5 an anesthesiologist. MUA can be performed under varying levels of anesthesia, including
6 general anesthesia, conscious sedation, and local anesthesia. General anesthesia is the most
7 complete form of anesthesia and requires intubation of the patient to help control their
8 breathing and monitor their respiratory function. General anesthesia was more commonly
9 used for MUA in the past, but its use for this procedure has declined notably over the last
10 ten (10) years. Conscious sedation is an intermediary level of anesthesia where the patient
11 is given intravenous or oral sedation that depresses the central nervous system. At this stage
12 of anesthesia, a patient is conscious and does not require intubation. A patient under
13 conscious sedation would not respond to mildly painful stimuli such as being pinched;
14 however, they would respond to severely painful stimuli such as undergoing surgery.
15 Proponents of MUA claim that conscious sedation allows for more patient feedback during
16 treatment than general anesthesia. However, the use of conscious sedation does not allow
17 for the same level of patient feedback as manipulation without any anesthesia. Local
18 anesthesia is another option for MUA, though it is less frequently used than conscious
19 sedation. A local anesthesia involves the injection of an anesthetizing substance at the site
20 where the manipulation will be performed. In this type of anesthesia, the patient remains
21 completely awake and aware of the procedure, but sensations of pain are blocked in the
22 specific area of manipulation. In addition, there are inherent risks to any type of anesthesia.

23
24 Comment on spinal MUA: while MUA of the spine may be considered professionally
25 recognized by certain physician groups (e.g., chiropractors and osteopaths), it may also
26 pose a health and safety risk greater than traditional high-velocity, low-amplitude (HVLA)
27 manipulation for the spine in particular. The use of any anesthesia during joint
28 manipulation does not allow the same level of patient feedback as manipulation without
29 anesthesia. Patient feedback during manipulation is an important safeguard in the
30 prevention of treatment related injury. Although safer than both general anesthesia and
31 conscious sedation, local anesthesia is often considered inappropriate for MUA of the
32 spine.

33 34 **EVIDENCE REVIEW**

35 **Spine**

36 Within the realm of chiropractic, spinal MUA is generally performed daily for 1 to 5
37 consecutive days on an outpatient basis, and is followed by a post-SMUA rehabilitation
38 regimen, which entails 1 week of daily manipulation to maintain joint mobility and avoid
39 re-adhesion of fibrotic tissue. Anesthesia is usually induced by intravenous Pentothal
40 (sodium thiopental), and manipulation of the affected joints takes about 7 to 10 minutes.

1 An old randomized controlled trial by Siehl et al. (1971) evaluated MUA for patients with
 2 spinal nerve root compression. This study could not determine the benefits of MUA due to
 3 the design of the study, which would have required very large differences between groups
 4 to have significance.

5
 6 Review of the literature revealed numerous case series and reports that expounded the
 7 benefits of MUA (Aspegren et al., 1997; Ben-David et al., 1994; Cremata et al., 2005;
 8 Dreyfuss et al., 1995; Herzog, 1999; Maxwell et al., 1994; Tsai and Chou, 2005; West et
 9 al., 1999; Xiong et al., 1998). There were also two non-randomized studies evaluating the
 10 efficacy of MUA. Palmieri and Smoyak (2002) evaluated MUA versus traditional spinal
 11 manipulation in the treatment of low back pain, but their objectives were to evaluate
 12 methods useful for studying the procedure, not to determine the efficacy of MUA for spinal
 13 pain. Although more of the patients reported more improvement in pain with MUA, the
 14 intervention group received treatments other than MUA (e.g., physical therapy) that the
 15 control group did not receive. Due to the design and goal of this study, it is not possible to
 16 attribute the effects seen in the study to MUA. Kohlbeck et al. (2005) found that MUA
 17 offered benefits exceeding those of traditional spinal manipulation in chronic low back
 18 pain patients. However, this study has many limitations. The authors state that their pre-
 19 study analysis found that a sample size of 80 patients (half in each group) would be
 20 necessary to detect group differences similar to the differences they found, but their study
 21 was much smaller than this. In addition, patient selection protocols allowed patients to
 22 choose which therapy they would receive and all of those with the worst baseline pain
 23 chose to receive MUA. As such, the conclusions of this study cannot be taken to show that
 24 MUA is beneficial. Digiorgi (2013) states the evidence to support the efficacy of MUA of
 25 the spine remains largely anecdotal. There is a lack of high-quality evidence in the peer-
 26 reviewed medical literature of the effectiveness. Evidence of spinal manipulation under
 27 anesthesia consists primarily of case reports and uncontrolled case series. Limitations of
 28 current literature include small sample sizes, lack of random assignment, and limited
 29 evidence of long term benefit. Other issues include lack of detail regarding patient selection
 30 criteria, and differences in protocols reported in studies, making generalizations difficult.
 31 Guidelines from the American College of Occupational and Environmental Medicine
 32 (2007, 2008) and the Work Loss Data Institute (2011) state that spinal manipulation under
 33 anesthesia is not recommended. Colorado Division of Workers' Compensation's guidelines
 34 on "Low back pain medical treatment" (2014) did not recommend MUA.

35 36 **Shoulder**

37 In a Cochrane review, Green et al. (2000) examined the effectiveness of common
 38 interventions for shoulder pain. Intervention of interest included NSAIDs, intra-articular
 39 or subacromial glucocorticosteroid injection, oral glucocorticosteroid treatment,
 40 physiotherapy, MUA, hydrodilatation, or surgery. The authors concluded that there is little
 41 evidence to support or refute the effectiveness of common interventions for shoulder pain.
 42 They stated that there is a need for further well-designed clinical trials to establish a

1 uniform method of defining shoulder disorders. An updated review in 2007 was
2 withdrawn. A systematic review in BMJ Clinical Evidence (Speed, 2006) found that MUA
3 plus intra-articular injection is "likely to be beneficial" for persons with frozen shoulder.
4 The conclusions were based upon the results of two randomized controlled trials (RCTs).
5 One RCT ($n = 30$) found that, in people with adhesive capsulitis, MUA plus intra-articular
6 hydrocortisone injection increased recovery rates compared with intra-articular
7 hydrocortisone injection alone at 3 months (Thomas et al., 1980). Another, weaker RCT
8 ($n = 98$) found limited evidence that subjects having MUA plus intra-articular saline
9 injection versus manipulation alone or manipulation plus intra-articular injection of
10 methylprednisolone had greater improvements in range of motion (ROM), pain relief, and
11 return to normal activities (Hamdan and Al Essa, 2003). The review noted that potential
12 adverse effects of MUA of the shoulder include intra-articular lesions within the
13 glenohumeral joint (Speed, 2006).

14
15 Quraishi et al. (2007) assessed the outcome of MUA and hydrodilatation as treatments for
16 adhesive capsulitis. A total of 36 patients (38 shoulders) were randomized to receive either
17 method, with all patients being treated in stage II of the disease process. The visual analog
18 scale (VAS) in the hydro dilatation group were significantly better than those in the MUA
19 group over the 6-month follow-up period. The ROM improved in all patients over the 6
20 months but was not significantly different between the groups. At the final follow-up, 94%
21 of patients (17 of 18) were satisfied or very satisfied after hydrodilatation compared with
22 81% (13 of 16) of those who received MUA. Most patients were treated successfully, but
23 those undergoing hydrodilatation did better than those who underwent MUA. Kivimäki
24 and colleagues (2007) examined the effect of MUA in patients with frozen shoulder. A
25 blinded randomized trial with a 1-year follow-up was performed at 3 referral hospitals. A
26 total of 125 patients with clinically verified frozen shoulder were randomly assigned to the
27 manipulation group ($n = 65$) or control group ($n = 60$). Both the intervention group and the
28 control group were instructed in specific therapeutic exercises by physiotherapists. Clinical
29 data were gathered at baseline and at 6 weeks and 3, 6, and 12 months after randomization.
30 The two groups did not differ at any time of the follow-up in terms of shoulder pain or
31 working ability. Small differences in the ROM were detected favoring the manipulation
32 group. Perceived shoulder pain decreased during follow-up equally in the 2 groups, and at
33 1 year after randomization, only slight pain remained. Authors concluded that
34 manipulation under anesthesia does not add effectiveness to an exercise program
35 performed by patients.

36
37 Flannery et al. (2007) examined the influence of timing of MUA for adhesive capsulitis of
38 the shoulder on the long-term outcome. A total of 180 consecutive patients with a diagnosis
39 of adhesive capsulitis were selected from a shoulder surgery database; 145 were available
40 for follow-up after a mean period of 62 months (range of 12 to 125). All patients underwent
41 MUA with intra-articular steroid injection. A statistically significant improvement in range
42 of movement, function (Oxford Shoulder Score (OSS)) and VAS was obtained following

1 manipulation. Ninety percent of the 145 patients who successfully completed the study
 2 were satisfied with the procedure; 89% indicated that they would choose the same
 3 procedure again if the same problem arose in the opposite shoulder. Eighty-three percent
 4 of the patients had MUA performed less than 9 months from onset of symptoms (early
 5 MUA). The remainder had MUA performed after 9 to 40 months (late MUA). Patients who
 6 had early intervention had a significantly better OSS at final follow-up. There was no
 7 significant difference for mobility and pain. Theodorides et al. (2014) aimed to evaluate
 8 and determine the factors that affect short- and long-term outcome following MUA of
 9 patients with adhesive capsulitis. In total, 295 patients (315 shoulders) were sequentially
 10 recruited, and information was collected at baseline, as well as at a mean follow-up of 28
 11 days and 3.6 years. A significant improvement in OSS and ROM was noted 1 month post
 12 MUA with females benefiting more than males. Long-term follow-up revealed that the
 13 improvement in OSS was maintained. Secondary adhesive capsulitis significantly reduced
 14 the efficacy of MUA as assessed by ROM. Other factors (age, initial ROM and OSS, and
 15 length of symptoms prior to MUA) did not significantly affect the outcome over the short-
 16 or long-term. The findings of the present study show that all patient groups had a
 17 significantly improved ROM and OSS in the short-term with long-term maintenance of
 18 improved OSS. Woods and Loganathan (2017) aimed to address the issue of why not all
 19 patients benefit from MUA. Some have persistent or recurrent symptoms. There are no
 20 clear recommendations in the literature on the optimal management of recurrent frozen
 21 shoulder after a MUA. A total of 730 patients (792 shoulders) underwent MUA during the
 22 study period. A further MUA was undertaken in 141 shoulders (17.8%), for which we had
 23 complete data for 126. The mean improvement in OSS for all patients undergoing MUA
 24 was 16 (26 to 42), and the mean post-operative OSS in those requiring a further MUA was
 25 14 (28 to 42). Improvement was seen after a further MUA, regardless both of the outcome
 26 of the initial MUA, and of the time of recurrence. Patients with type-1 diabetes mellitus
 27 were at a 38% increased risk of requiring a further MUA, compared with the 18% increased
 28 risk of the group as a whole. Authors concluded that patients with a poor outcome or
 29 recurrent symptoms of a frozen shoulder after a MUA should be offered a further MUA
 30 with the expectation of a good outcome and a low complication rate.

31
 32 Rangan et al. (2020) compared these two surgical interventions with early structured
 33 physiotherapy plus steroid injection. In this multicenter, pragmatic, three-arm, superiority
 34 randomized trial, patients referred to secondary care for treatment of primary frozen
 35 shoulder were recruited from 35 hospital sites in the UK. Participants were adults (≥ 18
 36 years) with unilateral frozen shoulder, characterized by restriction of passive external
 37 rotation ($\geq 50\%$) in the affected shoulder. Participants were randomly assigned (2:2:1) to
 38 receive manipulation under anesthesia, arthroscopic capsular release, or early structured
 39 physiotherapy. Both forms of surgery were followed by postprocedural physiotherapy.
 40 Early structured physiotherapy involved mobilization techniques and a graduated home
 41 exercise program supplemented by a steroid injection. Both early structured physiotherapy
 42 and postprocedural physiotherapy involved 12 sessions during up to 12 weeks. The primary

1 outcome was the Oxford Shoulder Score (OSS). We sought a target difference of 5 OSS
2 points between physiotherapy and either form of surgery, or 4 points between manipulation
3 and capsular release. At 12 months, OSS data were available for 189 (94%) of 201
4 participants assigned to manipulation (mean estimate 38.3 points, 95% CI 36.9 to 39.7),
5 191 (94%) of 203 participants assigned to capsular release (40.3 points, 38.9 to 41.7), and
6 93 (94%) of 99 participants assigned to physiotherapy (37.2 points, 35.3 to 39.2). Eight
7 serious adverse events were reported with capsular release and two with manipulation.
8 Authors concluded that all mean differences on the assessment of shoulder pain and
9 function (OSS) at the primary endpoint of 12 months were less than the target differences.
10 Therefore, none of the three interventions were clinically superior. Arthroscopic capsular
11 release carried higher risks, and manipulation under anesthesia was the most cost-effective.
12

13 Brealey et al. (2020) compared the clinical effectiveness and cost-effectiveness of three
14 treatments in secondary care for adults with frozen shoulder; to qualitatively explore the
15 acceptability of these treatments to patients and health-care professionals; and to update a
16 systematic review to explore the trial findings in the context of existing evidence for the
17 three treatments. Participants were adults (aged ≥ 18 years) with unilateral frozen shoulder,
18 characterized by restriction of passive external rotation in the affected shoulder to $< 50\%$
19 of the opposite shoulder, and with plain radiographs excluding other pathology. The
20 interventions were early structured physiotherapy with a steroid injection, MUA with a
21 steroid injection and arthroscopic capsular release followed by manipulation. Post-
22 procedural physiotherapy followed both surgical interventions. The primary outcome and
23 end point was the Oxford Shoulder Score at 12 months post randomization. A difference
24 of five points was considered clinically important between early structured physiotherapy
25 and MUA or arthroscopic capsular release. Similarly, a four-point difference between
26 MUA and arthroscopic capsular release was considered significant. The mean age of the
27 503 participants was 54 years; 319 were female (63%) and 150 had diabetes (30%). The
28 primary analyses comprised 473 participants (94%). At the primary end point of 12
29 months, participants randomized to arthroscopic capsular release had, on average, a
30 statistically significantly higher (better) Oxford Shoulder Score than those randomized to
31 MUA or early structured physiotherapy. MUA did not result in statistically significantly
32 better Oxford Shoulder Score than early structured physiotherapy. No differences were
33 deemed of clinical importance. Serious adverse events were rare but occurred in
34 participants randomized to surgery (arthroscopic capsular release, $n = 8$; manipulation
35 under anesthesia, $n = 2$). Participants in the qualitative study wanted early medical help
36 and a quicker pathway to resolve their shoulder problem. Nine studies from the updated
37 systematic review, including UK FROST, of which only two could be pooled, and found
38 that arthroscopic capsular release was more effective than physiotherapy in the long-term
39 shoulder functioning of patients, but not to the clinically important magnitude used in UK
40 FROST. Authors concluded that none of the three interventions were clearly superior.
41 Early structured physiotherapy with a steroid injection is an accessible and low-cost option.

1 MUA is the most cost-effective option. Arthroscopic capsular release carries higher risks
2 and higher costs.

3
4 Song et al. (2021) aimed to evaluate the effect of MUA with intra-articular steroid injection
5 (ISI) or not on pain severity and function of the shoulder. Data on 141 patients receiving
6 MUA with primary frozen shoulder (FS) refractory to conservative treatments for at least
7 1 month were retrospectively obtained from medical records. Propensity score matching
8 analysis was performed between patients receiving MUA only and those receiving MUA
9 plus ISI, and then conducted logistic regression analysis to identify the risk factors for the
10 need to other treatments during 6-month follow-up. More improvement in terms of the
11 SPADI pain scores and passive ROM at 2 weeks after first intervention remained in
12 patients receiving MUA plus ISI after matching. The need to other treatments during 6-
13 month follow-up occurred in 10.6% patients ($n = 141$). Logistic regression analysis
14 revealed that a repeat MUA 1 week after first intervention was a protective factor and
15 duration of disease was the only one risk factor (OR 1.080; 95% CI 1.020-1.144; $P = .008$)
16 for the need to other treatments during follow-up. ISI immediately following MUA
17 provided additional benefits in rapid relief of pain and disability for patients with refractory
18 FS. Authors suggest that pain and disability of the shoulder may be rapidly alleviated by
19 an earlier MUA from the onset of the symptoms and a repeat MUA 1 week after first
20 intervention.

21
22 Rex et al. (2021) includes a recently completed multicenter randomized controlled trial
23 (RCT), UK FROST, in the context of existing randomized evidence for the management
24 of primary frozen shoulder in a systematic review. UK FROST compared the effectiveness
25 of pre-specified physiotherapy techniques with a steroid injection (PTSI), manipulation
26 under anesthesia (MUA) with a steroid injection, and arthroscopic capsular release (ACR).
27 This review updates a 2012 review focusing on the effectiveness of MUA, ACR, hydro
28 dilatation, and PTSI. Nine RCTs were included. The primary outcome of patient-reported
29 shoulder function at long-term follow-up (> 6 months and ≤ 12 months) was reported for
30 five treatment comparisons across four studies. Authors concluded that the findings from
31 a recent multicenter RCT provided the strongest evidence that, when compared with each
32 other, neither PTSI, MUA, nor ACR are clinically superior. Evidence from smaller RCTs
33 did not change this conclusion. The effectiveness of hydro dilatation based on four RCTs
34 was inconclusive and there remains an evidence gap.

35
36 Ko et al. (2021) aimed to assess how comorbidities influence the recovery speed and
37 clinical outcomes after MUA. Between April 2013 and September 2018, 281 consecutive
38 primary stiff shoulders in the frozen phase treated with MUA were included in this study.
39 They investigated the comorbidities of patients and divided them into the control ($n = 203$),
40 diabetes mellitus (DM) ($n = 32$), hyperlipidemia ($n = 26$), and thyroid disorder ($n = 20$)
41 groups. The range of motion (ROM) and clinical scores for each group before MUA and 1
42 week, 6 weeks, and 3 months after MUA were comparatively analyzed. They identified

1 the ROM recovery time after MUA and the responsiveness to MUA. Then, subjects were
2 subdivided into early and late recovery groups based on their recovery time and into
3 successful and unsuccessful MUA groups based on their responsiveness to MUA.
4 Significant improvements in ROM and clinical scores at 3 months after MUA were
5 observed in all groups. Significant differences in ROM among the 4 groups were also
6 observed during follow-up ($P < .05$). The DM group had significantly lower ROM values,
7 even at 3 months after MUA, compared with the control group. The ROM recovery speed
8 after MUA was slowest in the DM group, followed by the thyroid disorder, hyperlipidemia,
9 and control groups. Most (90.6%) of the DM group experienced late recovery. The
10 proportion of unsuccessful MUA was higher in the DM and thyroid disorder groups than
11 that in the control and hyperlipidemia groups ($P = .004$). During follow-up, there were no
12 differences among groups regarding the visual analog scale, University of California at
13 Los Angeles shoulder, and Constant scores. Authors concluded that the ROM recovery
14 speed and responsiveness to MUA for primary stiff shoulder were poorer for the DM and
15 thyroid disorder groups than for the control group. In particular, compared with any other
16 disease, outcomes were poorer when the comorbidity was DM. If patients have
17 comorbidities, then they should be informed before MUA that the comorbidity could affect
18 the outcomes of treatment.

19
20 Salomon et al. (2022) investigated the efficacy of manipulation under anesthesia (MUA)
21 compared to other non-surgical therapeutic strategies for patients with frozen shoulder
22 contracture syndrome (FSCS). Five randomized controlled trials were included. The
23 overall risk of bias (RoB) was high in 4 out of 5 of the included studies. MUA was found
24 to be not superior in terms of reduction of pain and improvement of function when
25 compared to cortisone injections with hydrodilatation and home exercise in the short term
26 (3 months), and cortisone injections with hydrodilatation in the long term (≥ 6 months).
27 Moreover, if compared to structured physiotherapy, MUA highlighted a higher Oxford
28 Shoulder Score at final 1-year follow up. Similar results were obtained for disability, with
29 statistically no significant long-term differences between MUA and home exercise or
30 structured physiotherapy. Only two trials reported adverse events. This review suggested
31 that limited and inconsistent evidence currently exists on the efficacy of MUA compared
32 to other non-surgical strategies in the management of patients with FSCS. Future research
33 should focus on clinical trials with higher methodological quality.

34
35 Kraal et al. (2023) evaluated the effectiveness of MUA followed by a physiotherapy (PT)
36 program compared to a PT program alone in patients with stage 2 Frozen Shoulder (FS).
37 Frozen shoulder (FS) is a common cause of shoulder pain and stiffness. Conservative
38 treatment is sufficient for the majority of patients with long-term recovery of shoulder
39 function. Manipulation under anesthesia (MUA) is known as a well-established treatment
40 option if conservative treatment fails. It is unknown whether MUA does indeed shorten the
41 duration of symptoms or leads to a superior outcome compared to conservative treatment.

1 For this study, patients between 18 and 70 years old with stage 2 FS were deemed eligible
2 if an initial course of conservative treatment consisting of PT and intra-articular
3 corticosteroid infiltration was considered unsatisfactory. MUA was performed by a single
4 surgeon under interscalene block, and intensive PT treatment protocol was started within
5 4 hours after MUA. In the PT group, patients were referred to instructed physiotherapist,
6 and treatment was guided by tissue irritability. The primary outcome was the Shoulder
7 Pain and Disability Index (SPADI) score. Secondary outcomes were pain, range of motion
8 (ROM), Oxford Shoulder Score, quality of life, and ability to work. In total, 82 patients
9 were included, 42 in the PT group and 40 in the MUA group. There was a significant
10 improvement in SPADI, Oxford Shoulder Score, pain, ROM, and quality of life in both
11 groups at 1-year follow-up. SPADI scores at three months were significantly improved in
12 favor of MUA. MUA showed a significantly bigger increase in flexion and abduction
13 compared to PT at all points of follow-up. No significant differences between both groups
14 were found for all other parameters. No fractures, dislocations, or brachial plexus injuries
15 occurred in this trial. Authors concluded that MUA in stage 2 FS can be considered safe
16 and results in a faster recovery of ROM and improved functional outcome, measured with
17 SPADI scores, compared to PT alone in the short term. After 1 year, except for slightly
18 better ROM scores for MUA, the result of MUA is equal to PT.

19
20 Mullen et al. (2025) authored a paper on adhesive capsulitis. They note that nonsurgical
21 treatment focuses on pain relief and restoring ROM and includes physical therapy (PT),
22 oral anti-inflammatory medications, corticosteroid injections, extracorporeal shock wave
23 therapy, and ultrasonography-guided hydrodistention. Early corticosteroid injections are
24 associated with shortened symptom duration and improved functional scores. Ultrasound-
25 guided hydrodilatation with hyaluronic acid combined with PT has also shown superior
26 outcomes compared with PT alone. For patients with diabetes, extracorporeal shock wave
27 therapy avoids potential metabolic complications from steroids. Adhesive capsulitis is self-
28 limiting, with most patients achieving symptom resolution without surgery. Surgical
29 intervention, typically considered after 9 to 12 months of failed nonsurgical management,
30 includes arthroscopic capsular release, manipulation under anesthesia (MUA), or both.
31 Both approaches are efficacious, with improved pain, ROM, and functional scores.
32 Although MUA may be more cost-effective, arthroscopy allows direct visualization and
33 treatment of pathology while reducing the risk of complications from MUA, such as
34 fracture, dislocation, or nerve injury.

35 36 **Knee**

37 MUA is indicated, with or without arthroscopy for arthrofibrosis of the knee (i.e., post
38 ACL reconstruction), when there is <90° range of motion following surgery or trauma
39 despite physical therapy (Magit et al., 2007). Manipulation under anesthesia has also been
40 used to treat fibroarthrosis following total knee replacement. Following total knee
41 arthroplasty, some patients who fail to achieve greater than 90 degrees of flexion in the
42 early peri-operative period may be considered candidates for MUA of the knee.

1 Manipulation under anesthesia is indicated in total knee arthroplasty having less than 90
2 degrees ROM 4 to 12 weeks following surgery, with no progression or regression in ROM
3 (Pariante et al., 2006; Magit et al., 2007). Keating et al. (2007) assessed the outcomes of
4 manipulation following total knee arthroplasty. A total of 113 knees in 90 patients
5 underwent manipulation for post-operative flexion of less than or equal to 90 degrees at a
6 mean of 10 weeks after surgery. Eighty-one (90%) of the 90 patients achieved
7 improvement of ultimate knee flexion following manipulation. The average improvement
8 in flexion from the measurement made before manipulation to that recorded at the 5-year
9 follow-up was 35 degrees. The investigators reported that there was no significant
10 difference in the mean improvement in flexion when patients who had manipulation within
11 12 weeks post-operatively were compared with those who had manipulation more than 12
12 weeks post-operatively. Patients who eventually underwent manipulation had significantly
13 more pain than those who had not had manipulation. The investigators concluded that
14 manipulation generally increases final flexion following total knee arthroplasty. They
15 noted that patients with severe pre-operative pain are more likely to require manipulation.

16
17 Sassoon et al. (2015) investigated the results of closed manipulations performed under
18 anesthesia (MUA) to evaluate whether it is an effective means to treat posttraumatic knee
19 arthrofibrosis. Twenty-two patients with a mean age of 40 underwent closed MUA for
20 posttraumatic knee arthrofibrosis. Injuries included fractures of the femur, tibia, and patella
21 as well as ligamentous injuries and traumatic arthrotomies. The mean time from treatment
22 to manipulation was 90 days. Mean follow-up after manipulation was 7 months. The mean
23 premanipulation ROM arc was 59 ± 25 degrees. The mean intraoperative arc of motion,
24 achieved at the time of the manipulation was 123 ± 14 degrees. No complications occurred
25 during the MUA procedure. At the most recent follow-up, the mean ROM arc was $110 \pm$
26 19 degrees. Tobacco use, associated injuries, elevated body mass index, open fracture, and
27 advanced age did not impact manipulation efficacy. Additionally, manipulations
28 performed 90 days or more after surgical treatment provided a benefit equaling those
29 performed more acutely. Authors concluded that MUA is a safe and effective method to
30 increase knee ROM in the setting of posttraumatic arthrofibrosis. Improvement in ROM
31 was noted in all patients.

32
33 Ekhtiari et al. (2017) reviewed the literature to: (a) describe existing definitions of
34 arthrofibrosis, and (b) characterize the management strategies and outcomes of
35 arthrofibrosis treatment in patients post ACL reconstruction. Twenty-five studies of
36 primarily level IV evidence (88%) were included. A total of 647 patients with a mean age
37 of 28.2 ± 1.8 years (range 14-62 years) were treated for arthrofibrosis following ACL
38 reconstruction and followed for a mean 30.1 ± 16.9 months (range 2 months-9.6 years).
39 Definitions of arthrofibrosis varied widely and included subjective definitions and the
40 Shelbourne classification system. Patients were treated by one or more of: arthroscopic
41 arthrolysis (570 patients), MUA (153 patients), oral corticosteroids (31 patients),
42 physiotherapy (81 patients), drop-casting (17 patients), epidural therapy combined with

1 inpatient physiotherapy (6 patients), and intra-articular interleukin-1 antagonist injection
2 (4 patients). All studies reported improvement in range of motion post-operatively, with
3 statistically significant improvement reported for 306 patients (6 studies, p range <0.001
4 to =0.05), and one study (18 patients) reporting significantly better results if arthrofibrosis
5 was treated within 8 months of reconstruction ($p < 0.03$). The greatest improvements for
6 extension loss were seen with drop-casting (mean $6.2^\circ \pm 0.6^\circ$ improvement), whereas
7 MUA produced the greatest improvement for flexion deficit (mean $47.8^\circ \pm 3.3^\circ$
8 improvement). Gu et al. (2018) performed a systematic review of the literature was
9 performed to identify studies that reported clinical outcomes for patients who underwent
10 MUA for post-operative stiffness treatment. Repeat MUA procedures were included in the
11 study but were analyzed separately. Twenty-two studies (1,488 patients) reported on ROM
12 after MUA, and 4 studies (81 patients) reported ROM after repeat MUA. All studies
13 reported pre-MUA motion of less than 90° , while mean ROM at last follow-up exceeded
14 90° in all studies except two. For studies reporting ROM improvement following repeat
15 MUA, the mean pre-manipulation ROM was 80° and the mean post-manipulation ROM
16 was 100.6° .

17
18 Authors concluded that MUA remains an efficacious, minimally invasive treatment option
19 for post-operative stiffness following total knee arthroplasty (TKA). MUA provides
20 clinically significant improvement in ROM for most patients, with the best outcomes
21 occurring in patients treated within 12 weeks post-operatively. Neuman et al. (2018)
22 completed a study on risk factors, outcomes, and timing of MUA after TKA. Clinical
23 variables were compared between patients who underwent MUA and those who did not;
24 variables that differed were utilized to identify an appropriately matched control group of
25 non-MUA patients. The MUA group was divided into early (MUA ≤ 6 weeks from index)
26 and late (>6 weeks) subgroups. Flexion values at multiple time points were compared. In
27 total, 1,729 TKA patients were reviewed; MUA was performed in 62 patients. TKA
28 patients undergoing MUAs were younger, more likely to be current smokers, and more
29 likely to have undergone prior knee surgery. Even in patients with severe initial
30 postoperative limitations in ROM, MUA within 6 weeks may allow for final outcomes that
31 are equivalent to those experienced by similar patients not requiring manipulation.

32
33 Archunan et al. (2021) aimed to ascertain the prevalence, determine the influencing factors,
34 and evaluate the efficacy of MUA as a treatment option. For the study, stiffness was
35 defined as flexion contracture of >15 degrees and/or flexion of <75 degrees. Demographic
36 data included co-morbidities, previous knee surgery, pre-operative and post-operative
37 ROM, anesthetic techniques and use of nerve blocks, type of prosthesis, ligament balancing
38 including release, mobility post-surgery, patient motivation, physiotherapy, complications,
39 and final ROM post-MUA. Of the 1,350 patients evaluated, 33 (2.44%) had stiffness
40 defined by the above-outlined criteria and required intervention. Thirty-one patients
41 underwent MUA as a first-line treatment. No complications arose following MUA. One
42 patient (0.07%) required arthroscopic arthrolysis while another patient (0.07%) required

1 revision arthroplasty due to patellar maltracking. Following manipulation, mean flexion
2 contracture decreased from 8 degrees to 3.6 degrees, and mean flexion improved from 51.8
3 degrees to 93.2 degrees. Arc of motion improved in 100% of patients but it is important to
4 note that multiple manipulations were performed in seven patients. Authors concluded that
5 stiffness after TKA can be difficult to treat and can result in prolonged morbidity and
6 dissatisfaction. This retrospective study highlights the effectiveness of MUA as a first-line
7 treatment option leading to improved outcomes especially if done early.

8
9 Sala et al. (2022) completed a retrospective study determined the outcome of MUA and
10 identified the factors affecting it. The final sample consisted of 150 MUAs performed on
11 145 patients. The parameters of interest were ROM and Knee Society Score (KSS) or
12 Oxford Knee Score (OKS). The mean of 26° gain in flexion and the mean of 3° gain in
13 extension were noticed at post-MUA follow-up when compared with the ROM preceding
14 MUA. The mean post-MUA-FU flexion was 99° and the mean post-MUA-FU extension
15 deficit was 4°. KSS (121 vs. 129) and OKS (29 vs. 28) were similar before and after MUA.
16 The early timing of MUA was associated with better gain in flexion -0.04, while we found
17 no association between the timing of MUA and flexion after MUA -0.004. High BMI was
18 associated with better gain in flexion 0.8. Authors found that ROM improved substantially
19 after MUA. The gain in flexion decreased as the time between TKA and MUA increased.
20 DeFrance et al. (2022) sought to determine whether MUA was associated with an increase
21 in the rate of revision TKA within 2 years of MUA. A total of 49,310 patients within a
22 single institution who underwent primary TKA were identified from 1999 to 2019. Data
23 were matched at a 1:3 ratio (TKA with and without MUA, respectively) based on age, sex,
24 and body mass index. A matched comparison cohort was conducted, with the MUA cohort
25 having 575 patients and the no MUA cohort having 1,725 patients. A statistically
26 significant increase in the rate of noninfectious etiology revision TKA was found in the
27 MUA cohort (7.3%) compared with the no MUA cohort (4.9%; P=.034). The most
28 common reason for revision TKA after MUA was persistent stiffness, including
29 arthrofibrosis and ankylosis; however, aseptic loosening, ligamentous instability, and
30 periprosthetic fracture were found to be responsible for 21.4% of revision TKA procedures.
31 Although MUA is a commonly performed procedure for treating stiffness after primary
32 TKA, the orthopedic surgeon should counsel patients on the association of increased rate
33 of revision TKA after MUA, most commonly, persistent stiffness.

34
35 Haffar et al. (2022) performed a systematic review to compare the outcomes of
36 manipulation under anesthesia (MUA), arthroscopic lysis of adhesions (aLOA), and
37 revision TKA (rTKA) for arthrofibrosis and stiffness following TKA. A total of 40 studies
38 were included: 21 on rTKA, 7 on aLOA, and 14 on MUA. The mean or median post-
39 operative arc ROM was > 90° in 6/20 (30%) rTKA, 5/7 (71%) aLOA, and 7/10 (70%)
40 MUA studies. Post-operative Knee Society (KSS) clinical and functional scores were the
41 greatest in patients who underwent MUA and aLOA. As many as 43% of rTKA patients
42 required further care compared to 25% of aLOA and 17% of MUA patients. Authors

1 concluded that stiffness following TKA remains a challenging condition to treat.
2 Nonetheless, current evidence suggests that patients who undergo rTKA have poorer
3 clinical outcomes and a greater need for further treatment compared to patients who
4 undergo MUA or aLOA.

5
6 Thomas et al. (2023) compared the 2-year complication rates of arthroscopic lysis of
7 adhesions (ALA) and MUA and range-of-motion (ROM) outcomes for ALA, early MUA
8 (<3 months after TKA), and delayed MUA (>3 months after TKA). This retrospective
9 cohort study included 425 patients undergoing ALA or MUA after primary TKA from
10 2001 to 2018. Demographics, clinical variables, and complication rates were collected
11 from clinical records. ALA patients were younger (55.2 versus 58.9 years, $P < 0.001$) and
12 underwent surgery later from the index TKA (12 versus 1.9 months, $P < 0.001$). The
13 Charlson Comorbidity Index was higher in the MUA group. Preoperative ROM was
14 significantly worse in the MUA cohort but did not differ between groups after the
15 procedure or at 2 years. Demographics and ROM outcomes were equivalent between early
16 MUA and delayed MUA. The incidence of repeat arthrofibrosis (7.1%) and revision
17 arthroplasty (2.4%) was similar between ALA and MUA cohorts while ALA patients had
18 significantly more surgical site infections (3.8%) compared with MUA patients (0.47%, P
19 = 0.017). Equivalent ROM outcomes were seen between ALA, early MUA, and delayed
20 MUA for the treatment of arthrofibrosis after TKA. However, this study demonstrated a
21 markedly higher complication rate, particularly surgical site infection, after ALA,
22 suggesting that MUA may be the preferred option for treating arthrofibrosis at both early
23 and late time points.

24
25 Akhtar et al. (2024) evaluated the functional and clinical outcomes of early versus delayed
26 MUA for stiffness following TKA. Stiffness following TKA is often treated with MUA.
27 However, there is debate regarding the timing of MUA, with many recommending against
28 MUA beyond 3 months after TKA. Included were 14 studies analyzing 13,445 knees,
29 72.1% of which underwent early MUA and 27.8% of which underwent delayed MUA. Of
30 the 14 studies, 10 defined early MUA as being performed within 3 months of the index
31 TKA. Pre-MUA and post-MUA knee flexion for the early/delayed groups was $71.3^\circ/77.9^\circ$
32 and $103.0^\circ/96.1^\circ$, respectively. Upon meta-analysis, pre-MUA knee flexion was
33 significantly higher in the delayed group, whereas post-MUA flexion was similar in both
34 groups. The mean gain in knee flexion for the early and delayed groups was $32.0^\circ/19.2^\circ$.
35 The surgical complication and revision TKA rates for the early and delayed groups were
36 4.9%/10.3% and 5%/9%, respectively. A meta-analysis found the risk of surgical or
37 medical complications and revision TKA to be significantly higher in the delayed MUA
38 group. Authors concluded that although post-MUA knee flexion was similar in patients
39 undergoing early and delayed MUA following TKA, the mean gain in flexion for early
40 patients was nearly double that of delayed patients. Delayed patients also had significantly
41 higher risks of surgical or medical complications and revision TKA following MUA.

1 Brown et al. (2024) sought to determine whether MUA had any advantage over routine
2 care in the treatment of patients who developed arthrofibrosis following TKA.
3 Arthrofibrosis is a multifactorial process that results in decreased knee range of motion
4 (ROM). Manipulation under anesthesia (MUA) is commonly regarded as the preferred
5 initial treatment of arthrofibrosis following total knee arthroplasty (TKA). There have been
6 no well-controlled studies demonstrating that MUA effectively increases ROM in patients
7 who develop arthrofibrosis after TKA when compared with routine care. The authors
8 identified patients who underwent primary TKA at the authors' institution between 2010
9 and 2014 and had flexion ≤ 100 degrees at early follow-up. Knees were grouped based on
10 how the arthrofibrosis was treated: those who underwent MUA and those who received
11 routine care. Knee flexion was captured preoperatively (prior to TKA), at early follow-up
12 (prior to MUA or routine care), and at 1-year follow up. Flexion change from early follow-
13 up to 1 year was calculated. The average flexion at 1-year follow-up was not significantly
14 different between the two groups (106.1 ± 11.7 degrees in the routine care group versus
15 106.3 ± 12.8 degrees in the MUA group). The MUA group had a greater proportion of
16 patients with flexion gains > 20 degrees at final follow-up when compared with patients
17 who underwent routine care (56% vs. 8%, $p < 0.0001$). This study demonstrates that
18 patients with decreased ROM at early follow-up after primary TKA can expect greater
19 ROM increase at 1-year follow-up if they undergo MUA compared with patients who
20 undergo routine care.

21
22 Marquez-Lara et al. (2024) evaluated the safety and efficacy of early (<3 mo
23 postoperatively) MUA for the treatment of knee arthrofibrosis in adolescent patients.
24 Authors hypothesized that early MUA could restore normal knee motion with a low
25 complication rate and without the need for more invasive intervention. In a retrospective
26 review, 57 patients who underwent MUA for postoperative knee arthrofibrosis were
27 identified. The median age of the cohort at time of MUA was 14.5 years. 54.4% were male.
28 Median time to MUA was 64 days after index surgery. ROM before MUA was 90.0
29 degrees, which improved to 130 degrees (120 to 135) after MUA. At final median follow-
30 up of 8.9 months, mean ROM was 133 degrees (130 to 140). There were no iatrogenic
31 fractures or physeal separations associated with MUA. 12.3% ($n=7/57$) failed MUA either
32 due to the need for subsequent repeat MUA ($n=2$), need for lysis of adhesions ($n=3$) or
33 need for surgery after MUA ($n=2$). Those who failed early MUA and required subsequent
34 procedures had ROM >120 degrees at final follow-up. Authors concluded that
35 postoperative knee arthrofibrosis can be safely and effectively treated with early (<3 mo
36 postoperative) MUA. There were no iatrogenic fractures or physeal separations during
37 MUA. Patients who had recurrence of motion deficits after early MUA and required further
38 intervention, regained satisfactory knee motion at final follow-up. Although further
39 research is warranted to better characterize risk factors for knee arthrofibrosis in adolescent
40 patients, early recognition and MUA is a safe and effective treatment for arthrofibrosis to
41 help patients regain full ROM without invasive intervention.

Fracture and/or Dislocation

MUA is also considered a well-established and successful treatment for some types of fractures (e.g., vertebral, long bones) and acute/traumatic dislocations (e.g., perched cervical facet). It is typically performed with surgical repair and other medically necessary procedures such as arthroscopy. When performed in this context, MUA is considered incidental to the base procedure.

Chronic Joint Contracture

A joint contracture is a limitation in the passive range of motion of a joint. Joint contractures prevent normal movement of the associated body part and can result from a variety of causes such as spasticity or prolonged immobilization. Intra-articular adhesions and peri-articular adhesions, as well as capsular, ligament and muscle shortening, and tightness may develop. As a result, activities of daily living and other functions may be adversely affected due to the decreased mobility. In many cases, contractures can be successfully treated nonoperatively with aggressive physical therapy or splinting with restoration of functional range of motion. When conservative treatment fails more aggressive treatment may necessary and includes anesthetic block, maximal stretching, and in some cases, serial casting (Garden, 2002). For joint contracture deformities, extra-articular and intra-articular soft tissue releases are considered standard treatment (Paley, 2003). Surgical treatments include tenotomy, tendon lengthening and joint capsule release. Manipulation under anesthesia, involving maximal passive stretching may be considered standard treatment and is often performed in combination with serial casting and/or surgical release when less aggressive treatments have failed.

Elbow

Published peer reviewed supporting the safety and effectiveness of using manipulation under anesthesia of the elbow is limited to retrospective case series, involve small sample populations and lack control groups (Araghi et al., 2012; Tan. Et al., 2006; Chao et al., 2002; Gaur et al., 2003). Few studies support clinical effectiveness for the treatment of joint stiffness/fibrosis when other conservative measures, such as bracing and splinting, have failed to improve range of motion. There is insufficient evidence in the peer-reviewed published literature and lack of consensus among professional societies to support the effectiveness of MUA as treatment for arthrofibrosis of the elbow. Spitler et al. (2018) evaluated the safety and efficacy of manipulation under anesthesia (MUA) for posttraumatic elbow stiffness. Comparison of improvement between the early and late MUA groups found a significant difference ($P < 0.001$) in mean flexion arc improvement from premanipulation to postmanipulation, favoring the early group. Authors concluded that MUA is a safe and effective adjunct to improving motion in posttraumatic elbow stiffness when used within 3 months from the original injury or time of surgical fixation. After 3 months, MUA does not reliably increase elbow motion.

TMJ

Available evidence for MUA for temporomandibular joint syndrome is limited to small, uncontrolled studies with limited follow-up. Foster et al. (2000) conducted an uncontrolled prospective study of manipulation of the temporomandibular joint under anesthesia. The investigators reported that of the 55 patients available for participation in this study, 15 improved, 15 did not, 6 showed partial improvement, and 19 were not treated. The median pre-treatment opening was 20 mm (range of 13 to 27). Among those who improved after manipulation, the median opening after treatment was 38 mm (range of 35 to 56). The investigators reported that some of those who improved experienced a return of TMJ clicking but not of joint or muscle tenderness. There is insufficient evidence in the peer-reviewed published literature to support the effectiveness of MUA as treatment for TMJ syndrome.

Other Joints and Conditions

Evidence in the medical literature evaluating the use of MUA for management of pain conditions involving one or more (i.e., multiple joints, whole body MUA) of other major joints such as the hip, ankle, toe, elbow, and wrist, is lacking. Due to insufficient evidence conclusions cannot be made regarding the clinical utility or safety and efficacy of MUA involving other single or multiple joints for pain management. There is a paucity of evidence supporting the use of MUA for the treatment of disorders of other body joints such as the hip, ankle, knee, and wrist.

PRACTITIONER SCOPE AND TRAINING

Practitioners should practice only in the areas in which they are competent based on their education, training, and experience. Levels of education, experience, and proficiency may vary among individual practitioners. It is ethically and legally incumbent on a practitioner to determine where they have the knowledge and skills necessary to perform such services and whether the services are within their scope of practice.

It is best practice for the practitioner to appropriately render services to a member only if they are trained, equally skilled, and adequately competent to deliver a service compared to others trained to perform the same procedure. If the service would be most competently delivered by another health care practitioner who has more skill and training, it would be best practice to refer the member to the more expert practitioner.

Best practice can be defined as a clinical, scientific, or professional technique, method, or process that is typically evidence-based and consensus driven and is recognized by a majority of professionals in a particular field as more effective at delivering a particular outcome than any other practice (Joint Commission International Accreditation Standards for Hospitals, 2020).

1 Depending on the practitioner’s scope of practice, training, and experience, a member’s
 2 condition and/or symptoms during examination or the course of treatment may indicate the
 3 need for referral to another practitioner or even emergency care. In such cases it is prudent
 4 for the practitioner to refer the member for appropriate co-management (e.g., to their
 5 primary care physician) or if immediate emergency care is warranted, to contact 911 as
 6 appropriate. See the *Managing Medical Emergencies (CPG 159 – S)* policy for
 7 information.

9 **References**

- 10 Akhtar M, Razick D, Seibel A, Asad S, Shekhar A, Shelton T. Outcomes of Early Versus
 11 Delayed Manipulation Under Anesthesia for Stiffness Following Total Knee
 12 Arthroplasty: A Systematic Review and Meta-Analysis. *J Arthroplasty*.
 13 2024;39(11):2872-2879. doi:10.1016/j.arth.2024.05.059
- 14
- 15 Araghi A, Celli A, Adams R, Morrey B. The outcome of examination (manipulation) under
 16 anesthesia on the stiff elbow after surgical contracture release. *Shoulder Elbow Surg*.
 17 2010 Mar;19(2):202-8
- 18
- 19 Archunan M, Swamy G, Ramasamy A. Stiffness After Total Knee Arthroplasty:
 20 Prevalence and Treatment Outcome. *Cureus*. 2021;13(9):e18271. Published 2021
 21 Sep 25
- 22
- 23 Aspegren, D. D., Wright, R. E., & Hemler, D. E. (1997). Manipulation under epidural
 24 anesthesia with corticosteroid injection: two case reports. *Journal of Manipulative and*
 25 *Physiological Therapeutics*, 20(9), 618-621
- 26
- 27 Ben-David, B., & Raboy, M. (1994). Manipulation under anesthesia combined with
 28 epidural steroid injection. *Journal of Manipulative and Physiological Therapeutics*,
 29 17(9), 605-609
- 30
- 31 Brealey S, Northgraves M, Kottam L, Keding A, Corbacho B, Goodchild L, Srikesavan C,
 32 Rex S, Charalambous CP, Hanchard N, Armstrong A, Brooksbank A, Carr A, Cooper
 33 C, Dias J, Donnelly I, Hewitt C, Lamb SE, McDaid C, Richardson G, Rodgers S, Sharp
 34 E, Spencer S, Torgerson D, Toye F, Rangan A. Surgical treatments compared with
 35 early structured physiotherapy in secondary care for adults with primary frozen
 36 shoulder: the UK FROST three-arm RCT. *Health Technol Assess*. 2020 Dec;24(71):1-
 37 162. Doi: 10.3310/hta24710. PMID: 33292924; PMCID: PMC7750869
- 38
- 39 Brown ML, Vaz KM, McCauley JC, May L, Colwell CW Jr. Who Benefits from
 40 Manipulation under Anesthesia Following Total Knee Arthroplasty?. *J Surg Orthop*
 41 *Adv*. 2024;33(1):33-36

- 1 Buchalter D, Schaffler BC, Manjunath A, et al. Stiffness After Total Knee Arthroplasty A
2 Review. *Bull Hosp Jt Dis* (2013). 2024;82(1):15-20
3
- 4 Chao EK, Chen AC, Lee MS, Ueng SW. Surgical approaches for nonneurogenic elbow
5 heterotopic ossification with ulnar neuropathy. *J Trauma*. 2002 Nov;53(5):928-33
6
- 7 Cremata, E., Collins, S., Clauson, W., Solinger, A. B., & Roberts, E. S. (2005).
8 Manipulation under anesthesia: a report of four cases. *Journal of Manipulative and*
9 *Physiological Therapeutics*, 28(7), 526-533
10
- 11 DeFrance MJ, Cheesman QT, Hameed D, DiCiurcio WT, Harrer MF. Manipulation Under
12 Anesthesia Is Associated With an Increased Rate of Early Total Knee Arthroplasty
13 Revision. *Orthopedics*. 2022;45(5):270-275. Doi:10.3928/01477447-20220608-01
14
- 15 Digiorgi D. Spinal manipulation under anesthesia: a narrative review of the literature and
16 commentary. *Chiropr Man Therap*. 2013 May 14;21(1):14
17
- 18 Dodenhoff RM, Levy O, Wilson A, Copeland SA. Manipulation under anesthesia for
19 primary frozen shoulder: effect on early recovery and return to activity. *J Shoulder*
20 *Elbow Surg*. 2000;9:23–6
21
- 22 Dreyfuss, P., Michaelsen, M., & Horne, M. (1995). MUJA: manipulation under joint
23 anesthesia/analgesia: a treatment approach for recalcitrant low back pain of synovial
24 joint origin. *Journal of Manipulative and Physiological Therapeutics*, 18(8), 537-546
25
- 26 ECRI. (2003). Manipulation Under Anesthesia for Low-Back Pain. *Health Technology*
27 *Assessment Information Service: Windows on Medical Technology*, 1-33
28
- 29 Ekhtiari S, Horner NS, de Sa D, et al. Arthrofibrosis after ACL reconstruction is best
30 treated in a step-wise approach with early recognition and intervention: a systematic
31 review. *Knee Surg Sports Traumatol Arthrosc*. 2017;25(12):3929-3937
32
- 33 Fitzsimmons SE, Vazquez EA, Bronson MJ. How to treat the stiff total knee arthroplasty?:
34 a systematic review. *Clin Orthop Relat Res*. 2010 Apr;468(4):1096-106
35
- 36 Flannery O, Mullett H, Colville J. Adhesive shoulder capsulitis: Does the timing of
37 manipulation influence outcome? *Acta Orthop Belg*. 2007;73(1):21-25
38
- 39 Gaur A, Sinclair M, Caruso E, Peretti G, Zaleske D. Heterotopic ossification around the
40 elbow following burns in children: results after excision. *J Bone Joint Surg Am*. 2003
41 Aug;85-A(8):1538-43

- 1 Ghani H, Maffulli N, Khanduja V. Management of stiffness following total knee
2 arthroplasty: A systematic review. *Knee*. 2012 Apr 23
3
- 4 Gordon, R. C. (2001). An evaluation of the experimental and investigational status and
5 clinical validity of manipulation of patients under anesthesia: a contemporary opinion.
6 *Journal of Manipulative and Physiological Therapeutics*, 24(9), 603-611
7
- 8 Gordon R, Cremata E, Hawk C. Guidelines for the practice and performance of
9 manipulation under anesthesia. *Chiropr Man Therap*. 2014 Feb 3;22(1):7
10
- 11 Green S, Buchbinder R, Glazier R, Forbes A. Interventions for shoulder pain. *Cochrane*
12 *Database Syst Rev*. 2000;(2):CD001156. Review. Update in: *Cochrane Database Syst*
13 *Rev*. 2006;(4):CD001156
14
- 15 Greenman, P. E. (1992). Manipulation with the patient under anesthesia. *The Journal of*
16 *the American Osteopathic Association*, 92(9), 1159-1160, 1167-1170
17
- 18 Gu A, Michalak AJ, Cohen JS, Almeida ND, McLawhorn AS, Sculco PK. Efficacy of
19 Manipulation Under Anesthesia for Stiffness Following Total Knee Arthroplasty: A
20 Systematic Review. *J Arthroplasty*. 2018 May;33(5):1598-1605
21
- 22 Haffar A, Goh GS, Fillingham YA, Torchia MT, Lonner JH. Treatment of arthrofibrosis
23 and stiffness after total knee arthroplasty: an updated review of the literature. *Int*
24 *Orthop*. 2022;46(6):1253-1279. Doi:10.1007/s00264-022-05344-x
25
- 26 Haldeman, S., Chapman-Smith, D., & Petersen, D., Jr. (1993). *Guidelines for Chiropractic*
27 *Quality Assurance and Practice Parameters: Proceedings of the Mercy Center*
28 *Consensus Conference*
29
- 30 Hamdan TA, Al-Essa KA. Manipulation under anaesthesia for the treatment of frozen
31 shoulder. *Int Orthop*. 2003;27(2):107-9. Epub 2002 Sep 13
32
- 33 Herzog, J. (1999). Use of cervical spine manipulation under anesthesia for management of
34 cervical disk herniation, cervical radiculopathy, and associated cervicogenic headache
35 syndrome. *Journal of Manipulative and Physiological Therapeutics*, 22(3), 166-170
36
- 37 Hughes, B. L. (1993). Management of cervical disk syndrome utilizing manipulation under
38 anesthesia. *Journal of Manipulative and Physiological Therapeutics*, 16(3), 174-181
39
- 40 Hyman, S. A., Rogers, W. D., & Bullington, J. C., 3rd. (1990). Cervical osteotomy and
41 manipulation in ankylosing spondylitis: successful general anesthesia after failed local
42 anesthesia with sedation. *Journal of Spinal Disorders*, 3(4), 423-426

- 1 Ipach I, Mittag F, Lahrmann J, Kunze B, Kluba T. Arthrofibrosis after TKA - Influence
 2 factors on the absolute flexion and gain in flexion after manipulation under anaesthesia.
 3 BMC Musculoskelet Disord. 2011 Aug 12;12:184
 4
- 5 Issa K, Banerjee S, Kester MA, Khanuja HS, Delanois RE, Mont MA. The effect of timing
 6 of manipulation under anesthesia to improve range of motion and functional outcomes
 7 following total knee arthroplasty. J Bone Joint Surg Am. 2014 Aug 20;96(16):1349-57
 8
- 9 Issa K, Kapadia BH, Kester M, Khanuja HS, Delanois RE, Mont MA. Clinical, objective,
 10 and functional outcomes of manipulation under anesthesia to treat knee stiffness
 11 following total knee arthroplasty. J Arthroplasty. 2014 Mar;29(3):548-52
 12
- 13 Keating EM, Ritter MA, Harty LD, Haas G, Meding JB, Faris PM, Berend ME.
 14 Manipulation after total knee arthroplasty. J Bone Joint Surg Am. 2007 Feb;89(2):282-
 15 6
 16
- 17 Kivimäki J, Pohjolainen T, Malmivaara A, et al. Manipulation under anesthesia with home
 18 exercises versus home exercises alone in the treatment of frozen shoulder: A
 19 randomized, controlled trial with 125 patients. J Shoulder Elbow Surg.
 20 2007;16(6):722-726
 21
- 22 Ko YW, Park JH, Youn SM, Rhee YG, Rhee SM. Effects of comorbidities on the outcomes
 23 of manipulation under anesthesia for primary stiff shoulder. J Shoulder Elbow Surg.
 24 2021;30(8):e482-e492
 25
- 26 Kohlbeck, F. J., Haldeman, S., Hurwitz, E. L., & Dagenais, S. (2005). Supplemental care
 27 with medication-assisted manipulation versus spinal manipulation therapy alone for
 28 patients with chronic low back pain. *Journal of Manipulative and Physiological*
 29 *Therapeutics*, 28(4), 245-252
 30
- 31 Kraal T, de Wit Y, The B, et al. Improved range of motion after manipulation under
 32 anesthesia versus physiotherapy for stage two frozen shoulder: a randomized controlled
 33 trial. JSES Int. 2023;8(2):293-298. Published 2023 Dec 6
 34
- 35 Lee, A. S., MacLean, J. C., & Newton, D. A. (1994). Rapid Traction for Reduction of
 36 Cervical Spine Dislocation. *Journal of Bone and Joint Surgery: Britain*, 76(B), 352-
 37 356
 38
- 39 Magit D, Wolff A, Sutton K, Medvecky MJ. Arthrofibrosis of the knee. J Am Acad Orthop
 40 Surg. 2007 Nov;15(11):682-94

- 1 Marquez-Lara A, Padget W, Wall EJ, Parikh SN. Manipulation Under Anesthesia is Safe
2 and Effective for Management of Early Postoperative Knee Arthrofibrosis in
3 Adolescent Patients. *J Pediatr Orthop*. 2024;44(1):e84-e90
4
- 5 Maund E, Craig D, Suekarran S, Neilson A, Wright K, Brealey S, Dennis L, Goodchild L,
6 Hanchard N, Rangan A, Richardson G, Robertson J, McDaid C. Management of frozen
7 shoulder: a systematic review and cost-effectiveness analysis. *Health Technol Assess*.
8 2012;16(11):1-264
9
- 10 Maxwell, H. A., & Turner, P. G. (1994). Dislocation of the Austin Moore hemiarthroplasty:
11 is closed manipulation justified? *Journal of the Royal Colleges of Surgeons of*
12 *Edinburgh and Ireland*, 39(6), 370-371
13
- 14 Mohammed R, Syed S, Ahmed N. Manipulation under anesthesia for stiffness following
15 knee arthroplasty. *Ann R Coll Surg Engl*. 2009 Apr;91(3):220-3
16
- 17 Mullen JP, Hauer TM, Lau EN, Lin A. Adhesive Capsulitis of the Shoulder. *Arthroscopy*.
18 2025;41(7):2176-2178. doi:10.1016/j.arthro.2025.03.027
19
- 20 Namba RS, Inacio M. Early and late manipulation improve flexion after total knee
21 arthroplasty. *J Arthroplasty*. 2007 Sep;22(6 Suppl 2):58-61
22
- 23 Newman ET, Herschmiller TA, Attarian DE, Vail TP, Bolognesi MP, Wellman SS. Risk
24 Factors, Outcomes, and Timing of Manipulation Under Anesthesia After Total Knee
25 Arthroplasty. *J Arthroplasty*. 2018 Jan;33(1):245-249
26
- 27 Palmieri, N. F., & Smoyak, S. (2002). Chronic low back pain: a study of the effects of
28 manipulation under anesthesia. *Journal of Manipulative and Physiological*
29 *Therapeutics*, 25(8), E8-E17
30
- 31 Pivec R, Issa K, Kester M, Harwin SF, Mont MA. Long-term outcomes of MUA for
32 stiffness in primary TKA. *Knee Surg*. 2013 Dec;26(6):405-10
33
- 34 Quraishi NA, Johnston P, Bayer J, et al. Thawing the frozen shoulder. A randomised trial
35 comparing manipulation under anaesthesia with hydrodilatation. *J Bone Joint Surg Br*.
36 2007;89(9):1197-1200
37
- 38 Randsborg PH, Tajet J, Negård H, Røtterud JH. Manipulation under Anesthesia for
39 Stiffness of the Knee Joint after Total Knee Replacement. *Arthroplast Today*. 2020 Jun
40 28;6(3):470-474

- 1 Rangan A, Brealey SD, Keding A, Corbacho B, Northgraves M, Kottam L, Goodchild L,
2 Srikesavan C, Rex S, Charalambous CP, Hanchard N, Armstrong A, Brooksbank A,
3 Carr A, Cooper C, Dias JJ, Donnelly I, Hewitt C, Lamb SE, McDaid C, Richardson G,
4 Rodgers S, Sharp E, Spencer S, Torgerson D, Tøye F; UK FROST Study Group.
5 Management of adults with primary frozen shoulder in secondary care (UK FROST):
6 a multicentre, pragmatic, three-arm, superiority randomised clinical trial. *Lancet*. 2020
7 Oct 3;396(10256):977-989. doi: 10.1016/S0140-6736(20)31965-6
8
- 9 Rex SS, Kottam L, McDaid C, et al. Effectiveness of interventions for the management of
10 primary frozen shoulder : a systematic review of randomized trials. *Bone Jt Open*.
11 2021;2(9):773-784
12
- 13 Sala J, Jaroma A, Sund R, Huopio J, Kröger H, Sirola J. Manipulation under anesthesia
14 after total knee arthroplasty: a retrospective study of 145 patients. *Acta Orthop*.
15 2022;93:583-587. Published 2022 Jun 21. doi:10.2340/17453674.2022.3167
16
- 17 Salomon M, Pastore C, Maselli F, Di Bari M, Pellegrino R, Brindisino F. Manipulation
18 under Anesthesia versus Non-Surgical Treatment for Patients with Frozen Shoulder
19 Contracture Syndrome: A Systematic Review. *Int J Environ Res Public Health*.
20 2022;19(15):9715. Published 2022 Aug 7. doi:10.3390/ijerph19159715
21
- 22 Sheridan MA, Hannafin JA. Upper extremity: emphasis on frozen shoulder. *Orthop Clin*
23 *North Am*. 2006 Oct;37(4):531-9
24
- 25 Siehl, D., & Bradford, W. (1952). Manipulation of the low Back under General Anesthesia.
26 *Journal of the American Osteopathic Association*, 52(4), 239-242
27
- 28 Siehl, D., Olson, D. R., Ross, H. E., & Rockwood, E. E. (1971). Manipulation of the lumbar
29 spine with the patient under general anesthesia: evaluation by electromyography and
30 clinical-neurologic examination of its use for lumbar nerve root compression
31 syndrome. *Journal of the American Osteopathic Association*, 70(5), 433-440
32
- 33 Song C, Song C, Li C. Outcome of manipulation under anesthesia with or without intra-
34 articular steroid injection for treating frozen shoulder: A retrospective cohort study.
35 *Medicine (Baltimore)*. 2021;100(13):e23893
36
- 37 Speed C. Shoulder pain. In: *BMJ Clinical Evidence*. London, UK: BMJ Publishing Group;
38 February 2006
39
- 40 Spitler CA, Doty DH, Johnson MD, Nowotarski PJ, Kiner DW, Swafford RE, Jemison
41 DM. Manipulation Under Anesthesia as a Treatment of Posttraumatic Elbow Stiffness.
42 *J Orthop Trauma*. 2018 Aug;32(8):e304-e308

- 1 Tan V, Daluiski A, Simic P, Hotchkiss RN . Outcome of open release for post-traumatic
2 elbow stiffness. *J Trauma* 2006 Sep;6(13):673-8
3
- 4 Theodorides AA, Owen JM, Sayers AE, Woods DA. Factors affecting short- and long-
5 term outcomes of manipulation under anaesthesia in patients with adhesive capsulitis
6 of the shoulder. *Shoulder Elbow*. 2014 Oct;6(4):245-56
7
- 8 Tsai, S. W., & Chou, C. S. (2005). A case report of manipulation under anesthesia of
9 posttraumatic type II occipital-atlantoaxial rotatory subluxation in a 4-year-old girl.
10 *Journal of Manipulative and Physiological Therapeutics*, 28(5), 352-355
11
- 12 Thomas NP, Liu C, Varady N, Iban YC, Schwab PE, Chen AF. High Complication Rate
13 Associated With Arthroscopic Lysis of Adhesions Versus Manipulation Under
14 Anesthesia for Arthrofibrosis After Total Knee Arthroplasty. *J Am Acad Orthop Surg*.
15 2023 Feb 15;31(4):e216-e225. doi: 10.5435/JAAOS-D-22-00430. Epub 2022 Dec 21.
16 PMID: 36728979
17
- 18 Vastamäki H, Vastamäki M. Motion and pain relief remain 23 years after manipulation
19 under anesthesia for frozen shoulder. *Clin Orthop Relat Res*. 2013 Apr;471(4):1245-
20 50
21
- 22 Vezeridis PS, Goel DP, Shah AA, Sung SY, Warner JJ. Postarthroscopic arthrofibrosis of
23 the shoulder. *Sports Med Arthrosc*. 2010 Sep;18(3):198-206
24
- 25 W-Dahl A. Manipulation under anesthesia: to do or not to do, that is the question. *Acta*
26 *Orthop*. 2022;93:682-683. Published 2022 Jul 15. doi:10.2340/17453674.2022.4344
27
- 28 Wang JP, Huang TF, Hung SC, Ma HL, Wu JG, Chen TH. Comparison of idiopathic, post-
29 trauma and post-surgery frozen shoulder after manipulation under anesthesia. *Int*
30 *Orthop*. 2007 Jun;31(3):333-7. Epub 2006 Aug 23
31
- 32 West, D. T., Mathews, R. S., Miller, M. R., & Kent, G. M. (1999). Effective management
33 of spinal pain in one hundred seventy-seven patients evaluated for manipulation under
34 anesthesia. *Journal of Manipulative and Physiological Therapeutics*, 22(5), 299-308
35
- 36 Witvrouw E, Bellemans J, Victor J. Manipulation under anaesthesia versus low stretch
37 device in poor range of motion after TKA. *Knee Surg Sports Traumatol Arthrosc*. 2012
38 Aug 3
39
- 40 Woods DA, Loganathan K. Recurrence of frozen shoulder after manipulation under
41 anaesthetic (MUA): the results of repeating the MUA. *Bone Joint J*. 2017 Jun;99-
42 B(6):812-817

- 1 Xiong, X. H., Bean, A., Anthony, A., Inglis, G., & Walton, D. (1998). Manipulation for
- 2 cervical spinal dislocation under general anaesthesia: serial review for 4 years. *Spinal*
- 3 *Cord*, 36(1), 21-24